

Thin-layer chromatography of the peyote alkaloids*

The peyote cactus, *Lophophora williamsii* (Lem. ex SD.) Coult. (T.) (syn. *Anhalonium lewinii* Hennings), contains, besides the narcotic mescaline, a number of other bases, which are derivatives of phenylethylamine or tetrahydroisoquinoline^{1,2}. The use of peyote ("mescal buttons") by the natives of Mexico as a hallucinogenic drug has lately also spread to other countries. Until the recent paper by McLAUGHLIN AND PAUL³ apparently no thin-layer or paper chromatographic procedures for the rapid identification of *Lophophora* bases had been published.

In this note several thin-layer chromatographic systems suitable for the separation and identification of the peyote alkaloids are described.

Methods and materials

Thin-layer chromatography was carried out as described earlier⁴ on silica gel coated glass plates (20 × 20 cm, 0.25 mm layer) except that the coated plates were dried overnight at room temperature. For details regarding solvent systems, see Table I.

The base fraction from a peyote cactus (fresh wt. ca. 100 g–0.4 g alkaloids) was

TABLE I

R_F VALUES × 100 OF PEYOTE ALKALOIDS

Silica Gel G chromatoplates with the following solvent mixtures:

- (A) chloroform-ethanol-diethylamine (85:5:10 by vol.)
- (B) chloroform-ethanol-diethylamine (85:10:5)
- (C) chloroform-ethanol-conc. NH_3 (85:15:0.4)
- (D) chloroform-*n*-butanol-conc. NH_3 (50:50:2.5)
- (E) pyridine-conc. NH_3 (90:10)

Alkaloid	Solvent system					Colour ^a
	A	B	C	D	E	
<i>Phenolic</i>						
Anhalamine	11	20	—	—	40	purple
N-Methyltyramine	31	31	—	—	32	yellow
Tyramine	34	33	—	—	42	yellow
Anhalonidine	39	51	—	—	51	purple
Hordenine	51	56	—	—	60	yellow
Anhalidine	55	65	—	—	72	purple
Pellotine	63	70	—	—	69	purple
<i>Non-phenolic</i>						
N-Methylmescaline	—	—	22	20	25	yellow
Mescaline	—	—	24	31	36	brown
Anhalinine	—	—	30	41	48	yellow
O-Methylanhalonidine	—	—	33	45	50	yellow
Anhalonine	—	—	45	58	56	yellow
Lophophorine	—	—	68	80	72	blue gray
N-Acetylmescaline	—	—	82	95	68	pale brown

^a Colour with *o*-dianisidine reagent.

* Supported by the Swedish National Research Council. The technical assistance of Miss B. BURELL is appreciated.

isolated by chloroform extraction⁵. The evaporated chloroform extract was dissolved in 100 ml chloroform and passed through a 2×15 cm column of acid Celite (15 g Celite 545 and 4 ml 0.5 M H_3PO_4). The column was washed with 200 ml chloroform to remove non-basic compounds. The alkaloids were eluted with chloroform saturated with ammonia⁶. A solution of the alkaloids in methanol was applied to a column (1×20 cm) of Amberlite IRA 400 (OH) ion-exchange resin. The column was washed with 100 ml of 30 % aqueous methanol to yield the non-phenolic alkaloids. The phenolic alkaloid fraction was obtained by elution with 200 ml of a solution of 120 ml methanol, 60 ml water and 20 ml glacial acetic acid.

Alkaloids were located by the use of an *o*-dianisidine reagent (equal volumes of 0.5 % *o*-dianisidine in dilute HCl and 10 % NaNO_2 in water) or iodoplatinate reagent⁷.

Reference alkaloids were kindly supplied by Drs. A. BROSSI, Hoffman-La Roche Inc., and G. KAPADIA, Howard University, or isolated or synthesized according to known procedures (*cf.* ref. 1).

Results and discussion

The thin-layer chromatographic behaviour of the peyote alkaloids in several solvent systems and their colour reactions with the dianisidine reagent are recorded in Table I. This reagent produces a red colour with phenolic tetrahydroisoquinolines and a yellow or brown, fading colour with non-phenolic alkaloids.

Solvent system A was found to be most suitable for the separation of phenolic alkaloids and system D for non-phenolic alkaloids. With the exception of solvent system E, no system was found to resolve satisfactorily both phenolic and non-phenolic alkaloids.

Details of thin-layer and gas chromatographic separation of peyote alkaloids will be published at a later date.

Department of Pharmacognosy, Kungliga
Farmaceutiska Institutet, Kungstengatan 49,
Stockholm (Sweden)

JAN LUNDSTRÖM
STIG AGURELL

1 H. G. BOIT, *Ergebnisse der Alkaloidchemie bis 1960*, Akademie-Verlag, Berlin, 1961, pp. 13-24, 210-215.

2 A. HOFMANN, *Planta Med.*, 12 (1964) 341.

3 J. L. McLAUGHLIN AND A. G. PAUL, *Lloydia*, 29 (1966) 315.

4 S. AGURELL, *Acta Pharm. Suecica*, 2 (1965) 357.

5 E. LEETE, *J. Am. Chem. Soc.*, 88 (1966) 4218.

6 *Pharmacopea Nordica, Editio Suecica*, Vol. I, Apotekarsocietetens Förlag, Stockholm, 1964, pp. 74-75.

7 E. STAHL, *Dünnschicht-Chromatographie*, Springer-Verlag, Berlin, 1962, p. 506.

Received April 13th, 1967

J. Chromatog., 30 (1967) 271-272