

Psychedelic 5-Methoxy-*N,N*-Dimethyltryptamine: Metabolism, Pharmacokinetics, Drug Interactions, and Pharmacological Actions

Hong-Wu Shen¹, Xi-Ling Jiang¹, Jerrold C. Winter² and Ai-Ming Yu^{1,2,*}

¹Department of Pharmaceutical Sciences, School of Pharmacy and Pharmaceutical Sciences; ²Department of Pharmacology and Toxicology, School of Medicine and Biomedical Sciences, University at Buffalo, The State University of New York, Buffalo, NY 14260-1200, USA

Abstract: 5-Methoxy-*N,N*-dimethyltryptamine (5-MeO-DMT) belongs to a group of naturally-occurring psychoactive indolealkylamine drugs. It acts as a nonselective serotonin (5-HT) agonist and causes many physiological and behavioral changes. 5-MeO-DMT is *O*-demethylated by polymorphic cytochrome P450 2D6 (CYP2D6) to an active metabolite, bufotenine, while it is mainly inactivated through the deamination pathway mediated by monoamine oxidase A (MAO-A). 5-MeO-DMT is often used with MAO-A inhibitors such as harmaline. Concurrent use of harmaline reduces 5-MeO-DMT deamination metabolism and leads to a prolonged and increased exposure to the parent drug 5-MeO-DMT, as well as the active metabolite bufotenine. Harmaline, 5-MeO-DMT and bufotenine act agonistically on serotonergic systems and may result in hyperserotonergic effects or serotonin toxicity. Interestingly, CYP2D6 also has important contribution to harmaline metabolism, and CYP2D6 genetic polymorphism may cause considerable variability in the metabolism, pharmacokinetics and dynamics of harmaline and its interaction with 5-MeO-DMT. Therefore, this review summarizes recent findings on biotransformation, pharmacokinetics, and pharmacological actions of 5-MeO-DMT. In addition, the pharmacokinetic and pharmacodynamic drug-drug interactions between harmaline and 5-MeO-DMT, potential involvement of CYP2D6 pharmacogenetics, and risks of 5-MeO-DMT intoxication are discussed.

Keywords: 5-MeO-DMT, harmaline, metabolism, drug interaction, pharmacokinetics, pharmacogenetics, CYP2D6, serotonin toxicity.

INTRODUCTION

Indolealkylamine drugs consist of many antimigraine triptans (e.g. sumatriptan, naratriptan and almotriptan) and psychedelic substances of abuse (e.g., 5-methoxy-*N,N*-dimethyltryptamine or 5-MeO-DMT) [1]. 5-MeO-DMT was initially isolated from the bark of *Dictyoloma incanescens* D.C. [2]. It is a major active ingredient of South American *Virola* snuffs and *Ayahuasca* beverage [3-5]. 5-MeO-DMT also represents the active constituent of the venom of Colorado River *Bufo alvarius* toads, and it accounts for 15% of the dry weight of parotoid and tibial glands [6]. In addition, 5-MeO-DMT may be synthesized in human pineal and retina, and has been identified in human body fluids including urine, blood, and cerebrospinal fluid [7-11]. 5-MeO-DMT is regarded as an endogenous psychotoxin, and elevated concentrations of 5-MeO-DMT and its analogs in body fluids might be associated with psychotic disorders such as schizophrenic psychosis [12-19].

5-MeO-DMT is a potent, fast-acting hallucinogen with short duration in humans. Following different administration routes, e.g., inhalation (~6-20 mg), intravenous injection (~0.7-3.1 mg), sublingual or intranasal insufflation (~10 mg), and oral administration (~30 mg; with MAO inhibitor), 5-MeO-DMT produces psychedelic effects in human subjects [20]. 5-MeO-DMT also induces various physiological and behavioral changes in animal models [21-24]. 5-MeO-DMT has high affinity for the serotonin 5-HT_{1A} receptor, and it is 4- to 10-fold more potent than *N,N*-dimethyltryptamine (DMT) in human subjects [25, 26]. Unlike its chemically and pharmacologically related drugs, such as 5-hydroxy-*N,N*-dimethyltryptamine (bufotenine), DMT, 5-methoxy-*N,N*-diisopropyltryptamine (5-MeO-DiPT) and α -methyltryptamine (AMT) that all have been Schedule I controlled drugs for years, 5-MeO-DMT was not a federally controlled substance in United States until August 2009 when the Drug Enforcement Administration issued a notice intending to place 5-MeO-DMT into Schedule I of the Controlled Substances Act [20]. By contrast, 5-MeO-DMT had been a controlled substance in many European countries including the United Kingdom.

As a tryptamine derivative, 5-MeO-DMT is mainly inactivated through a deamination pathway mediated by monoamine oxidase A (MAO-A), and it is *O*-demethylated by cytochrome P450 2D6 (CYP2D6) enzyme to produce an active metabolite, bufotenine [27, 28], which binds to the 5-HT_{2A} receptor with much higher affinity than 5-MeO-DMT itself [29-31]. Concurrent use of 5-MeO-DMT with an MAO inhibitor (MAOI), or plant preparations (e.g., *ayahuasca*) or Syrian rue (*Peganum harmala*) seeds containing an MAOI (e.g., harmaline) [4, 32, 33], often leads to an enhanced and prolonged drug effect or more severe toxicity. Mechanistically, both MAOI and 5-MeO-DMT act agonistically on serotonergic systems that readily causes hyperserotonergic effects or serotonin toxicity [34-36]. In addition, MAOI increases exposure to the parent drug 5-MeO-DMT and the active metabolite bufotenine through the inhibition of deamination metabolism [37]. Such complex pharmacokinetic and pharmacodynamic interactions may even cause fatal toxicity [38, 39]. This review, therefore, aims to introduce the current understanding of 5-MeO-DMT biotransformation, pharmacokinetics and pharmacological actions, as well as drug-drug interactions (DDI) between 5-MeO-DMT and the MAOI, harmaline, and potential impact of CYP2D6 genetics.

PHARMACO/TOXICOLOGICAL EFFECTS AND DRUG ACTIONS OF 5-MEO-DMT

5-MeO-DMT is psychoactive in humans following inhalation of the vapor [20]. Human self-experiments have revealed that 5-MeO-DMT causes visionary and auditory changes, and distorts the perception of time. The effects start at 3-4 min, peak about 35-40 min, and end around 60-70 min after insufflation [3]. However, oral ingestion of 30-35 mg of 5-MeO-DMT produces either no psychoactive effect at all [40] or only one-third of the potency as that generated from intranasal or sublingual ingestion, or oral administration of *pharmahuasca* that contains both 5-MeO-DMT and the MAOI, harmaline [3, 26].

The toxicity of 5-MeO-DMT was firstly reported as a lethal syndrome called "staggers" in sheep after grazing on *Phalaris tuberosa*, a plant containing 5-MeO-DMT [41]. Studies with mouse, rat, sheep, and monkey models revealed remarkable ataxia, mydriasis, head nodding, tremor, convulsion and shivering after administration of 5-MeO-DMT. Sheep was found to be the most susceptible to 5-

*Address correspondence to this author at the Department of Pharmaceutical Sciences, School of Pharmacy and Pharmaceutical Sciences, University at Buffalo, The State University of New York, 541 Cooke Hall, Buffalo, NY 14260-1200, USA; Tel: 716-645-4817; Fax: 716-645-3693; E-mail: aimingyu@buffalo.edu

MeO-DMT, and exhibited stringy salivation, tachycardia, and respiratory failure when exposed to 1 mg/kg of 5-MeO-DMT. The LD₅₀ values of 5-MeO-DMT in mice ranged from 48 to 278 mg/kg for different administration routes [12, 21].

The 5-HT_{2A} receptor subtype has been shown to play a major role in the stimulus effects of indolealkylamine and phenethylamine hallucinogens, while the 5-HT_{2C} receptor acts as a modulator [42-44]. However, the 5-HT_{2A} receptor appears to be less important for the stimulus control of 5-MeO-DMT, and the potencies of drugs in substituting 5-MeO-DMT-induced stimulus effect are correlated well with their affinities with the 5-HT_{1A} receptor [29]. In addition, the discriminative stimuli induced by 5-MeO-DMT are attenuated by 5-HT_{1A} antagonists including TVX Q7821, WAY-100635 and pindolol [29, 45, 46], supporting the conclusion that 5-MeO-DMT induces stimulus control mainly via 5-HT_{1A} receptor. Nevertheless, a partial generalization of 5-MeO-DMT-induced stimulus by (-)-1-(2,5-dimethoxy-4-methylphenyl)-2-aminopropane [(+)-DOM] suggests that, besides the 5-HT_{1A} receptor, the 5-HT_{2A} receptor may also contribute to 5-MeO-DMT-mediated stimulus complex [45]. Interestingly, other structurally-related tryptamines including DMT and 5-MeO-DIPT also exhibit considerable binding affinities toward 5-HT_{1A} receptor, as well as partial or full agonistic activities against 5-HT_{2A} receptor [46-49].

Besides drug-induced discriminative stimulus control, 5-MeO-DMT provokes a variety of other behavioral effects in animal models, such as head shaking, forepaw treading, flat-body posture, straub tail, and hindlimb abduction [50-55] that are shared with many other hallucinogens including lysergic acid diethylamide (LSD), DOM and 1-(2,5-dimethoxy-4-iodophenyl)-2-aminopropane (DOI) [24, 56-58]. The similar effects of 5-MeO-DMT and 8-hydroxy-N,N-dipropyl-2-aminotetralin (8-OH-DPAT), a 5-HT_{1A} receptor agonist, on forepaw treading behavior in rats support the action of 5-MeO-DMT on postsynaptic 5-HT_{1A} receptor [51, 54]. In mice, 5-MeO-DMT induces head-twitch and head-weaving responses via the activation of 5-HT_{2A} and 5-HT_{1A} receptors, respectively [50, 59]. Among a series of methylated serotonin derivatives, 5-MeO-DMT has been identified as one of the most potent agents to induce "sham rage" responses in cats [12]. 5-MeO-DMT also decreases locomotor activity and investigatory behavior in rats that can be attenuated by WAY-100635, a selective 5-HT_{1A/7} antagonist but not by the 5-HT_{2A}-selective antagonist, M100907, suggesting the involvement of the 5-HT_{1A} receptor [55]. Pretreatment with MAOI significantly modifies the locomotor activity induced by 5-MeO-DMT to a biphasic effect in rats, which may involve the activation of 5-HT_{2A} receptor [60]. In addition, 5-MeO-DMT induces body temperature change in rats, causing hypothermia at low doses (0.5-1.0 mg/kg) and hyperthermia at high dose (3-10 mg/kg). The hyperthermic effect may be completely attenuated or even converted into hypothermia by the 5-HT_{2A} antagonist, ketanserin. Chronic treatment with MAOI nialamide not only diminishes the hypothermic effect but also attenuates the hyperthermic response [61, 62].

5-MeO-DMT indeed binds to the 5-HT_{1A} receptor subtype with much higher affinity (K_i , < 10 nM) than 5-HT₂ receptor (>1000 nM) [29]. Studies using rat brain synaptosomes [63] show that 5-MeO-DMT also inhibits 5-HT re-uptake with an IC₅₀ value comparable to other psychostimulants such as cocaine and methamphetamine, whereas it has little effect on dopamine re-uptake or the release of monoamine neurotransmitters. Furthermore, 5-MeO-DMT is the most potent tryptamine in stimulating G protein binding, with an EC₅₀ value around 100 nM, which is approximately 115% of the maximum activation of 5-HT itself [64]. In addition, 5-MeO-DMT and other endogenous hallucinogens (e.g., DMT and bufotenine) are thought to act as ligands for the recently discovered trace amine receptors that may be involved in sensory perception [65].

BIOTRANSFORMATION OF 5-MEO-DMT

Similar to tryptamine, DMT and bufotenine [66-69], 5-MeO-DMT primarily undergoes MAO-A-mediated deamination (Fig. (1)). Other metabolic pathways reported for 5-MeO-DMT include *O*-demethylation, *N*-demethylation and *N*-oxygenation. After intraperitoneal (i.p.) administration of ¹⁴C-labeled 5-MeO-DMT, 5-methoxyindoleacetic acid (5-MIAA) was identified as the major metabolite (54%) in rat urine, followed by 5-hydroxy-*N,N*-dimethyltryptamine glucuronide (23%), 5-hydroxyindoleacetic acid (5-HIAA, 14%), and bufotenine (9%) [70]. Because 5-HIAA cannot be transformed from 5-MIAA but from bufotenine, the 5-HIAA radioactivity may be attributed to consequential *O*-demethylation and deamination, suggesting that oxidative deamination and *O*-demethylation are the two major biotransformation pathways for 5-MeO-DMT in rats. 5-MeO-DMT was also found to be *N*-oxidized in rats [71-74]. In rat models, though the role of *N*-oxidation is uncertain, deaminated and *O*-demethylated metabolites were found in all studies. Our studies [27, 28, 37, 75] using human liver microsomes, hepatocytes and recombinant enzymes, as well as wild-type and CYP2D6-humanized mouse models, support the conclusion that 5-MeO-DMT is mainly inactivated through an MAO-A-catalyzed deamination pathway in humans although some 5-MeO-DMT is *O*-demethylated to bufotenine.

It is noteworthy that bufotenine, the *O*-demethylated product of 5-MeO-DMT, is also a hallucinogenic compound co-existing with 5-MeO-DMT in many plants and toad species [4-6, 76]. Bufotenine exhibits about 5- to 10-fold higher affinity to the 5-HT_{2A} receptor than 5-MeO-DMT *in vitro*, and about 3-fold higher potency than 5-MeO-DMT when they are present at similar level in brain [31, 77, 78]. Although the psychoactivity of bufotenine had been questioned due to its lower ability to cross blood-brain barrier (BBB), bufotenine does produce psychoactive effects in humans after intravenous injection or intranasal and sublingual administration [76, 79, 80]. Thus, the production of bufotenine from 5-MeO-DMT may be viewed as an activation process similar to the production of serotonin from 5-methoxytryptamine [81]. Indeed, severe toxicity or even lethality associated with the use of bufotenine has been documented [82, 83], and bufotenine has been a Schedule I controlled substance in United State for decades.

The production of bufotenine from 5-MeO-DMT is predominantly mediated by human CYP2D6 (Fig. (1)), which is well known for its genetic polymorphism and clinical importance [28, 84-87]. This biotransformation was initially discovered using recombinant CYP2D6 enzyme [27]. Our recent studies [37] indicate that 5-MeO-DMT *O*-demethylation is significantly correlated with bufuralol 1'-hydroxylation and CYP2D6 content in human liver microsomes when the deamination pathway is blocked by an MAO inhibitor. In addition, the CYP2D6.10 allelic isoform has much lower enzymatic activity in catalyzing 5-MeO-DMT *O*-demethylation than wild-type CYP2D6.1 enzyme, suggesting that subjects carrying the CYP2D6*10 allelic variant may produce less bufotenine from 5-MeO-DMT. Given the important role for CYP2D6 in 5-MeO-DMT *O*-demethylation activation, elucidation of the formation of bufotenine would be necessary to understand 5-MeO-DMT pharmacokinetics and risk of intoxication.

PHARMACOKINETICS OF 5-MEO-DMT

After i.p. administration, 5-MeO-DMT reaches the maximum drug concentration (C_{max}) at around 5-7 min and is then eliminated with a terminal half-life ($t_{1/2}$) of 12-19 min in mice [75]. The fast absorption and short $t_{1/2}$ are also true for 5-MeO-DMT in rats [72]. 5-MeO-DMT is predominantly eliminated through MAO-A-mediated metabolism, as supported by the low urinary recovery and biliary excretion of the parent compound [70, 71] and an over 4-fold increase in systemic exposure to 5-MeO-DMT when co-administered with an MAOI [37]. 5-MeO-DMT also shows a rela-




Fig. (1). 5-MeO-DMT biotransformation and the metabolic interactions with harmaline. O-demethylation of 5-MeO-DMT by CYP2D6 produces an active metabolite bufotenine. Both 5-MeO-DMT and bufotenine are readily deaminated by MAO-A to indoleacetic acid derivatives. The MAOI harmaline, which is inactivated by CYP2D6, blocks the deamination metabolism of 5-MeO-DMT and bufotenine.

tively high oil/water partition coefficient (3.30) [88], suggesting that 5-MeO-DMT may easily penetrate various lipoprotein barriers including the BBB. Indeed, 5-MeO-DMT significantly accumulates in many organs (e.g., liver, kidney and brain) in different animal models (e.g., mouse, rat and rabbit) [72, 73, 89]. The brain concentration of 5-MeO-DMT is about 1.7-fold higher than that in blood at 45 min after i.p. administration [72], and the drug is widely distributed in different rat brain regions including cortex, thalamus, hippocampus, basal ganglia, medulla, pons and cerebellum [90]. Our unpublished results also support the idea that 5-MeO-DMT is readily distributed and accumulated in mouse cortex, hippocampus, hypothalamus, and striatum after i.p. administration.

As mentioned above, monitoring bufotenine formation may provide improved understanding of the complex pharmacological and toxicological effects of 5-MeO-DMT because bufotenine is the active metabolite of 5-MeO-DMT, and has a higher binding affinity for the 5-HT_{2A} receptor than 5-MeO-DMT does. Bufotenine is also rapidly eliminated from the body. In healthy volunteers receiving an intravenous infusion of ¹⁴C-labeled bufotenine, nearly all the radioactivity is recovered in the first 12 hr urine sample [91]. Furthermore, only 1-6% of total recovered radioactivity is identified as unchanged bufotenine, while the deaminated metabolite 5-HIAA accounts for 68-74% of radioactivity. Compared to 5-MeO-DMT, bufotenine has a poor partition coefficient and shows a low penetration across the BBB but a high accumulation in rat lungs after subcutaneous injection [68, 88].

Bufotenine is readily detected in mouse blood samples, following the administration of a lower dose of 5-MeO-DMT (2 mg/kg, i.p.) [37, 75]. Bufotenine reaches the C_{max} at around 13 min, and is eliminated with an apparent half-life of about 25 min. The systemic exposure (AUC) to bufotenine appears to be less than 10% of that of the parent 5-MeO-DMT in mice [37, 75]. Considering the low systemic exposure to bufotenine produced from 5-MeO-DMT and the low BBB penetration for bufotenine, the peripheral bufotenine produced from 5-MeO-DMT may not be able to cross the BBB and thus maintain sufficient concentration in brain to exert neurotoxicity. In addition, our unpublished findings indicate that bufotenine may not be extensively produced from 5-MeO-DMT within mouse brain. Therefore, it is likely that only peripherally-generated bufotenine contributes to the apparent activity of 5-MeO-DMT.

DRUG INTERACTIONS BETWEEN 5-MEO-DMT AND MAOI HARMALINE

DDI may be arisen when subjects are exposed to polypharmacy [36, 92-95]. At the pharmacokinetic level, a perpetrator drug alters the absorption, disposition, metabolism, or excretion of the victim drug, which may be translated into a significant change in drug efficacy and/or toxicity. The metabolic DDI is often observed because metabolism represents the major route of drug elimination. At the pharmacodynamic level, concurrent drugs both act on the common targets, leading to synergistic or antagonistic responses. It should be noted that some drugs may interact at both pharmacokinetic and pharmacodynamic levels, and lead to severe or even fatal toxicity. This may be true for the concomitant use of 5-MeO-DMT and harmaline or another MAOI. By inhibiting MAO-A-mediated 5-HT degradation, MAOI itself promotes serotonergic transmission Fig. (2). Furthermore, harmaline is also a 5-HT agonist [96, 97] that could potentiate serotonergic actions of 5-MeO-DMT. In addition, harmaline reduces 5-MeO-DMT deamination metabolism, leading to an increased and prolonged exposure to 5-MeO-DMT, as well as the psychoactive metabolite bufotenine that depends upon CYP2D6 status. Indeed, cases of severe and lethal intoxication due to the combined use of a tryptamine (e.g., 5-MeO-DMT or 5-MeO-DiPT) and an MAOI drug (e.g., harmaline) have been reported [38, 39, 98, 99].

Pharmacokinetic and Pharmacodynamic Interactions

The pharmacokinetics of 5-MeO-DMT can be significantly altered by a co-administered MAOI. Co-incubation of 5-MeO-DMT with harmaline in human hepatocytes completely blocks the depletion of 5-MeO-DMT, and reduces the *in vitro* intrinsic clearance (CL_{int}) by over 24-fold [37]. In rats, pretreatment with iproniazid significantly increases 5-MeO-DMT levels in urine, blood, and other tissues [71, 72]. In mice, pretreatment with harmaline decreases the clearance of 5-MeO-DMT by 4.4-fold, leading to a 4.4-fold higher systemic exposure to 5-MeO-DMT Fig. (3) [37]. The concomitant administration of harmaline not only affects the pharmacokinetics of 5-MeO-DMT, but also influences formation of the active metabolite bufotenine Fig. (3). When deamination metabolism is inhibited, more 5-MeO-DMT is diverted to other metabolic pathways including O-demethylation with increased production of bufotenine. In human CYP2D6 extensive metabolizer hepatocytes,




Fig. (2). Hyperserotonergic effects may be induced when harmaline blocks 5-HT degradation, and the concurrent 5-MeO-DMT activates 5-HT receptors within the synaptic cleft, beside their pharmacokinetic interactions.




Fig. (3). Harmaline (5 mg/kg, i.p.) co-administered with 5-MeO-DMT (2 mg/kg, i.p.) markedly alters blood concentrations of the parent drug 5-MeO-DMT (A), the active metabolite bufotenine (B), and the ratio of their systemic exposures (C) in wild-type and/or Tg-CYP2D6 mouse models [37].

bufotenine formation is significantly increased when 5-MeO-DMT is co-incubated with harmaline [37]. Since bufotenine is also mainly inactivated through the MAO-A-mediated deamination pathway [91], MAOI may not only increase bufotenine production but also reduce its elimination. As a result, the C_{max} and AUC of bufotenine are increased to 2.6- and 6-fold, respectively, in mice after harmaline pretreatment [37].

Besides the impact on 5-MeO-DMT pharmacokinetics, harmaline acts on serotonergic systems and exhibits a variety of neuropharmacological activities. Harmaline is one of the most potent inhibitors of MAO-A [100], and it may interact with 5-HT, dopamine, gamma-amino-butyric acid (GABA), and N-methyl-D-aspartate (NMDA) receptors [101-105]. The pharmacokinetic and dynamic interactions of harmaline and 5-MeO-DMT are manifested by a remarkable change in drug responses. For instance, pretreatment with harmaline and clorgyline in rats significantly modifies 5-MeO-DMT-induced locomotor activity to a biphasic effect, of which the late hyperactivity is attenuated by MDL 11939, a selective 5-HT_{2A} antagonist, suggesting the important contribution of the 5-HT_{2A} receptor to the hyperactivity induced by combined use of 5-MeO-DMT and MAOI [60]. Studies in our laboratory indicate that harmaline enhances the hyperthermic effect of 5-MeO-DMT and potentiates stimulus control in mice (unpublished data). Anecdotal reports by human self-experimenters suggest an enhancement of the psychedelic effects of 5-MeO-DMT when combined with harmaline or harmine [3, 40]. For instance, 10 mg of 5-MeO-DMT administered intranasally or sublingually causes significant visionary response, whereas the same dose of oral 5-MeO-DMT does not show any effects in humans, which is probably due to the extensive first-pass metabolism by MAO-A. However, when used with harmaline, the same oral dose 5-MeO-DMT (10 mg) shows nearly equal intensity of psychedelic effects as 10 mg of 5-MeO-DMT itself dosed intranasally or sublingually [3]. These findings indicate that 5-MeO-DMT pharmacological and toxicological effects are generally potentiated by a concurrent MAOI because of pharmacokinetic and dynamic interactions.

The Impact of CYP2D6 Genetic Polymorphism

CYP2D6 is an important P450 enzyme that has more than 90 allelic variants, leading to considerable interindividual variability in

metabolism of some CYP2D6 substrate drugs including therapeutic agents and drugs of abuse [1, 106-108]. Because 5-MeO-DMT *O*-demethylation is dependent upon CYP2D6 [27], the impact of CYP2D6 phenotype/genotype on the production of bufotenine from 5-MeO-DMT has been examined [37]. The blockage of MAO activity results in a strong correlation between 5-MeO-DMT *O*-demethylation and CYP2D6 activity in human liver microsomes. As expected, the CYP2D6.1, CYP2D6.2 and CYP2D6.10 allelic isoforms shows variable catalytic activities in producing bufotenine from 5-MeO-DMT, and human CYP2D6 poor hepatocytes lacking CYP2D6 activity do not produce any bufotenine. In addition, CYP2D6-humanized (Tg-CYP2D6) mice [109] show a significantly higher systemic exposure (AUC) to the active metabolite bufotenine than wild-type mice treated i.p. with 20 mg/kg of 5-MeO-DMT. It should be noted that the Tg-CYP2D6 and wild-type control mice are useful animal models in delineating the impact of CYP2D6 on drug metabolism and pharmacokinetics [28, 110, 111]. Therefore, it is expected that subjects with regular or increased CYP2D6 activity would be exposed to both the substrate drug 5-MeO-DMT and the active metabolite bufotenine, and thus might have more complex drug effects [37].




Fig. (4). Studies using wild-type and Tg-CYP2D6 mouse model demonstrate that CYP2D6 status has significant impact on harmaline pharmacokinetics (A) and harmaline-induced hypothermia (B) [113].

Harmaline co-administered with 5-MeO-DMT is also metabolized by CYP2D6 [112], and the effects of CYP2D6 status on harmaline metabolism, pharmacokinetics and pharmacodynamics have been defined separately [113]. Depletion of harmaline is 2.5 times slower in human CYP2D6 poor metabolizer hepatocytes than that in CYP2D6 extensive metabolizer hepatocytes. Further studies using Tg-CYP2D6 and wild-type mouse models reveal that wild-type mice have a longer and higher exposure to harmaline, as well as a more severe hypothermia Fig. (4). In addition, wild-type mice lacking CYP2D6 activity are more sensitive to harmaline in marble-

burying test [113]. Therefore, a lower exposure to harmaline in Tg-CYP2D6 mice might counteract the effect of CYP2D6 on 5-MeO-DMT *O*-demethylation. Indeed, levels of bufotenine formed from 5-MeO-DMT do not differ significantly between wild-type and Tg-CYP2D6 mice after co-administration of harmaline [37], which may be due to the low dose (2 mg/kg, i.p.) of 5-MeO-DMT used in the study and the significant contribution of mouse P450s to 5-MeO-DMT *O*-demethylation [27, 28]. Nevertheless, systemic exposure to bufotenine is about 24% of that to 5-MeO-DMT in Tg-CYP2D6 mice, whereas it is 15% in wild-type mice, indicating the impact of CYP2D6 status on harmaline-5-MeO-DMT pharmacokinetic drug interaction.

CONCLUSIONS

5-MeO-DMT represents a natural psychoactive tryptamine indolealkylamine drug of abuse. As a fast-acting drug, 5-MeO-DMT induces many physiological and behavioral changes in humans and animal models, such as visionary and auditory distortion, hyperthermia, head-twitch, and stimulus control, which involve the actions of 5-HT receptors. It is readily inactivated through the MAO-A-mediated deamination pathway, while a small portion is transformed to the active metabolite bufotenine by polymorphic CYP2D6. Receptor binding studies have revealed that 5-MeO-DMT is more selective for the 5-HT_{1A} receptor, while bufotenine mainly acts as a 5-HT_{2A} agonist. The apparent non-selectivity of 5-MeO-DMT may be attributed, at least in part, to bufotenine produced from 5-MeO-DMT, which is dependent on CYP2D6 enzymatic activity.

5-MeO-DMT is often co-abused with an MAOI such as harmaline to enhance hallucinations. There are two levels of interactions between harmaline and 5-MeO-DMT, pharmacokinetic and pharmacodynamic. When deamination metabolism is inhibited by harmaline, the systemic and cerebral exposure to 5-MeO-DMT, as well as to the metabolite bufotenine, is sharply elevated and prolonged. Meanwhile, harmaline and 5-MeO-DMT both act agonistically on the serotonergic systems. As a result, coadministration of harmaline potentiates 5-MeO-DMT drug responses and sometimes leads to severe or fatal serotonin toxicity in animal models. Several cases of intoxication or even death have also been reported in humans associated with the abuse of 5-MeO-DMT and harmaline. In addition, depending upon the dose combination, an influence of CYP2D6 genotype/phenotype on harmaline-5-MeO-DMT DDI may be seen, despite the fact that the CYP2D6 enzyme inactivates harmaline whereas it activates 5-MeO-DMT. Therefore, CYP2D6 may serve as a marker for bufotenine production during pharmacokinetic interaction between 5-MeO-DMT and harmaline.

ACKNOWLEDGEMENT

This project was supported by Award Number R01DA021172 from the National Institute On Drug Abuse, National Institutes of Health (NIH).

ABBREVIATIONS

5-MeO-DMT	=	5-methoxy- <i>N,N</i> -dimethyltryptamine
Bufotenine	=	5-hydroxy- <i>N,N</i> -dimethyltryptamine
CYP2D6	=	Cytochrome P450 2D6
MAO-A	=	Monoamine oxidase A
MAOI	=	Monoamine oxidase inhibitor
DDI	=	Drug-drug interaction
Serotonin	=	5-HT
DMT	=	<i>N,N</i> -dimethyltryptamine
5-MeO-DiPT	=	5-methoxy- <i>N,N</i> -diisopropyltryptamine
BBB	=	Blood-brain barrier
Tg-CYP2D6	=	CYP2D6-humanized

REFERENCES

- [1] Yu, A.M. Indolealkylamines: biotransformations and potential drug-drug interactions. *Aaps. J.*, **2008**, *10*, 242-253.
- [2] Pachter, I.J.; Zacharias, D.E.; Ribeiro, O. Indole alkaloids of acer saccharinum (the Silver Maple), *Dictyoloma incanescens*, *Piptadenia colubrina*, and *Mimosa hostilis*. *J. Org. Chem.*, **1959**, *24*, 1285-1287.
- [3] Ott, J. Pharmepena-Psychonautics: Human intranasal, sublingual and oral pharmacology of 5-methoxy-N,N-dimethyl-tryptamine. *J. Psychoactive Drugs*, **2001**, *33*, 403-407.
- [4] McKenna, D.J. Clinical investigations of the therapeutic potential of ayahuasca: rationale and regulatory challenges. *Pharmacol. Ther.*, **2004**, *102*, 111-129.
- [5] Gambelunghe, C.; Aroni, K.; Rossi, R.; Moretti, L.; Bacci, M. Identification of N,N-dimethyltryptamine and beta-carbolines in psychotropic ayahuasca beverage. *Biomed. Chromatogr.*, **2008**, *22*, 1056-1059.
- [6] Weil, A.T.; Davis, W. Bufo alvarius: a potent hallucinogen of animal origin. *J. Ethnopharmacol.*, **1994**, *41*, 1-8.
- [7] Narasimhachari, N.; Heller, B.; Spaide, J.; Haskovec, L.; Meltzer, H.; Strahilevitz, M.; Himwich, H.E. N,N-dimethylated indoleamines in blood. *Biol. Psychiatry*, **1971**, *3*, 21-23.
- [8] Oon, M.C.; Murray, R.M.; Rodnight, R.; Murphy, M.P.; Birley, J.L. Factors affecting the urinary excretion of endogenously formed dimethyltryptamine in normal human subjects. *Psychopharmacology (Berl.)*, **1977**, *54*, 171-175.
- [9] Smythies, J.R.; Morin, R.D.; Brown, G.B. Identification of dimethyltryptamine and O-methylbufotenine in human cerebrospinal fluid by combined gas chromatography/mass spectrometry. *Biol. Psychiatry*, **1979**, *14*, 549-556.
- [10] Sitaram, B.R.; Blackman, G.L.; McLeod, W.R.; Vaughan, G.N. The ion-pair extraction, purification, and liquid chromatographic analysis of indolealkylamines in human urine. *Anal. Biochem.*, **1983**, *128*, 11-20.
- [11] Forsstrom, T.; Tuominen, J.; Karkkainen, J. Determination of potentially hallucinogenic N-dimethylated indoleamines in human urine by HPLC/ESI-MS-MS. *Scand. J. Clin. Lab. Invest.*, **2001**, *61*, 547-556.
- [12] Benington, F.; Morin, R.D.; Clark, L.C., Jr. 5-methoxy-N, N-dimethyltryptamine, a possible endogenous psychotoxin. *Ala. J. Med. Sci.*, **1965**, *2*, 397-403.
- [13] Nishimura, T.; Gjessing, L.R. Failure to detect 3, 4-dimethoxyphenylethylamine and bufotenine in the urine from a case of periodic catatonia. *Nature*, **1965**, *206*, 963-964.
- [14] Narasimhachari, N.; Heller, B.; Spaide, J.; Haskovec, L.; Fujimori, M.; Tabushi, K.; Himwich, H.E. Urinary studies of schizophrenics and controls. *Biol. Psychiatry*, **1971**, *3*, 9-20.
- [15] Gillin, J.C.; Wyatt, R.J. Evidence for and against the involvement of N,N-dimethyl-tryptamine (DMT) and 5-methoxy-N,N-dimethyltryptamine (5-MeO-DMT) in schizophrenia. *Psychopharmacol. Bull.*, **1976**, *12*, 12-13.
- [16] Angrist, B.; Gershon, S.; Sathananthan, G.; Walker, R.W.; Lopez-Ramos, B.; Mandel, L.R.; Vandenheuvel, W.J. Dimethyltryptamine levels in blood of schizophrenic patients and control subjects. *Psychopharmacology (Berl.)*, **1976**, *47*, 29-32.
- [17] Takeda, N.; Ikeda, R.; Ohba, K.; Kondo, M. Bufotenine reconsidered as a diagnostic indicator of psychiatric disorders. *Neuroreport*, **1995**, *6*, 2378-2380.
- [18] Cichon, S.; Nothen, M.M.; Rietschel, M.; Propping, P. Pharmacogenetics of schizophrenia. *Am. J. Med. Genet.*, **2000**, *97*, 98-106.
- [19] Emanuele, E.; Colombo, R.; Martinelli, V.; Brondino, N.; Marini, M.; Boso, M.; Barale, F.; Politi, P. Elevated urine levels of bufotenine in patients with autistic spectrum disorders and schizophrenia. *Neuro Endocrinol. Lett.*, **2010**, *31*, 117-121.
- [20] DEA-2009-0008. Placement of 5-Methoxy-N,N-Dimethyltryptamine Into Schedule I of the Controlled Substances Act. <http://edocket.access.gpo.gov/2009/E9-20204.htm> (Accessed Sep 23, 2010).
- [21] Gillin, J.C.; Tinklenberg, J.; Stoff, D.M.; Stillman, R.; Shortlidge, J.S.; Wyatt, R.J. 5-Methoxy-N,N-dimethyltryptamine: behavioral and toxicological effects in animals. *Biol. Psychiatry*, **1976**, *11*, 355-358.
- [22] Glennon, R.A. Do classical hallucinogens act as 5-HT2 agonists or antagonists? *Neuropsychopharmacology*, **1990**, *3*, 509-517.
- [23] Duvvuri, V.; Risbrough, V.B.; Kaye, W.H.; Geyer, M.A. 5-HT1A receptor activation is necessary for 5-MeODMT-dependent potentiation of feeding inhibition. *Pharmacol. Biochem. Behav.*, **2009**, *93*, 349-353.
- [24] Fantegrossi, W.E.; Murnane, K.S.; Reissig, C.J. The behavioral pharmacology of hallucinogens. *Biochem. Pharmacol.*, **2008**, *75*, 17-33.
- [25] McKenna, D.J.; Towers, G.H. Biochemistry and pharmacology of tryptamines and beta-carbolines. A minireview. *J. Psychoactive Drugs*, **1984**, *16*, 347-358.
- [26] Ott, J. Pharmahuasca: human pharmacology of oral DMT plus harmine. *J. Psychoactive Drugs*, **1999**, *31*, 171-177.
- [27] Yu, A.M.; Idle, J.R.; Herraiz, T.; Kupfer, A.; Gonzalez, F.J. Screening for endogenous substrates reveals that CYP2D6 is a 5-methoxyindolethylamine O-demethylase. *Pharmacogenetics*, **2003**, *13*, 307-319.
- [28] Yu, A.M.; Idle, J.R.; Gonzalez, F.J. Polymorphic cytochrome P450 2D6: humanized mouse model and endogenous substrates. *Drug. Metab. Rev.*, **2004**, *36*, 243-277.
- [29] Spencer, D.G., Jr.; Glaser, T.; Traber, J. Serotonin receptor subtype mediation of the interoceptive discriminative stimuli induced by 5-methoxy-N,N-dimethyltryptamine. *Psychopharmacology (Berl.)*, **1987**, *93*, 158-166.
- [30] Glennon, R.A.; Dukat, M.; el-Bermawy, M.; Law, H.; De los Angeles, J.; Teitler, M.; King, A.; Herrick-Davis, K. Influence of amine substituents on 5-HT2A versus 5-HT2C binding of phenylalkyl- and indolylalkylamines. *J. Med. Chem.*, **1994**, *37*, 1929-1935.
- [31] Roth, B.L.; Choudhary, M.S.; Khan, N.; Uluer, A.Z. High-affinity agonist binding is not sufficient for agonist efficacy at 5-hydroxytryptamine2A receptors: evidence in favor of a modified ternary complex model. *J. Pharmacol. Exp. Ther.*, **1997**, *280*, 576-583.
- [32] Lamchouri, F.; Settaf, A.; Cherrah, Y.; El Hamidi, M.; Tligui, N.; Lyoussi, B.; Hassar, M. Experimental toxicity of Peganum harmala seeds. *Ann. Pharm. Fr.*, **2002**, *60*, 123-129.
- [33] Herraiz, T.; Gonzalez, D.; Ancin-Azpilicueta, C.; Aran, V.J.; Guillen, H. beta-Carboline alkaloids in Peganum harmala and inhibition of human monoamine oxidase (MAO). *Food. Chem. Toxicol.*, **2010**, *48*, 839-845.
- [34] Boyer, E.W.; Shannon, M. The serotonin syndrome. *N. Engl. J. Med.*, **2005**, *352*, 1112-1120.
- [35] Sun-Edelstein, C.; Tepper, S.J.; Shapiro, R.E. Drug-induced serotonin syndrome: a review. *Expert Opin. Drug. Saf.*, **2008**, *7*, 587-596.
- [36] Gillman, P.K. A review of serotonin toxicity data: implications for the mechanisms of antidepressant drug action. *Biol. Psychiatry*, **2006**, *59*, 1046-1051.
- [37] Shen, H.W.; Wu, C.; Jiang, X.L.; Yu, A.M. Effects of monoamine oxidase inhibitor and cytochrome P450 2D6 status on 5-methoxy-N,N-dimethyltryptamine metabolism and pharmacokinetics. *Biochem. Pharmacol.*, **2010**, *80*, 122-128.
- [38] Brush, D.E.; Bird, S.B.; Boyer, E.W. Monoamine oxidase inhibitor poisoning resulting from Internet misinformation on illicit substances. *J. Toxicol. Clin. Toxicol.*, **2004**, *42*, 191-195.
- [39] Sklerov, J.; Levine, B.; Moore, K.A.; King, T.; Fowler, D. A fatal intoxication following the ingestion of 5-methoxy-N,N-dimethyltryptamine in an ayahuasca preparation. *J. Anal. Toxicol.*, **2005**, *29*, 838-841.
- [40] Shulgin, A.T.; Shulgin, A. *TIKAL The Continuation*, Transform Press, Berkeley, CA, **1997**.
- [41] Gallagher, C.H.; Koch, J.H.; Moore, R.M.; Steel, J.D. Toxicity of Phalaris Tuberosa for sheep. *Nature*, **1964**, *204*, 542-545.
- [42] Glennon, R.A.; Titeler, M.; McKenney, J.D. Evidence for 5-HT2 involvement in the mechanism of action of hallucinogenic agents. *Life. Sci.*, **1984**, *35*, 2505-2511.
- [43] Darmani, N.A.; Martin, B.R.; Pandey, U.; Glennon, R.A. Do functional relationships exist between 5-HT1A and 5-HT2 receptors? *Pharmacol. Biochem. Behav.*, **1990**, *36*, 901-906.
- [44] Winter, J.C.; Fiorella, D.J.; Timineri, D.M.; Filipink, R.A.; Helsley, S.E.; Rabin, R.A. Serotonergic receptor subtypes and hallucinogen-

- induced stimulus control. *Pharmacol. Biochem. Behav.*, **1999**, *64*, 283-293.
- [45] Winter, J.C.; Filipink, R.A.; Timineri, D.; Helsley, S.E.; Rabin, R.A. The paradox of 5-methoxy-N,N-dimethyltryptamine: an indoleamine hallucinogen that induces stimulus control via 5-HT1A receptors. *Pharmacol. Biochem. Behav.*, **2000**, *65*, 75-82.
- [46] Winter, J.C. Hallucinogens as discriminative stimuli in animals: LSD, phenethylamines, and tryptamines. *Psychopharmacology (Berl)*, **2009**, *203*, 251-263.
- [47] Sadzot, B.; Baraban, J.M.; Glennon, R.A.; Lyon, R.A.; Leonhardt, S.; Jan, C.R.; Titeler, M. Hallucinogenic drug interactions at human brain 5-HT2 receptors: implications for treating LSD-induced hallucinogenesis. *Psychopharmacology (Berl)*, **1989**, *98*, 495-499.
- [48] Fantegrossi, W.E.; Harrington, A.W.; Kiessel, C.L.; Eckler, J.R.; Rabin, R.A.; Winter, J.C.; Coop, A.; Rice, K.C.; Woods, J.H. Hallucinogen-like actions of 5-methoxy-N,N-diisopropyltryptamine in mice and rats. *Pharmacol. Biochem. Behav.*, **2006**, *83*, 122-129.
- [49] Rabin, R.A.; Regina, M.; Doat, M.; Winter, J.C. 5-HT2A receptor-stimulated phosphoinositide hydrolysis in the stimulus effects of hallucinogens. *Pharmacol. Biochem. Behav.*, **2002**, *72*, 29-37.
- [50] Matsumoto, K.; Mizowaki, M.; Takayama, H.; Sakai, S.; Aimi, N.; Watanabe, H. Suppressive effect of mitragynine on the 5-methoxy-N,N-dimethyltryptamine-induced head-twitch response in mice. *Pharmacol. Biochem. Behav.*, **1997**, *57*, 319-323.
- [51] Tricklebank, M.D.; Forler, C.; Middlemiss, D.N.; Fozard, J.R. Subtypes of the 5-HT receptor mediating the behavioural responses to 5-methoxy-N,N-dimethyltryptamine in the rat. *Eur. J. Pharmacol.*, **1985**, *117*, 15-24.
- [52] Eison, A.S.; Wright, R.N. 5-HT1A and 5-HT2 receptors mediate discrete behaviors in the Mongolian gerbil. *Pharmacol. Biochem. Behav.*, **1992**, *43*, 131-137.
- [53] Cancela, L.M.; Volosin, M.; Molina, V.A. Gangliosides attenuate stress-induced changes on body weight, motor activity and on the behavioral response to 5-methoxy-N,N-dimethyltryptamine. *Brain. Res. Bull.*, **1996**, *40*, 105-110.
- [54] Sanchez, C.; Arnt, J.; Moltzen, E. Assessment of relative efficacies of 5-HT1A receptor ligands by means of *in vivo* animal models. *Eur. J. Pharmacol.*, **1996**, *315*, 245-254.
- [55] Krebs-Thomson, K.; Ruiz, E.M.; Masten, V.; Buell, M.; Geyer, M.A. The roles of 5-HT1A and 5-HT2 receptors in the effects of 5-MeO-DMT on locomotor activity and prepulse inhibition in rats. *Psychopharmacology (Berl)*, **2006**, *189*, 319-329.
- [56] Knapp, D.J.; Sim-Selley, L.J.; Breese, G.R.; Overstreet, D.H. Selective breeding of 5-HT(1A) receptor-mediated responses: application to emotion and receptor action. *Pharmacol. Biochem. Behav.*, **2000**, *67*, 701-708.
- [57] Fox, M.A.; Andrews, A.M.; Wendland, J.R.; Lesch, K.P.; Holmes, A.; Murphy, D.L. A pharmacological analysis of mice with a targeted disruption of the serotonin transporter. *Psychopharmacology (Berl)*, **2007**, *195*, 147-166.
- [58] Fantegrossi, W.E.; Harrington, A.W.; Eckler, J.R.; Arshad, S.; Rabin, R.A.; Winter, J.C.; Coop, A.; Rice, K.C.; Woods, J.H. Hallucinogen-like actions of 2,5-dimethoxy-4-(n)-propylthiophenethylamine (2C-T-7) in mice and rats. *Psychopharmacology (Berl)*, **2005**, *181*, 496-503.
- [59] Matsumoto, K.; Morishige, R.; Murakami, Y.; Tohda, M.; Takayama, H.; Sakakibara, I.; Watanabe, H. Suppressive effects of isorhynchophylline on 5-HT2A receptor function in the brain: behavioural and electrophysiological studies. *Eur. J. Pharmacol.*, **2005**, *517*, 191-199.
- [60] Halberstadt, A.L.; Buell, M.R.; Masten, V.L.; Risbrough, V.B.; Geyer, M.A. Modification of the effects of 5-methoxy-N,N-dimethyltryptamine on exploratory behavior in rats by monoamine oxidase inhibitors. *Psychopharmacology (Berl)*, **2008**, *201*, 55-66.
- [61] Gudelsky, G.A.; Koenig, J.I.; Jackman, H.; Meltzer, H.Y. Suppression of the hypo- and hyperthermic responses to 5-HT agonists following the repeated administration of monoamine oxidase inhibitors. *Psychopharmacology (Berl)*, **1986**, *90*, 403-407.
- [62] Gudelsky, G.A.; Koenig, J.I.; Meltzer, H.Y. Thermoregulatory responses to serotonin (5-HT) receptor stimulation in the rat. Evidence for opposing roles of 5-HT2 and 5-HT1A receptors. *Neuropharmacology*, **1986**, *25*, 1307-1313.
- [63] Nagai, F.; Nonaka, R.; Satoh Hisashi Kamimura, K. The effects of non-medically used psychoactive drugs on monoamine neurotransmission in rat brain. *Eur. J. Pharmacol.*, **2007**, *559*, 132-137.
- [64] Nonaka, R.; Nagai, F.; Ogata, A.; Satoh, K. *In vitro* screening of psychoactive drugs by [(35)S]GTPgammaS binding in rat brain membranes. *Biol. Pharm. Bull.*, **2007**, *30*, 2328-2333.
- [65] Wallach, J.V. Endogenous hallucinogens as ligands of the trace amine receptors: a possible role in sensory perception. *Med. Hypotheses*, **2009**, *72*, 91-94.
- [66] Suzuki, O.; Katsumata, Y.; Oya, M. Characterization of eight biogenic indoleamines as substrates for type A and type B monoamine oxidase. *Biochem. Pharmacol.*, **1981**, *30*, 1353-1358.
- [67] Raynaud, F.; Pevet, P. 5-Methoxytryptamine is metabolized by monoamine oxidase A in the pineal gland and plasma of golden hamsters. *Neurosci. Lett.*, **1991**, *123*, 172-174.
- [68] Fuller, R.W.; Snoddy, H.D.; Perry, K.W. Tissue distribution, metabolism and effects of bufotenine administered to rats. *Neuropharmacology*, **1995**, *34*, 799-804.
- [69] Yu, A.M.; Granvil, C.P.; Haining, R.L.; Krausz, K.W.; Corchero, J.; Kupfer, A.; Idle, J.R.; Gonzalez, F.J. The relative contribution of monoamine oxidase and cytochrome p450 isozymes to the metabolic deamination of the trace amine tryptamine. *J. Pharmacol. Exp. Ther.*, **2003**, *304*, 539-546.
- [70] Agurell, S.; Holmstedt, B.; Lindgren, J.E. Metabolism of 5-methoxy-N,N dimethyltryptamine- 14 C in the rat. *Biochem. Pharmacol.*, **1969**, *18*, 2771-2781.
- [71] Sitaram, B.R.; Lockett, L.; Blackman, G.L.; McLeod, W.R. Urinary excretion of 5-methoxy-N,N-dimethyltryptamine, N,N-dimethyltryptamine and their N-oxides in the rat. *Biochem. Pharmacol.*, **1987**, *36*, 2235-2237.
- [72] Sitaram, B.R.; Lockett, L.; Talomsin, R.; Blackman, G.L.; McLeod, W.R. *In vivo* metabolism of 5-methoxy-N,N-dimethyltryptamine and N,N-dimethyltryptamine in the rat. *Biochem. Pharmacol.*, **1987**, *36*, 1509-1512.
- [73] Sitaram, B.R.; McLeod, W.R. Observations on the metabolism of the psychotomimetic indolealkylamines: implications for future clinical studies. *Biol. Psychiatry*, **1990**, *28*, 841-848.
- [74] Sitaram, B.R.; Talomsin, R.; Blackman, G.L.; McLeod, W.R. Study of metabolism of psychotomimetic indolealkylamines by rat tissue extracts using liquid chromatography. *Biochem. Pharmacol.*, **1987**, *36*, 1503-1508.
- [75] Shen, H.W.; Jiang, X.L.; Yu, A.M. Development of a LC-MS/MS method to analyze 5-methoxy-N,N-dimethyltryptamine and bufotenine: application to pharmacokinetic study. *Bioanalysis*, **2009**, *1*, 87-95.
- [76] Ott, J. Pharamnopo-psychonautics: human intranasal, sublingual, intrarectal, pulmonary and oral pharmacology of bufotenine. *J. Psychoactive Drugs*, **2001**, *33*, 273-281.
- [77] McBride, M.C. Bufotenine: toward an understanding of possible psychoactive mechanisms. *J. Psychoactive Drugs*, **2000**, *32*, 321-331.
- [78] Vogel, W.H.; Evans, B.D. Structure-activity-relationships of certain hallucinogenic substances based on brain levels. *Life Sci.*, **1977**, *20*, 1629-1635.
- [79] Fabing, H.D.; Hawkins, J.R. Intravenous bufotenine injection in the human being. *Science*, **1956**, *123*, 886-887.
- [80] McLeod, W.R.; Sitaram, B.R. Bufotenine reconsidered. *Acta. Psychiatr. Scand.*, **1985**, *72*, 447-450.
- [81] Yu, A.M.; Idle, J.R.; Byrd, L.G.; Krausz, K.W.; Kupfer, A.; Gonzalez, F.J. Regeneration of serotonin from 5-methoxytryptamine by polymorphic human CYP2D6. *Pharmacogenetics*, **2003**, *13*, 173-181.
- [82] Chilton, W.S.; Bigwood, J.; Jensen, R.E. Psilocin, bufotenine and serotonin: historical and biosynthetic observations. *J. Psychedelic Drugs*, **1979**, *11*, 61-69.
- [83] Kostakis, C.; Byard, R.W. Sudden death associated with intravenous injection of toad extract. *Forensic Sci. Int.*, **2009**, *188*, e1-5.
- [84] Zanger, U.M.; Raimundo, S.; Eichelbaum, M. Cytochrome P450 2D6: overview and update on pharmacology, genetics, biochemistry. *Naunyn Schmiedeberg's Arch. Pharmacol.*, **2004**, *369*, 23-37.
- [85] Looers, H.M.; van der Weide, J. Implementation of CYP2D6 genotyping in psychiatry. *Expert Opin. Drug Metab. Toxicol.*, **2009**, *5*, 1065-1077.

- [86] Dorado, P.; Penas-Lledo, E.M.; Llerena, A. CYP2D6 polymorphism: implications for antipsychotic drug response, schizophrenia and personality traits. *Pharmacogenomics*, **2007**, *8*, 1597-1608.
- [87] De Gregori, M.; Allegri, M.; De Gregori, S.; Garbin, G.; Tinelli, C.; Regazzi, M.; Govoni, S.; Ranzani, G.N. How and why to screen for CYP2D6 interindividual variability in patients under pharmacological treatments. *Curr. Drug Metab.*, **2010**, *11*, 276-282.
- [88] Migliaccio, G.P.; Shieh, T.L.; Byrn, S.R.; Hathaway, B.A.; Nichols, D.E. Comparison of solution conformational preferences for the hallucinogens bufotenin and psilocin using 360-MHz proton NMR spectroscopy. *J. Med. Chem.*, **1981**, *24*, 206-209.
- [89] Berger, G.; Maziere, M.; Marazano, C.; Comar, D. Carbon 11 labeling of the psychoactive drug o-methyl-bufotenine and its distribution in the animal organism. *Eur. J. Nucl. Med.*, **1978**, *3*, 101-104.
- [90] Barker, S.A.; Littlefield-Chabaud, M.A.; David, C. Distribution of the hallucinogens N,N-dimethyltryptamine and 5-methoxy-N,N-dimethyltryptamine in rat brain following intraperitoneal injection: application of a new solid-phase extraction LC-APCI-MS-MS-isotope dilution method. *J. Chromatogr. B. Biomed. Sci. Appl.*, **2001**, *751*, 37-47.
- [91] Sanders-Bush, E.; Oates, J.A.; Bush, M.T. Metabolism of bufotenine-2'-14C in human volunteers. *Life Sci.*, **1976**, *19*, 1407-1411.
- [92] Boobis, A.; Watelet, J.B.; Whomsley, R.; Benedetti, M.S.; Demoly, P.; Tipton, K. Drug interactions. *Drug Metab. Rev.*, **2009**, *41*, 486-527.
- [93] Yu, A.M. Role of microRNAs in the regulation of drug metabolism and disposition. *Expert Opin. Drug Metab. Toxicol.*, **2009**, *5*, 1513-1528.
- [94] He, S.M.; Yang, A.K.; Li, X.T.; Du, Y.M.; Zhou, S.F. Effects of herbal products on the metabolism and transport of anticancer agents. *Expert Opin. Drug Metab. Toxicol.*, **2010**, *6*, 1195-1213.
- [95] Ulbricht, C.; Chao, W.; Costa, D.; Rusie-Seamon, E.; Weissner, W.; Woods, J. Clinical evidence of herb-drug interactions: a systematic review by the natural standard research collaboration. *Curr. Drug Metab.*, **2008**, *9*, 1063-1120.
- [96] Abdel-Fattah, A.F.; Matsumoto, K.; Gammaz, H.A.; Watanabe, H. Hypothermic effect of harmala alkaloid in rats: involvement of serotonergic mechanism. *Pharmacol. Biochem. Behav.*, **1995**, *52*, 421-426.
- [97] Glennon, R.A.; Dukat, M.; Grella, B.; Hong, S.; Costantino, L.; Teitler, M.; Smith, C.; Egan, C.; Davis, K.; Mattson, M.V. Binding of beta-carbolines and related agents at serotonin (5-HT₂) and 5-HT_{1A}), dopamine (D₂) and benzodiazepine receptors. *Drug Alcohol Depend.*, **2000**, *60*, 121-132.
- [98] Tanaka, E.; Kamata, T.; Katagi, M.; Tsuchihashi, H.; Honda, K. A fatal poisoning with 5-methoxy-N,N-diisopropyltryptamine, foxy. *Forensic Sci. Int.*, **2006**, *163*, 152-154.
- [99] Bjornstad, K.; Hulten, P.; Beck, O.; Helander, A. Bioanalytical and clinical evaluation of 103 suspected cases of intoxications with psychoactive plant materials. *Clin. Toxicol. (Phila)*, **2009**, *47*, 566-572.
- [100] Kim, H.; Sablin, S.O.; Ramsay, R.R. Inhibition of monoamine oxidase A by beta-carboline derivatives. *Arch. Biochem. Biophys.*, **1997**, *337*, 137-142.
- [101] Monsef, H.R.; Ghobadi, A.; Iranshahi, M.; Abdollahi, M. Antinociceptive effects of Peganum harmala L. alkaloid extract on mouse formalin test. *J. Pharm. Pharm. Sci.*, **2004**, *7*, 65-69.
- [102] Hilber, P.; Chapillon, P. Effects of harmaline on anxiety-related behavior in mice. *Physiol. Behav.*, **2005**, *86*, 164-167.
- [103] Moura, D.J.; Rorig, C.; Vieira, D.L.; Henriques, J.A.; Roesler, R.; Saffi, J.; Boeira, J.M. Effects of beta-carboline alkaloids on the object recognition task in mice. *Life Sci.*, **2006**, *79*, 2099-2104.
- [104] Hutchinson, D.; Ho, V.; Dodd, M.; Dawson, H.N.; Zumwalt, A.C.; Schmitt, D.; Colton, C.A. Quantitative measurement of postural sway in mouse models of human neurodegenerative disease. *Neuroscience*, **2007**, *148*, 825-832.
- [105] Paterson, N.E.; Malekiani, S.A.; Foreman, M.M.; Olivier, B.; Hanania, T. Pharmacological characterization of harmaline-induced tremor activity in mice. *Eur. J. Pharmacol.*, **2009**, *616*, 73-80.
- [106] Zhou, S.F. Polymorphism of human cytochrome P450 2D6 and its clinical significance: part II. *Clin. Pharmacokinet.*, **2009**, *48*, 761-804.
- [107] Zhou, S.F. Polymorphism of human cytochrome P450 2D6 and its clinical significance: Part I. *Clin. Pharmacokinet.*, **2009**, *48*, 689-723.
- [108] Narimatsu, S.; Yonemoto, R.; Saito, K.; Takaya, K.; Kumamoto, T.; Ishikawa, T.; Asanuma, M.; Funada, M.; Kiryu, K.; Naito, S.; Yoshida, Y.; Yamamoto, S.; Hanioka, N. Oxidative metabolism of 5-methoxy-N,N-diisopropyltryptamine (Foxy) by human liver microsomes and recombinant cytochrome P450 enzymes. *Biochem. Pharmacol.*, **2006**, *71*, 1377-1385.
- [109] Corchero, J.; Granvil, C.P.; Akiyama, T.E.; Hayhurst, G.P.; Pimpale, S.; Feigenbaum, L.; Idle, J.R.; Gonzalez, F.J. The CYP2D6 humanized mouse: effect of the human CYP2D6 transgene and HNF4alpha on the disposition of debrisoquine in the mouse. *Mol. Pharmacol.*, **2001**, *60*, 1260-1267.
- [110] Gonzalez, F.J.; Yu, A.M. Cytochrome P450 and xenobiotic receptor humanized mice. *Annu. Rev. Pharmacol. Toxicol.*, **2006**, *46*, 41-64.
- [111] Lin, J.H. Applications and limitations of genetically modified mouse models in drug discovery and development. *Curr. Drug. Metab.*, **2008**, *9*, 419-438.
- [112] Yu, A.M.; Idle, J.R.; Krausz, K.W.; Kupfer, A.; Gonzalez, F.J. Contribution of individual cytochrome P450 isozymes to the O-demethylation of the psychotropic beta-carboline alkaloids harmaline and harmine. *J. Pharmacol. Exp. Ther.*, **2003**, *305*, 315-322.
- [113] Wu, C.; Jiang, X.L.; Shen, H.W.; Yu, A.M. Effects of CYP2D6 status on harmaline metabolism, pharmacokinetics and pharmacodynamics, and a pharmacogenetics-based pharmacokinetic model. *Biochem. Pharmacol.*, **2009**, *78*, 617-624.