

cervical spine radiography prior to intubation. When radiographic resources can only provide films of limited diagnostic power, and the clinical setting demands rapid intervention, resuscitation must proceed with in line cervical spine stabilization.

The maintenance of the airway in all patients with head injury is well known to be a vital first aid measure. It is probable that in flight difficulties with airway management during initial transfer resulted in relative hypercapnia, exacerbating the trauma related increase in intracranial pressure⁶, suggested by the history of a deteriorating conscious level and dilated left pupil. Prompt intervention with intubation and hyperventilation rapidly improved the clinical picture. We speculate that rapid normalization of the left pupil following airway control indicates reduction in brain oedema.

The Triservice Anaesthetic Apparatus³ was modified to incorporate a single Oxford Miniature Vaporiser⁷ factory calibrated to deliver isoflurane.

It is concluded that early neurosurgical intervention is possible in forward field dressing stations and we stress the importance of airway maintenance in such cases. Early correction of hypercapnia is probably crucial to outcome.

Although the eventual outcome was disappointing there is no doubt that the history and physical signs justified early craniotomy.

References

- 1 Teasdale G, Jennett B. Assessment of coma and impaired consciousness. A practical scale. *Lancet* 1974;ii:81-4
- 2 Sellick BA. Cricoid pressure to control regurgitation of stomach contents during induction of anaesthesia. *Lancet* 1961;ii:404-6
- 3 Houghton IT. The triservice anaesthetic apparatus. *Anaesthesia* 1981;36:1094-109
- 4 Jones RM. Clinical comparison of inhaled anaesthetic agents. *Br J Anaesthes* 1984;56S:57-69
- 5 American college of surgeons. *The advanced trauma life support student manual*. Chicago: American college of surgeons, 1989
- 6 Campkin TV, Turner JM. *Neurosurgical anaesthesia and intensive care*, 2nd edn. London: Butterworths, 1986
- 7 Parkhouse J. Clinical performance of the OMV inhaler. *Anaesthesia* 1966;21:498

(Accepted 5 November 1991)

Chronic nutmeg psychosis

N Brenner MRCPsych **O S Frank** MRCP FRCPsych
E Knight MB BS *Gordon Hospital,*
Bloomburg Street, London SW1

Keywords: nutmeg; chronic psychosis; hyponatraemia; convulsions

The nutmeg tree (*Myristica fragrans* Houtt. Family Myristicaceae) is found in the East Indies. The dried fruit yields the spice nutmeg. The first documented case of poisoning was in 1576 and used as a hallucinogen since the Crusades¹.

In the more recent literature there have been several reports of acute psychosis caused by nutmeg². However to date there have been no reported cases of chronic nutmeg abuse and its sequelae. Here we report such a case.

Case report

A 25-year-old Caucasian man of no fixed address, was admitted into hospital on a Section 136 of the Mental Health Act. He was dirty, dishevelled and very disturbed. He was unable to give a history, at that time he was overtly psychotic, showing thought disorder, perseveration of speech and appeared to be responding intermittently to auditory hallucinations. He thought he was about to die. He required antipsychotic medication (haloperidol). All routine blood tests were normal and urine drug screen was negative. Brain scan was normal.

It was discovered that his mental state fluctuated with his ingestion of large quantities of nutmeg, two to six packets at a time (120-650 mg). After taking this amount of nutmeg he would become agitated, disruptive, complain of 'uncomfortable feelings' in his head, a sense of impending death, dry tongue polydipsia and vomiting. Even though he was given large amounts of antipsychotic he remained intermittently agitated.

He was also noted to be polydipsic, often drinking up to five litres of fluid per hour. On at least one occasion he drank his own urine. Blood chemistry confirmed water overload

hyponatraemia (plasma sodium 115 mmol/l, urine osmolality 50 mmol/l; plasma osmolality 242 mmol/l). After a nutmeg binge these periods of polydipsia would last 1 or 2 days if water was not restricted. At one point he suffered a grand mal seizure probably related to water overload.

More history was obtained from his mother, whom he had not seen for 10 years. His parents had divorced when he was 5 years old, his father was a heavy drinker, violent with a possible mental illness. He was a 'loner' as a child, and worked for Customs and Excise after leaving school aged 16 years. His mother and sister noted a personality change about then and it was at this time that he started abusing nutmeg. The family lost touch with him soon afterwards. He had been admitted briefly to hospital on two occasions in the past. On both occasions he responded to an antipsychotic but nutmeg abuse was noted during his second admission.

His personal hygiene was always very poor, he responded to a behavioural programme and antipsychotic medication slowly. He continued to binge on nutmeg bought for him by other patients. He was never nutmeg free for more than 2 weeks.

Discussion

The mind altering experience (a high) is usually achieved by the ingestion of 5-15 g of nutmeg (one to three whole nutmegs or two teaspoons of grated nutmeg⁷). Toxic overdose can be seen with as little as 5 g⁴.

The effect starts within 6 h of ingestion. Overdose resembles an anticholinergic episode, with cutaneous flushing, tachycardia, decreased salivation, fever and CNS excitation. Classically nutmeg produces a miosis rather than a mydriasis but this is an unreliable sign⁸. Shock, coma and death have been reported in severe cases^{9,10} but usually complete recovery occurs within 24 h, treatment is mainly supportive.

Although myristicin is one of the major components¹¹, it is still unclear if it is the psychoactive element of nutmeg. Synthetic myristicin does not always produce hallucination¹². The structural similarities of the metabolized components of nutmeg to amphetamine-like compounds could be partly responsible for the activity of nutmeg¹³.

This case is complicated by the possible underlying mental illness. As nutmeg is so readily available repeated toxic states have led to episodes of polydipsia. The prognosis is poor for this man, with little motivation to control his

abuse of nutmeg even while under close supervision. He awaited a place in a behavioural unit.

References

- 1 Cushny AR. Nutmeg poisoning. *Proc R Soc Med* 1908;3:39-44
- 2 Painter JC, Shanor SP, Winek CL. Nutmeg poisoning: a case report. *Clin Toxicol* 1971;4:1-4
- 3 Payne RB. Nutmeg intoxication. *N Engl J Med* 1963;269:36-8
- 4 Green RC. Nutmeg poisoning. *JAMA* 1959;171:1342
- 5 Reekie JS. Nutmeg poisoning. *JAMA* 1909;52:62
- 6 Hammond PW. Nutmeg poisoning. *BMJ* 1906;ii:778
- 7 Shafran I. Nutmeg toxicology. *N Engl J Med* 1976;294:849

- 8 Ahmed A, Thompson HS. Nutmeg mydriasis. *JAMA* 1975;234:3
- 9 Faguet RA, Rowland KF. Spice cabinet intoxication. *Am J Psychiatry* 1978;135:860
- 10 Dodge WT. Nutmeg poisoning. *Med Rec* 1887;32:624
- 11 Suligin AT. Possible implication of myristicin as a psychotropic substance. *Nature* 1966;210:380-4
- 12 Framsworth NR. Nutmeg poisoning. *Am J Psychiatry* 1979;136:858-9
- 13 Weil AT. The structure of myristicin. *Economic Botany* 1965;19:194

(Accepted 12 November 1991)

Some recent books

Biotechnology Current Progress. P N Cheremisinoff, L M Ferrante, eds. (pp 266, SFr. 150) ISBN 0-87762-776-2, Pennsylvania: Technomic Publishing, 1991

Choices in Health Care: A Report by the Government Committee on Choices in Health Care, the Netherlands 1992 (pp 160) ISBN 90 346 2840 X, Rijswijk: Ministry of Welfare, 1992

Comparative Molecular Carcinogenesis: Progress in Clinical and Biological Research, Vol. 376. A J P Klein-Szanto, M W Anderson, J C Barrett, T J Slaga, eds. (pp 385) ISBN 0-471-56205-X, New York: Wiley-Liss, Inc, 1992

Delusions and Hallucinations in Old Age. C Katona, R Levy, eds. (pp 248, £12.50) ISBN 0 902241 47 8, London: The Royal College of Psychiatrists, 1992

Diseases of the Spinal Cord. E Critchley and A Eisen, eds. (pp 453, DM 236.00) ISBN 3-540-19684-6, London: Springer-Verlag, 1992

Down Syndrome: Advances in Medical Care. I T Lott, E E McCoy, eds. (pp 196) ISBN 0-471-56181-9, New York: Wiley-Liss, 1992

Gender Dysphoria, Vol. 1, No. 1. C Mate-Kole, ed. (pp 77) ISSN 0057-4689, London: Whiting & Birch Ltd, 1991

Guanidino Compounds in Biology and Medicine (The Third International Symposium on Guanidino Compounds in Biology and Medicine). P P De Deyn, B Marescau, Stalon and I A Qureshi, eds. (pp 480) ISBN 0 86196 330 X, London: John Libbey & Co Ltd, 1992

How Psychiatrists Look at Aging (Mental Health Library Series). G H Pollock, ed. (pp 244, £32.50) ISBN 0-8236-2364-5, Madison: International Universities Press, Inc, 1992

Innovative Competition in Medicine: A Schumpeterian Analysis of the Pharmaceutical Industry and the NHS. G Teeling Smith, ed. (pp 111, £7.50) London: Office of Health Economics, 1992

Interviewing the Sexually Abused Child: Investigation of Suspected Abuse. D P H Jones (pp 67, £5.50) ISBN 0 902241 46 X, London: The Royal College of Psychiatrists, 1992

Management Training for Psychiatrists. D Bhugra, A Burns, eds (pp 260, £15.00) ISBN 0 902241 43 5, London: The Royal College of Psychiatrists, 1992

Melanie Klein: From Theory to Reality. O Weinger (pp 210, £16.95) ISBN 1 85575 011 2, London: Karnac Books, 1992

Molecular Biology of Atherosclerosis (Proceedings of the 57th European Atherosclerosis Society Meeting). M J Halpern, ed. (pp 662) ISBN 0 86196 3547, London: John Libbey & Co Ltd, 1992

Molecular Genetic Medicine, Vol. 2. T Friedmann, ed. (pp 233, \$55.00) ISBN 0-12-4620-7, San Diego: Academic Press Inc, 1992

Obesity in Europe 91 (Proceedings of the 3rd European Congress on Obesity). G Ailhaud, B Guy-Grand, M Lafontan and D Ricquier (pp 463) ISBN 0 86196 273 7, London: John Libbey & Co Ltd, 1992

Ovarian Cancer, Vol 2: Biology, Diagnosis and Management. F Sharp, W P Mason, W Creasman, eds (pp 453, £65.00) ISBN-0-412-45190-5, London: Chapman and Hall Medical, 1992

Promoting Health and Preventing Disease. D J Scherl, J Noren, M Osterweis (pp 212, \$20.00) ISBN 1-879694-04-2, Washington: Association of Academic Health Centers, 1992

Psychosocial Aspects of Genetic Counselling. G Evers-Kiebooms, J P Fryns, J J Cassiman, H Van de Berghe, eds (pp 203, £95.00) ISBN 0-471-56185-1, New York: Wiley-Liss, 1992

Quality Counts: Achieving Quality in Social Care Services. D Kelly and B Warr, eds. (pp 230) ISBN 1-87-1177-21-9, London: Whiting & Birch, 1992

Receptor Subunits and Complexes. A Burgen, E A Barnard, eds. (pp 468, £60.00) ISBN 0-521-36612-7, Cambridge: Cambridge University Press, 1992

Regulating British Medicine: The General Medical Council. M Stacey (pp 310, £15.95) ISBN 0 471 93189 6, Chichester: John Wiley & Sons Ltd, 1992

Reproductive Biology of Invertebrates, Vol. IV, Part B. K G Adiyodi, R G Adiyodi, eds. (pp 527) ISBN 0 471 922714, Chichester: John Wiley & Sons, 1990

Risk Markers for Oral Diseases, Vol. 3: Periodontal Diseases. N W Johnson, ed. (pp 464, £60.00) ISBN 0 521 38566 0, New York: Cambridge University Press, 1991

Spinal Cord Dysfunction, Vol. III, Functional Stimulation. L S Illis, ed. (pp 360, £50.00) ISBN 0 19 2619195, Oxford: Oxford University Press, 1991

Spinal Manipulation. J F Bourdillon, E A Day, M R Bookhout, (pp 381, £35.00) ISBN 0 7506 0576 6, Oxford: Butterworth-Heinemann, 1992

Szycher's Dictionary of Biomaterials and Medical Devices. M Szycher (pp 264, SF. 187.00) ISBN 0-87762-882-3, Pennsylvania: Technomic, 1992

Textbook of Biochemistry with Clinical Correlations, 3rd edn. T M Devlin, ed. (pp 1185, £55.95) ISBN 0 47151 3482, New York: Wiley-Liss, 1992

The Amygdala: Neurobiological Aspects of Emotion, Memory, and Mental Dysfunction. J P Aggleton, ed. (pp 615, £125.00) ISBN 0-471-56129-0, New York: Wiley-Liss, 1992

The Harvard Lectures. A Freud (pp 142, £10.95) ISBN 1 85575 030 9, London: Karnac-Books, 1992

The NHS Transformed. I Holliday (pp 117, £4.99) ISBN 1-897626-00-2, Manchester: Baseline Book Co, 1992

The Royal College of Nursing Manual of Family Health (pp 692, £25.00) ISBN 0-356-20051-5, London: Little, Brown & Co, 1992