



Cannabinoids and Pain: Sites and Mechanisms of Action

Katarzyna Starowicz*, David P. Finn^{†,1}

*Institute of Pharmacology, Polish Academy of Sciences, Laboratory of Pain Pathophysiology, Krakow, Poland

[†]Pharmacology and Therapeutics, School of Medicine, Galway Neuroscience Centre and Centre for Pain Research, NCBES, National University of Ireland, Galway, Ireland

¹Corresponding author: e-mail address: david.finn@nuigalway.ie

Contents

1. Introduction	438
2. Anatomical Localization of the Endocannabinoid System Throughout the Pain Pathway	445
3. Supraspinal Sites and Mechanisms of Action	449
3.1 Evidence From Acute Pain Models	449
3.2 Evidence From Inflammatory Pain Models	452
3.3 Evidence From Neuropathic Pain Models	454
4. Spinal Sites and Mechanisms of Action	455
4.1 Evidence From Acute Pain Models	455
4.2 Evidence From Inflammatory Pain Models	455
4.3 Evidence From Neuropathic Pain Models	456
5. Peripheral Sites and Mechanisms of Action	458
5.1 Evidence From Acute Pain Models	458
5.2 Evidence From Inflammatory Pain Models	459
5.3 Evidence From Neuropathic Pain Models	459
6. Conclusion	461
Conflict of Interest	462
Acknowledgments	462
References	462
Further Reading	475

Abstract

The endocannabinoid system, consisting of the cannabinoid₁ receptor (CB₁R) and cannabinoid₂ receptor (CB₂R), endogenous cannabinoid ligands (endocannabinoids), and metabolizing enzymes, is present throughout the pain pathways. Endocannabinoids, phytocannabinoids, and synthetic cannabinoid receptor agonists have antinociceptive effects in animal models of acute, inflammatory, and neuropathic pain. CB₁R and CB₂R located at peripheral, spinal, or supraspinal sites are important targets mediating these antinociceptive effects. The mechanisms underlying the analgesic effects of

cannabinoids likely include inhibition of presynaptic neurotransmitter and neuropeptide release, modulation of postsynaptic neuronal excitability, activation of the descending inhibitory pain pathway, and reductions in neuroinflammatory signaling. Strategies to dissociate the psychoactive effects of cannabinoids from their analgesic effects have focused on peripherally restricted CB₁R agonists, CB₂R agonists, inhibitors of endocannabinoid catabolism or uptake, and modulation of other non-CB₁R/non-CB₂R targets of cannabinoids including TRPV1, GPR55, and PPARs. The large body of preclinical evidence in support of cannabinoids as potential analgesic agents is supported by clinical studies demonstrating their efficacy across a variety of pain disorders.

ABBREVIATIONS

2-AG 2-arachidonoyl glycerol

AEA anandamide

BLA basolateral nucleus of the amygdala

CB₁R cannabinoid receptor type 1

CB₂R cannabinoid receptor type 2

CCI chronic constriction injury

CeA central nucleus of the amygdala

CFA complete Freund's adjuvant

CGRP calcitonin gene-related peptide

DRG dorsal root ganglia

FAAH fatty acid amide hydrolase

GPR55 G protein-coupled receptor 55

i.p. intraperitoneal

i.t. intrathecal

MAGL monoacylglycerol lipase

PAG periaqueductal gray

PEA N-palmitoylethanolamide

PPARs peroxisome proliferator-activated receptors

RVM rostral ventromedial medulla

SNI spared nerve injury

SNL spinal nerve ligation

TRPV1 transient receptor potential subfamily V member 1

Δ⁹-THC Δ⁹-tetrahydrocannabinol



1. INTRODUCTION

Cannabis sativa has been used for medicinal purposes, including relief of pain, for thousands of years (Grinspoon & Bakalar, 1993). The isolation and identification of the principal psychoactive constituent of cannabis,

Δ^9 -tetrahydrocannabinol (Δ^9 -THC), in the 1960s (Mechoulam & Gaoni, 1967), sparked a search for its mechanism of action which in turn led to the discovery of two cannabinoid receptors, the cannabinoid₁ receptor (CB₁R) (Devane, Dysarz, Johnson, Melvin, & Howlett, 1988; Matsuda, Lolait, Brownstein, Young, & Bonner, 1990) and cannabinoid₂ receptor (CB₂R) (Munro, Thomas, & Abu-Shaar, 1993). Endogenous ligands (endocannabinoids) which exert their effects upon binding to these cannabinoid receptors were also discovered, the two best characterized being arachidonoyl ethanolamide (anandamide, AEA) (Devane et al., 1992) and 2-arachidonoyl glycerol (2-AG) (Mechoulam et al., 1995; Sugiura et al., 1995). The receptors, endocannabinoids, transport proteins, and enzymes that synthesize or degrade the endocannabinoids together comprise the endocannabinoid system. A large body of preclinical and clinical research indicates that this lipid signaling system modulates a broad range of physiological processes and behaviors including, but not limited to, pain, mood, appetite, emesis, neuronal activity, memory, immunity, cell development and cell fate, and the cardiovascular system. In particular, the antinociceptive effects of cannabinoids and endocannabinoid signaling have received a lot of attention over the past 30 years, with thousands of peer-reviewed publications reporting antinociceptive/analgesic effects in preclinical and clinical studies and elucidating the sites and mechanisms of action. The impact of this research has started to be seen in clinical practice with the introduction of the Δ^9 -THC/Cannabidiol buccal spray nabiximols (Sativex[®]) for the adjunctive treatment of neuropathic pain in multiple sclerosis patients and severe cancer pain in Canada, and with many US states and countries around the world relaxing their laws to allow patients to use cannabis or cannabinoids for a range of conditions including chronic pain. The present review will focus primarily on the evidence from preclinical studies utilizing animal models of acute, inflammatory, and neuropathic pain with an emphasis on the sites and mechanisms underlying cannabinoid-mediated antinociception. For excellent recent reviews and meta-analyses of clinical studies in this area, please see Barnes (2006), Boychuk, Goddard, Mauro, and Orellana (2015), Canadian Agency for Drugs and Technologies in Health (2016), Iskedjian, Bereza, Gordon, Piwko, and Einarson (2007), Lynch and Ware (2015), McCormick et al. (2017), Russo (2016), Vermersch (2011), and Whiting et al. (2015).

With regards preclinical studies in rodents, both genetic and pharmacological (Table 1) approaches have been used to demonstrate and understand

Table 1 Compounds Referred to in the Text and Used to Elucidate the Role of Cannabinoids and the Endocannabinoid System in Pain Modulation

Pharmacological Substance	IUPAC Name	Target
THC	(-)-(6 <i>aR</i> ,10 <i>aR</i>)-6,6,9-Trimethyl-3-pentyl-6 <i>a</i> ,7,8,10 <i>a</i> -tetrahydro-6 <i>H</i> -benzo[<i>c</i>]chromen-1-ol; Δ^9 -tetrahydrocannabinol	The main psychotropic constituent of cannabis, CB ₁ /CB ₂ receptor partial agonist
ACEA	<i>N</i> -(2-Chloroethyl)-5 <i>Z</i> ,8 <i>Z</i> ,11 <i>Z</i> ,14 <i>Z</i> -eicosatetraenamide	Potent and highly selective synthetic CB ₁ receptor agonist has low affinity for CB ₂
2-AG	(5 <i>Z</i> ,8 <i>Z</i> ,11 <i>Z</i> ,14 <i>Z</i>)-5,8,11,14-Eicosatetraenoic acid, 2-hydroxy-1-(hydroxymethyl)ethyl ester; 2-arachidonoyl glycerol	Endogenous CB ₁ and CB ₂ receptor agonists without any marked selectivity for either subtype. AEA is also an agonist at TRPV1
AEA	<i>N</i> -(2-Hydroxyethyl)-5 <i>Z</i> ,8 <i>Z</i> ,11 <i>Z</i> ,14 <i>Z</i> -eicosatetraenamide; anandamide	
WIN55,212-2	{(<i>R</i>)-(+)-[2,3-Dihydro-5-methyl-3-[(4-morpholino)methyl]pyrrolo-[1,2,3- <i>de</i>]-1,4-benzoxazin-6-yl](1-naphthyl)methanone}	Mixed CB ₁ /CB ₂ receptor agonist
CP-55,940	{(-)-3-[2-Hydroxy-4-(1,1-dimethylheptyl)phenyl]-4-(3-hydroxypropyl)cyclohexan-1-ol}	
HU210	3-(1,1'-Dimethylheptyl)-6 <i>aR</i> ,7,10,10 <i>aR</i> -tetrahydro-1-hydroxy-6,6-dimethyl-6 <i>H</i> -dibenzo[<i>b,d</i>]pyran-9-methanol	
AM251	<i>N</i> -(Piperidin-1-yl)-5-(4-iodophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1 <i>H</i> -pyrazole-3-carboxamide	CB ₁ -selective antagonists
AM281	1-(2,4-Dichlorophenyl)-5-(4-iodophenyl)-4-methyl- <i>N</i> -4-morpholinyl-1 <i>H</i> -pyrazole-3-carboxamide	
SR141716A (rimonabant)	[<i>N</i> -(Piperidin-1-yl)-5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1 <i>H</i> -pyrazole-3-carboxamide hydrochloride] (rimonabant)	

A-836339	[<i>N</i> (<i>Z</i>)]- <i>N</i> -[3-(2-Methoxyethyl)-4,5-dimethyl-2(3 <i>H</i>)-thiazolylidene]-2,2,3,3-tetramethyl-cyclopropanecarboxamide	CB ₂ -selective agonists
AM1241	(2-Iodo-5-nitrophenyl)-(1-(1-methylpiperidin-2-ylmethyl)-1 <i>H</i> -indol-3-yl)methanone	
GW-405,833	1-(2,3-Dichlorobenzoyl)-5-methoxy-2-methyl-3-[2-(4-morpholinyl)ethyl]-1 <i>H</i> -indole	
JTE-907	<i>N</i> -(1,3-Benzodioxol-5-ylmethyl)-7-methoxy-2-oxo-8-pentoxo-1 <i>H</i> -quinoline-3-carboxamide	
JWH-015	(2-Methyl-1-propylindol-3-yl)-naphthalen-1-ylmethanone	
JWH-133	(6 <i>aR</i> ,10 <i>aR</i>)-6,6,9-Trimethyl-3-(2-methylpentan-2-yl)-6 <i>a</i> ,7,10,10 <i>a</i> -tetrahydrobenzo[<i>c</i>]chromene	
NESS400	1-(2',4'-Dichlorophenyl)-6-methyl- <i>N</i> -cyclohexylamine-1,4-dihydroindeno[1,2- <i>c</i>]pyrazole-3-carboxamide	CB ₂ -selective antagonists
O-3223	(6 <i>aR</i> ,10 <i>aR</i>)-6,6,9-Trimethyl-3-(2-methylpentan-2-yl)-6 <i>a</i> ,7,10,10 <i>a</i> -tetrahydrobenzo[<i>c</i>]chromene	
AM630	6-Iodopravadoline	
SR144528	5-(4-Chloro-3-methylphenyl)-1-[(4-methylphenyl)methyl]- <i>N</i> -[(1 <i>R</i> ,3 <i>S</i> ,4 <i>S</i>)-2,2,4-trimethyl-3-bicyclo[2.2.1]heptanyl]pyrazole-3-carboxamide	FAAH inhibitor
OL-135	7-Phenyl-1-(5-pyridin-2-yl-1,3-oxazol-2-yl)heptan-1-one	
PF-3845	<i>N</i> -Pyridin-3-yl-4-[[3-[5-(trifluoromethyl)pyridin-2-yl]oxyphenyl]methyl]piperidine-1-carboxamide	
URB597	(3-Phenylphenyl) <i>N</i> -(4-methoxyphenyl)carbamate	

Continued

Table 1 Compounds Referred to in the Text and Used to Elucidate the Role of Cannabinoids and the Endocannabinoid System in Pain Modulation—cont'd

Pharmacological

Substance	IUPAC Name	Target
JZL184	4-Nitrophenyl-4-[bis(1,3-benzodioxol-5-yl)(hydroxy)methyl]piperidine-1-carboxylate	MAGL inhibitor
URB602	[1,1'-Biphenyl]-3-yl-carbamic acid, cyclohexyl ester	
I-RTX	6,7-Deepoxy-6,7-didehydro-5-deoxy-21-dephenyl-21-(phenylmethyl)-daphnetoxin,20-(4-hydroxy-5-iodo-3-methoxybenzeneacetate); iodoresiniferatoxin	Potent TRPV1 antagonist
AA-5-HT	(5 <i>Z</i> ,8 <i>Z</i> ,11 <i>Z</i> ,14 <i>Z</i>)- <i>N</i> -(2-(5-Hydroxy-1 <i>H</i> -indol-3-yl)ethyl)icosa-5,8,11,14-tetraenamide; <i>N</i> -arachidonoyl-serotonin	FAAH inhibitor and TRPV1 antagonist
ML-193 (alternative name CID 1261822)	<i>N</i> -[4-[(3,4-Dimethyl-5-isoxazolyl)amino]sulfonyl]phenyl]-6,8-dimethyl-2-(2-pyridinyl)-4-quinolinecarboxamide	Potent and selective GPR55 antagonist

the modulation of pain by cannabinoids and the endocannabinoid system. Enhanced thermal analgesia and reduced nociceptive behavior in the formalin and carrageenan models were observed in mice lacking the enzyme fatty acid amide hydrolase (FAAH) which catabolizes AEA and other *N*-acylethanolamines including *N*-palmitoylethanolamide (PEA) and *N*-oleoylethanolamide, compared with wild-type controls (Carey et al., 2016; Cravatt et al., 2001; Lichtman, Shelton, Advani, & Cravatt, 2004). These results suggest that one or more FAAH substrates exert anti-nociceptive actions in these models. FAAH knockout mice, and mice that express FAAH exclusively in nervous tissue, have also been shown to display antiinflammatory and antihyperalgesic effects in both the carrageenan and collagen-induced arthritis models, effects prevented by administration of a CB₂R, but not CB₁R, antagonist (Kinsey, Naidu, Cravatt, Dudley, & Lichtman, 2011; Lichtman et al., 2004). Thus, the augmented levels of AEA in these mice appear to exert tonic analgesia via CB₂R. However, a pronociceptive phenotype of FAAH knockout mice can be unmasked following intradermal injection of the transient receptor potential subfamily V member 1 (TRPV1) agonist capsaicin (Carey et al., 2016). Similarly, mice lacking the 2-AG-catabolizing enzyme monoacylglycerol lipase (MAGL) exhibited significantly augmented nociceptive behavior in the formalin and acetic acid tests and no alterations in thermal tail-withdrawal latency, effects that were likely due to desensitization of CB₁R (Petrenko, Yamazaki, Sakimura, Kano, & Baba, 2014; Schlosburg et al., 2010). In a recent study, nitroglycerin-induced mechanical allodynia and neuronal activation of the trigeminal nucleus to model migraine were abolished in FAAH-deficient mice, results also seen in mice administered FAAH inhibitors (Nozaki, Markert, & Zimmer, 2015). These effects were shown to be CB₁R mediated and they infer that one or more FAAH substrates mediate antinociception via CB₁R. Knockouts of the CB₁R have also been generated and exhibit hypoalgesia in the hot plate, tail immersion, and formalin tests (Valverde, Karsak, & Zimmer, 2005; Zimmer, Zimmer, Hohmann, Herkenham, & Bonner, 1999), suggesting, somewhat paradoxically, a pronociceptive role for CB₁R. However, a different CB₁^{-/-} mouse line displayed similar basal responses to noxious stimuli compared to wild-type animals (Castane et al., 2006; Ledent et al., 1999). Development of mechanical hypersensitivity following partial sciatic nerve ligation was unaltered in CB₁R knockout mice (Castane et al., 2006; Racz, Nent, Erxlebe, & Zimmer, 2015); however, these mice did exhibit more pronounced behavioral manifestations of anxiety-related behaviors compared to wild-type

mice (Racz et al., 2015), suggesting an anxiolytic role for CB₁R. Mice lacking the CB₂R have also been generated as have CB₁/CB₂ double knock-outs (Buckley, 2008; Buckley et al., 2000) and mice overexpressing the CB₂R (La Porta, Bura, Aracil-Fernandez, Manzanares, & Maldonado, 2013). The affective manifestations of osteoarthritis pain in the monosodium iodoacetate model were enhanced in CB₁R knockout mice and absent in CB₂R knockouts, suggesting that the presence of CB₁R attenuates the affective component in this model, while CB₂R is required for expression of the affective component. Both the CB₁R agonist ACEA and the CB₂R agonist JWH-133 ameliorated the nociceptive and affective alterations, with ACEA also improving the associated memory impairment (La Porta et al., 2015). It had previously been shown that development of mechanical allodynia in this model was unaltered in CB₁R and CB₂R knockout mice, but attenuated in those overexpressing CB₂R, compared with wild-type mice (La Porta et al., 2013). In another recent study, paclitaxel-induced mechanical and cold allodynia developed to an equivalent degree in mice lacking CB₁R, CB₂R, and wild-type mice (Deng et al., 2015), suggesting that CB₁R and CB₂R do not impact on the development of the pain-related phenotype in this model. Following intraplantar administration of complete Freund's adjuvant (CFA), or partial nerve ligation, mechanical hyperalgesia was absent in mice lacking GPR55 (Staton et al., 2008), a receptor sensitive to some cannabinoids. However, another study reported thermal hyperalgesia in GPR55 knockout mice (Bjursell et al., 2016) and, most recently, it was shown that genetic deletion of GPR55 did not alter the development of pain-related behavior in a number of mechanistically distinct models of inflammatory and neuropathic pain (Carey et al., 2017).

Shortly after its discovery, Bicher and Mechoulam (1968) showed that Δ^9 -THC was antinociceptive in rabbits (Bicher & Mechoulam, 1968). Since then, many studies have shown that cannabinoids are antinociceptive following systemic administration (for comprehensive review, see Pertwee, 2001). The animal (usually rodent) models used can be divided into three broad groups: (1) acute pain, (2) inflammatory pain involving tissue injury, and (3) neuropathic pain involving peripheral nerve injury. Cannabinoid receptor agonists, administered intraperitoneally, intravenously, subcutaneously, or orally, demonstrate analgesic efficacy (to greater or lesser degrees depending on the compound and model under investigation) across these models (Finn & Chapman, 2004; Pertwee, 2001) with a potency that is comparable with, or greater than, some opiates and cyclooxygenase inhibitors

(Bloom & Dewey, 1978; Smith, Cichewicz, Martin, & Welch, 1998; Sofia, Vassar, & Knobloch, 1975; Thorat & Bhargava, 1994). Many of the earlier studies used nonselective cannabinoid receptor agonists; however, the involvement of CB₁ and/or CB₂ receptors has been probed with selective antagonists, and more recent studies have assessed the efficacy of agonists with selectivity for CB₁ or CB₂ receptors. Interpretation of the results of studies employing systemic administration of cannabinoids can, however, sometimes be complicated by cannabinoid-mediated suppression of motor activity. To examine the specific sites and mechanisms through which cannabinoids reduce pain, studies have investigated the antinociceptive activity of cannabinoids and endocannabinoid system modulators administered supraspinally, spinally, and peripherally. These site-specific studies are the key focus of the present review.



2. ANATOMICAL LOCALIZATION OF THE ENDOCANNABINOID SYSTEM THROUGHOUT THE PAIN PATHWAY

Two major ascending pain pathways in mammals, the spinothalamic pathway and the spinoparabrachial pathway, encode the sensory-discriminatory and affective aspects of pain, respectively (see Fig. 1). In addition, the descending pain pathway originates in higher cortical regions and in the amygdala and hypothalamus, and projects (via the periaqueductal gray (PAG)) to the lower brain stem and spinal cord. Descending control of pain can be either inhibitory or facilitatory depending on the precise circuitry and receptors that are engaged (Millan, 2002; Ossipov, Morimura, & Porreca, 2014; Suzuki & Dickenson, 2005; Suzuki, Rygh, & Dickenson, 2004). The endocannabinoid system is expressed throughout the ascending and descending pain pathways at peripheral, spinal, and supraspinal sites (Fig. 1). CB₁ receptors are located on peripheral endings and central terminals of primary afferent neurons (Hohmann, Briley, & Herkenham, 1999; Hohmann & Herkenham, 1998, 1999a). CB₁ receptors are also found in the dorsal root ganglion (DRG) and in the superficial laminae of the spinal cord (Farquhar-Smith et al., 2000; Glass, Dragunow, & Faull, 1997; Herkenham et al., 1991; Hohmann & Herkenham, 1999b; Ross et al., 2001; Sanudo-Pena, Strangman, Mackie, Walker, & Tsou, 1999). Ahluwalia, Urban, Bevan, Capogna, and Nagy (2002) reported that 80% of CB₁R-expressing neurons either contained calcitonin gene-related peptide (CGRP), a marker for peptidergic neurons, or bound IB4, a marker




Fig. 1 Cannabinoid receptor distribution throughout the pain pathways. Cannabinoid receptors are present at all three levels of pain processing: (A) in the periphery: CB₁R is present in the peripheral sensory nerve endings, both CB₁R and CB₂R are expressed in the dorsal root ganglion (DRG); (B) in the spinal cord: CB₁R is found in the dorsolateral funiculus, in the surroundings of the central canal, and in the superficial dorsal horn. CB₂R is expressed on glial cells highly restricted to lumbar spinal cord; its expression coincides with the appearance of activated microglia, and (C) in the supraspinal sites: CB₁R is distributed in areas of the brain involved in pain processing, perception, and modulation, e.g., thalamus, amygdala, parabrachial nucleus, periaqueductal gray matter, and rostral/ventromedial medulla. They are also present in caudate nucleus and putamen (n. accumbens), basal ganglia, hypothalamus, and cerebellum. CB₂R is expressed in some neurons within the brain stem, and also on glial cells in the cerebellum and cortex. CB₁R and CB₂R distribution in regions involved in pain transduction, transmission, perception, and modulation provides the anatomical basis for the well-known ability of CB₁/CB₂R agonists to decrease pain.

for an unmyelinated neurons which express glycoproteins (Ahluwalia et al., 2002), suggesting a functional role for CB₁R on peripheral nerve terminals. However, there is also evidence that CB₁R mRNA is expressed predominantly in medium- and large-sized DRG neurons, with lower levels in DRG neurons expressing substance P or CGRP mRNA (Hohmann & Herkenham, 1999b). In addition to its peripheral and spinal localization, CB₁R is also located in all of the major brain regions involved in pain processing and modulation. Receptor autoradiography and immunohistochemistry studies have demonstrated the presence of CB₁R in the cortex, amygdala, hypothalamus, thalamus, PAG, parabrachial nucleus, and in brain stem regions including the rostral ventromedial medulla (RVM) (Glass et al., 1997; Herkenham et al., 1991, 1990; Mailleux, Parmentier, & Vanderhaeghen, 1992; Thomas, Wei, & Martin, 1992; Tsou, Brown, Sanudo-Pena, Mackie, & Walker, 1998). CB₁R localization is predominantly presynaptic, and its direct activation by synthetic agonists, or by endocannabinoids that signal retrogradely, inhibits the release of neurotransmitters including GABA and glutamate (Rea, Roche, & Finn, 2007).

The clinical utility of cannabinoids acting at CB₁R can be limited due to adverse central side effects and the development of tolerance (De Vry, Jentsch, Kuhl, & Eckel, 2004; Gonzalez, Cebeira, & Fernandez-Ruiz, 2005). This has led to increased interest in the role of the CB₂R in pain. The CB₂R has been categorized classically as the peripheral cannabinoid receptor due to its presence on the cells and tissues of the immune, reproductive, cardiovascular, gastrointestinal, and respiratory systems and numerous reports which were unable to detect CB₂R transcripts in normal healthy brain (Derbenev, Stuart, & Smith, 2004; Facci et al., 1995; Griffin et al., 1999; Munro et al., 1993). However, more recent evidence suggests that CB₂R is present in the brain under normal and, in particular, under pathological/inflammatory conditions (Baek, Zheng, Darlington, & Smith, 2008; Concannon, Okine, Finn, & Dowd, 2015; Onaivi, Ishiguro, Gong, et al., 2006; Roche & Finn, 2010; Van Sickle et al., 2005; Zhang et al., 2014), although to a much lesser extent than the ubiquitously expressed CB₁R. CB₂R expression has been demonstrated within pain-related brain regions including the cerebral cortex, hippocampus, striatum, amygdala, thalamic nuclei, PAG, cerebellum, and several brain stem nuclei of the rodent brain (Ashton, Friberg, Darlington, & Smith, 2006; Brusco, Tagliaferro, Saez, & Onaivi, 2008; Gong et al., 2006; Onaivi et al., 2008; Onaivi, Ishiguro, Gong, et al., 2006; Onaivi, Ishiguro, Sejal, et al., 2006;

Suarez et al., 2008; Van Sickle et al., 2005). Although many studies have identified central CB₂R on glial and endothelial cells, there is also evidence to support the expression of CB₂R on subpopulations of neurons within the central nervous system (Ashton et al., 2006; Beltramo et al., 2006; Gong et al., 2006; Molina-Holgado et al., 2007; Onaivi, Ishiguro, Gong, et al., 2006; Palazuelos et al., 2006; Suarez et al., 2008; Van Sickle et al., 2005; Viscomi et al., 2009; Zhang et al., 2014). There is evidence for expression of CB₂R in DRG and in the dorsal horn of the spinal cord and upregulation during neuropathic or inflammatory pain (Anand et al., 2008; Hsieh et al., 2011; Romero-Sandoval & Eisenach, 2007; Romero-Sandoval, Natile-McMenemy, & DeLeo, 2008; Ross et al., 2001; Svizenska, Brazda, Klusakova, & Dubovy, 2013; Wotherspoon et al., 2005; Zhang et al., 2003). The high expression of the CB₂R in tissues of the immune system including the spleen and thymus as well as on specific immune cells including B lymphocytes, natural killer cells, monocytes, neutrophils and T lymphocytes (Berdyshev, 2000; Howlett et al., 2002; Klein, Newton, & Friedman, 2001; Munro et al., 1993; Sugiura et al., 1995) has focused research on the viability of the CB₂R as a therapeutic target in inflammatory pain conditions in particular, but also neuropathic pain which can have a neuroinflammatory/neuroimmune component (Milligan et al., 2003; Watkins, Milligan, & Maier, 2003).

In addition to the cannabinoid receptors, other components of the endocannabinoid system are also present throughout the ascending and descending pain pathways. Thus, the endocannabinoids, *N*-acylethanolamines, and their metabolizing enzymes are localized in peripheral tissues innervated by primary afferent nociceptive neurons (Calignano, La Rana, Giuffrida, & Piomelli, 1998; Felder et al., 1996), spinal cord (Di Marzo et al., 2000; Egertova, Giang, Cravatt, & Elphick, 1998; Tsou, Nogueron, et al., 1998), and brain (Devane et al., 1992; Egertova et al., 1998; Hanus et al., 2001; Huang et al., 2002; Porter et al., 2002; Stella, Schweitzer, & Piomelli, 1997; Tsou, Nogueron, et al., 1998) tissues, including regions important in pain. Elegant *in vivo* microdialysis experiments demonstrated that intraplantar injection of the chemical irritant formalin evokes the release of AEA in the midbrain PAG (Walker, Huang, Strangman, Tsou, & Sanudo-Pena, 1999). The endocannabinoids and *N*-acylethanolamines also have affinity for, and activity at, a number of non-CB₁/non-CB₂ receptors, including TRPV1, GPR55 (putative CB₃ receptor), and the peroxisome proliferator-activated receptors (PPARs) (Alexander & Kendall, 2007; Wiley & Martin, 2002), all of which are also expressed throughout the pain

pathways and likely play important roles in endocannabinoid-mediated regulation of pain. The remainder of this review will focus on functional *in vivo* studies of cannabinoids and the endocannabinoid system in models of acute, inflammatory, and neuropathic pain with a focus on supraspinal, spinal, and peripheral sites and mechanisms of action.



3. SUPRASPINAL SITES AND MECHANISMS OF ACTION

3.1 Evidence From Acute Pain Models

In the 1990s, it was demonstrated that the inhibitory effects of the cannabinoid receptor agonists CP-55,940, THC, and WIN55,212-2, administered systemically, on either tail-flick responding (Lichtman & Martin, 1991) or noxious-evoked responses of spinal neurons (Hohmann, Tsou, & Walker, 1999a) are abolished in rats following spinal transection, results that suggested an important role for descending inhibitory pathways in mediating cannabinoid-induced antinociception. The realization that CB₁R is present in moderate to high densities in brain regions which play an important role in nociceptive processing (see Section 2) also prompted investigation of supraspinal sites of action mediating cannabinoid-induced antinociception. Multiple studies have now shown that synthetic or plant-derived cannabinoid receptor agonists, or endogenous cannabinoid ligands, display antinociceptive activity in the mouse and rat tail-flick tests, following intracerebroventricular administration (Fang et al., 2012; Lichtman & Martin, 1997; Martin, Lai, Patrick, Tsou, & Walker, 1993; Pan et al., 2014; Raffa, Stone, & Hipp, 1999; Welch, 1994; Welch, Huffman, & Lowe, 1998; Welch, Thomas, & Patrick, 1995; Zheng et al., 2017).

Significant effort has also been directed at elucidating the specific brain regions that mediate the antinociceptive effects of cannabinoid receptor agonists (Corcoran, Roche, & Finn, 2015). In early work by Martin and colleagues, direct administration of WIN55,212-2 into a number of different brain regions including the amygdala, thalamus, superior colliculus, and A5 region was shown to be antinociceptive in the tail-flick test (Martin, Coffin, et al., 1999). Microinjection of the nonselective cannabinoid receptor agonists WIN55,212-2 and HU210 into the RVM also elevated tail-flick latencies in rats (Martin, Tsou, & Walker, 1998; Meng & Johansen, 2004). Moreover, the effects of HU210 were attenuated by co-administration with the CB₁R antagonist/inverse agonist rimonabant (Martin et al., 1998). Further evidence for the importance of the RVM in cannabinoid-induced attenuation of acute pain came from a study demonstrating that GABA_A

receptor agonist-mediated inactivation of the RVM prevented antinociceptive effects of systemically administered WIN55,212-2 in the rat tail-flick test (Meng, Manning, Martin, & Fields, 1998). In addition, it has been shown that the antinociceptive effects of intra-RVM administration of WIN55,212-2 in the tail-flick test are associated with inhibition of ON-cell activity and an increase in OFF-cell activity, effects blocked by rimonabant (Meng & Johansen, 2004). Within the RVM, the nucleus reticularis gigantocellularis pars alpha appears to be an important locus for cannabinoid-mediated antinociception (Monhemius, Azami, Green, & Roberts, 2001).

The PAG, another major component of the descending inhibitory pain pathway, is also an important locus for the antinociceptive effects of cannabinoids. Electrical stimulation of the dorsal or lateral columns of the PAG resulted in CB₁R-mediated antinociception in the rat tail-flick test which was accompanied by a marked increase in AEA release in the PAG (Walker et al. 1999). Intraplantar injection of formalin also resulted in increased AEA release, suggesting engagement of an endogenous cannabinergic pain modulatory system in this midbrain region. Direct administration of CP-55,940 into the ventrolateral (vl) PAG (Lichtman, Cook, & Martin, 1996) and of WIN55,212-2 into the dorsal (dl) PAG (Martin, Coffin, et al., 1999; Martin, Patrick, Coffin, Tsou, & Walker, 1995) had antinociceptive effects in the rat tail-flick test. In vitro studies of the mechanism of action of cannabinoids at the level of the PAG suggest that cannabinoids reduce neurotransmitter release from presynaptic terminals and inhibit GABAergic and glutamatergic transmission (Vaughan, Connor, Bagley, & Christie, 2000). Thus, the antinociceptive effects of cannabinoid agonists administered into the PAG may arise from the disinhibition of GABAergic interneurons and the activation of the descending inhibitory controls, with subsequent inhibition of excitatory transmission at the level of the spinal cord. There is also evidence for a CB₁–glutamatergic interaction in the dlPAG in mediating cannabinoid-induced antinociception in the plantar test in rats (Palazzo et al., 2001). The suppression of acute pain (tail-flick response) following exposure to acute stress (footshock) via the phenomenon of stress-induced analgesia has also been shown to be mediated by endocannabinoids acting at CB₁R in the dlPAG and RVM (Hohmann et al., 2005; Suplita, Farthing, Gutierrez, & Hohmann, 2005). There is also evidence that the cannabinoid receptor agonist HU210 can enhance the antinociceptive effects of morphine, and vice versa, with a site of action in the vlPAG (Wilson, Maher, & Morgan, 2008;

Wilson-Poe, Pocius, Herschbach, & Morgan, 2013). In addition to its activity at cannabinoid receptors, AEA also acts at TRPV1, a receptor that also plays an important role in supraspinal modulation of pain (Madasu, Roche, & Finn, 2015). The TRPV1 agonist capsaicin has been shown to induce initial hyperalgesia in the tail-flick test, followed by antinociception, when injected into the dIPAG (McGaraughty et al., 2003). Similarly, in the rat plantar test, biphasic effects of intra-dIPAG administration of capsaicin have been demonstrated (Palazzo et al., 2002) and intra-vIPAG administration of capsaicin results in glutamate release in the RVM, thereby activating OFF cells and producing antinociception (Starowicz et al., 2007). In further work using the rat plantar test, intra-vIPAG injection of a low dose of the FAAH inhibitor URB597 with the CB₁ receptor antagonist/inverse agonist AM251 converted the hyperalgesic effect of low dose URB597 to an antinociceptive effect, while coadministration of URB597 with both the TRPV1 antagonist capsazepine and AM251 abolished all effects (Maione et al., 2006). In comparison, the antinociceptive effect of high dose URB597 was converted to a hyperalgesic effect following TRPV1 antagonism. The URB597-induced antinociceptive effects (TRPV1 mediated) and pronociceptive effects (CB₁ receptor mediated) were associated with enhanced or reduced RVM OFF-cell activity, respectively, suggesting URB597-induced modulation of the activity of excitatory PAG output neurons (Maione et al., 2006). Intra-vIPAG injection of the dual FAAH inhibitor and TRPV1 antagonist AA-5-HT increased endocannabinoid levels and had an antinociceptive effect in the rat tail-flick test, with associated inhibition of RVM ON- and OFF-cell activity (de Novellis et al., 2008). These effects were blocked by the CB₁ receptor antagonist AM251 or the TRPV1 antagonist I-RTX and were mimicked by intra-vIPAG coadministration of the FAAH inhibitor URB597 with the TRPV1 antagonist I-RTX (de Novellis et al., 2008). Thus, activity of the descending pain pathway is regulated by the action of endocannabinoids at both CB₁R and TRPV1 in the vIPAG. For an excellent schematic of the possible mechanisms underlying endocannabinoid/endovanilloid-mediated control of nociception in the ventrolateral PAG and RVM, see scheme 1 within Maione et al. (2006). Recently, it has also been shown that intra-PAG administration of the GPR55 agonist lysophosphatidylinositol reduces the nociceptive threshold in the rat hot plate test, an effect blocked upon pretreatment with the GPR55 antagonist ML-193 (Deliu et al., 2015), thereby suggesting a role for this putative CB₃ receptor in the PAG in acute pain processing.

The amygdala is thought to play a role in the affective component of pain and is also a component of the descending pain pathway (Neugebauer, Galhardo, Maione, & Mackey, 2009; Neugebauer, Li, Bird, & Han, 2004). Direct administration of WIN55,212-2 into either the basolateral (BLA) or central (CeA) nucleus of the amygdala has been shown to increase tail-flick latency in rats (Hasanein, Parviz, Keshavarz, & Javanmardi, 2007; Martin, Coffin, et al., 1999). Intra-CeA, but not intra-BLA, administration of muscimol, significantly attenuated the antinociceptive effects of systemically administered WIN55,212-2 in rats (Manning, Martin, & Meng, 2003). Another study from the same group found that the amygdala also plays a role in cannabinoid-induced antinociception in nonhuman primates (Manning, Merin, Meng, & Amaral, 2001). Pharmacological blockade of CB₁R in the rat BLA attenuated the stress-induced suppression of nociceptive responding in the tail-flick test (Connell, Bolton, Olsen, Piomelli, & Hohmann, 2006). A role for CB₁R signaling in the rat prelimbic cortex in facilitation of stress-induced analgesia has also been demonstrated (Freitas, Salgado-Rohner, Hallak, Crippa, & Coimbra, 2013). Using fMRI, it has been shown that THC reduces the reported unpleasantness, but not the intensity of ongoing pain and hyperalgesia, induced by capsaicin in healthy human subjects, an effect positively correlated with amygdala activity. THC also reduced functional connectivity between the amygdala and primary sensorimotor areas during the ongoing-pain state (Lee et al., 2013).

3.2 Evidence From Inflammatory Pain Models

Some studies have investigated the effects of intracerebral administration of cannabinoids specifically in animal models of inflammatory pain. Direct microinjection of WIN55,212-2 into the nucleus reticularis gigantocellularis pars alpha, a major source of descending modulation, reduced formalin-evoked pain behavior, via the CB₁ receptor (Monhemius et al., 2001). Administration of the potent cannabinoid receptor agonist HU210 into the dlPAG inhibited formalin-evoked nociceptive behavior during the second phase and was antiaversive in rats (Finn et al., 2004, 2003). Intra-vPAG administration of AA-5-HT to rats prevented the changes in ON- and OFF-cell firing activity induced by intraplantar injection of formalin and reversed the formalin-induced increase in locus coeruleus adrenergic cell activity (de Novellis et al., 2008). Injection of the CB₁ receptor antagonist AM251 into the PAG or RVM reverses metazolinol-induced analgesia in the rat carrageenan model of inflammatory pain, suggesting a role for the

endocannabinoid system in these brain regions in NSAID-induced analgesia (Escobar et al., 2012). These data provide additional evidence that the RVM and PAG are important brain regions mediating the antinociceptive effects of cannabinoids in animal models of inflammatory pain. Evidence that pharmacological blockade of CB₁R in the dlPAG attenuates conditioned fear-induced suppression of formalin-evoked nociceptive behavior (i.e., fear-conditioned analgesia) further substantiates the key role of the endocannabinoid system in the PAG in stress-induced analgesia (Olango, Roche, Ford, Harhen, & Finn, 2012). Conversely, anxiety and depression may exacerbate pain and are frequently found comorbid with chronic pain. Finn and coworkers have demonstrated that hyperalgesia to intraplantar formalin injection in Wistar–Kyoto rats that exhibit an anxiodepressive phenotype (vs Sprague–Dawley counterparts) is associated with impaired endocannabinoid–CB₁R signaling in the RVM (Rea et al., 2014). Recently, it has been shown that while CB₁R-mediated inhibition of GABAergic neurons in the RVM is reduced in the rat CFA model, CB₂R functionality in this region is increased in this model of persistent inflammatory pain (Li, Suchland, & Ingram, 2017), supporting the contention that CB₂R may represent a viable analgesic target.

Unilateral inactivation of the CeA reduced the suppression of formalin-evoked c-Fos expression by WIN55,212-2 in the superficial dorsal horn of the spinal cord (Manning et al., 2003). Furthermore, intra-BLA administration of WIN55,212-2 has also been shown to reduce formalin-evoked nociceptive behavior in rats, an effect attenuated by intra-BLA administration of the CB₁R antagonist AM251 (Hasanein et al., 2007). Interestingly, intra-BLA administration of rimonabant has also been reported to attenuate formalin-evoked nociceptive behavior and associated increases in c-Fos immunoreactivity in the hippocampus and RVM in rats (Roche, O'Connor, Diskin, & Finn, 2007; Roche et al., 2010), although intra-BLA administration of AM251 did not have this effect (Rea et al., 2013). In contrast, intra-BLA administration of AM251 (Rea et al., 2013), but not rimonabant (Roche et al., 2010, 2007), attenuated fear-conditioned analgesia in rats. The same doses of rimonabant and AM251 were micro-injected into the BLA in these studies and under very similar methodological conditions. However, as discussed in Rea et al. (2013), discrepancies between the effects of the two CB₁R antagonists/inverse agonist may relate to dose–response differences between the two compounds when administered into this brain region or to differential activity of the two compounds at non-CB₁R targets expressed in the BLA (e.g., GPR55, TRPV1, or

PPARs). There is also evidence that fear-conditioned analgesia is mediated by endocannabinoid-CB₁R signaling in the ventral hippocampus (Ford, Kieran, Dolan, Harhen, & Finn, 2011).

In the rat kaolin/carrageenan intraarticular injection model of arthritis, coactivation of mGluR5 and CB₁R increased activity of prefrontal cortex neurons and inhibited pain-related neuronal activity in the CeA (Ji & Neugebauer, 2014). Further evidence for a role of the endocannabinoid system in the prefrontal cortex in arthritic conditions comes from work demonstrating that osteoarthritis pain is associated with increased 2-AG levels in the prefrontal cortex of mice in the monosodium iodoacetate model (La Porta et al., 2015). Recently, Finn and coworkers demonstrated that the antinociceptive effects of PEA injected into the anterior cingulate cortex in the rat formalin test are likely mediated by AEA-induced activation of CB₁R in this brain region arising from substrate competition between PEA and AEA at FAAH (Okine et al., 2016). A facilitatory role for PPARs and TRPV1 in the anterior cingulate cortex in formalin-evoked nociceptive behavior has also been suggested (Okine et al., 2016, 2014).

3.3 Evidence From Neuropathic Pain Models

Increased levels of AEA and 2-AG have been reported in the PAG and RVM of rats 7 days postchronic constriction injury (CCI) of the sciatic nerve, when hyperalgesia and mechanical allodynia were observed to be maximal (Petrosino et al., 2007). Partial sciatic nerve injury has been shown to reduce formalin-evoked pain behavior in rats (Monhemius et al., 2001). This effect was blocked by direct administration of rimonabant into the nucleus reticularis gigantocellularis pars, suggesting that increased endocannabinoid tone in neuropathic rats can modulate nociceptive behavior (Monhemius et al., 2001). In the thalamus, CB₁R mRNA is upregulated in a rat model of neuropathic pain (Siegling, Hofmann, Denzer, Mauler, & De Vry, 2001). Potentially, upregulation of thalamic CB₁R in neuropathic pain states may serve to enhance the analgesic effects of cannabinoids under these conditions. Interestingly, it has been shown that CB₂R plays a functional role in the modulation of responses of neurons in the ventral posterior nucleus of the thalamus in spinal nerve-ligated, but not sham-operated, rats (Jhaveri et al., 2008).

TRPV1 expression is increased in glutamatergic neurons of the medial prefrontal cortex following spared nerve injury (SNI) in rats (Giordano et al., 2012). Moreover, SNI-induced neuropathic pain is also associated with

increased levels of endovanilloids and endocannabinoids in the medial prefrontal cortex and direct administration of AA-5-HT into the prelimbic and infralimbic cortices reduces nociceptive behavior in rats following SNI (de Novellis et al., 2011; Giordano et al., 2012).



4. SPINAL SITES AND MECHANISMS OF ACTION

4.1 Evidence From Acute Pain Models

Early evidence that the synthetic cannabinoid levonantradol produced a dose-dependent increase in the hot plate and tail-flick response latencies following intrathecal (i.t.) administration (Yaksh, 1981), followed by studies elucidating mechanisms of THC-induced analgesia (Smith & Martin, 1992), indicated a spinal component in the antinociceptive action of the cannabinoids. Behavioral (Smith & Martin, 1992; Yaksh, 1981), electrophysiological (Hohmann, Tsou, & Michael Walker, 1998; Johannek, Simone, & Lisa, 2005; Sokal, Elmes, Kendall, & Chapman, 2003), and neurochemical (Hohmann, Tsou, & Walker, 1999a, 1999b) studies have demonstrated that cannabinoids act at the spinal level to suppress nociceptive processing. In a model of tonic pain, immunocytochemistry for the proto-oncogene *c-fos* (a marker for the activation of nociceptive neurons in the spinal cord) was used to demonstrate that cannabinoids reduce behavioral responses to noxious stimuli by decreasing spinal processing of nociceptive inputs (Tsou, Martin, & Bereiter, 1996).

4.2 Evidence From Inflammatory Pain Models

The CB₁R has been suggested to be tonically active in the spinal cord under normal conditions, and its activity is increased in response to injections of CFA in the plantar surface of the rat hind paw (Martin, Loo, & Basbaum, 1999). The synthetic mixed CB₁R/CB₂R agonist WIN55,212-2 reverses inflammation-induced allodynia at doses that do not produce analgesia; additionally rimonabant differentially affects the pattern of Fos expression in the spinal cord, depending on the presence or absence of inflammation (Martin, Coffin, et al., 1999).

A functional inhibitory effect of i.t. administration of the CB₂R-selective agonists A-836339 and AM1241 has been demonstrated in CFA-induced chronic inflammatory pain (Hsieh et al., 2011). These data complement the findings that CB₂R mRNA is upregulated in the spinal cord only from rats under inflammatory conditions, suggesting that

CB₂R agonists may elicit analgesic effects by acting not only at peripheral DRG sites but also at central levels of the spinal cord, making CB₂ an attractive target for chronic pain treatment, avoiding the adverse psychotropic effects that can accompany CB₁R-based therapies. The antinociceptive effects of A-836339 were not sensitive to pretreatment with naloxone, and thus are not mediated by μ -opioid receptors. Interestingly, the blockade of AM1241 by naloxone was observed in the CFA model of inflammatory pain (Hsieh et al., 2011).

4.3 Evidence From Neuropathic Pain Models

Cannabinoids suppress C-fiber-evoked responses of dorsal horn neurons recorded in a rat model of neuropathic pain (Elmes, Jhaveri, Smart, Kendall, & Chapman, 2004). The synaptic processes that produce “windup,” the phenomenon whereby repeated stimulation of cutaneous C-fibers at frequencies >0.3 Hz gives increasing responses of dorsal horn cells and withdrawal reflexes, are sufficient to produce central sensitization, which appears to be an important component of hyperalgesia and allodynia. The effect of cannabinoids, namely of the potent, synthetic cannabinoid receptor agonist WIN55,212-2 on windup of spinal dorsal horn neurons was investigated in 1999 (Strangman, Walker, & Strangman, 1999). Strangman and Walker provided the first direct evidence that cannabinoids inhibit the activity-dependent facilitation of spinal nociceptive responses. These authors suggested that cannabinoids may act as general inhibitors of central sensitization by inhibiting calcium entry (Strangman et al., 1999).

The effectiveness of cannabinoids is inconsistent in preclinical neuropathic pain models. WIN55,212-2 delivered i.t. is effective in mitigating mechanical allodynia in the CCI model (Lim, Sung, Ji, & Mao, 2003), while Costa et al. (2005) demonstrated that systemic administration of a CB₁R antagonist significantly reduces mechanical and thermal hyperalgesia in CCI rats and in mice. Others (Toniolo et al., 2014; Ueda et al., 2014) have also suggested that CB₁R expression and activation can be maladaptive. Very recent research indicates that CB₁R expression contributes to the development of persistent mechanical hypersensitivity, protects against the development of cold allodynia, but is not involved in motor impairment following SNI in mice (Sideris et al., 2016).

Although nerve injury increased CB₂R expression in spinal microglia (Zhang et al., 2003), CB₂R agonists suppressed microglial activation and reduced neuropathic pain symptoms (Wilkerson et al., 2012). I.t. delivery

of the CB₂R agonist JWH-015 reverses hypersensitivity following nerve injury in a CB₂R- and not CB₁R-dependent manner (an effect blocked by AM630 but not AM281) (Romero-Sandoval & Eisenach, 2007). Interestingly, CB₂R knockout mice displayed increased microglial and astrocytic reactivity in the spinal cord and enhanced neuropathic pain symptoms, whereas transgenic mice overexpressing CB₂R showed attenuated glial reactivity and neuropathic pain (Racz et al., 2008). CB₂R is upregulated on both microglia and astrocytes following SNI in mice, and chronic systemic administration of the CB₂R agonist NESS400 reduces pain behavior, astrogliosis, microglial activation, and levels of proinflammatory cytokines, while promoting levels of antiinflammatory cytokines (Luongo et al., 2010).

Interestingly, the CB₂-selective agonists A-836339 and AM1241, which have previously been shown to counteract inflammatory pain, have also been proven to alleviate neuropathic pain in the rat spinal nerve ligation (SNL) model (Hsieh et al., 2011). As in the case of CFA-induced inflammation, A-836339 action was opioid insensitive, while the blockade of AM1241 by naloxone was not observed. The reason for the difference between two drugs is currently unknown. AM1241 may interact with additional targets that may contribute to the antinociceptive efficacy through the regulation of the opioid receptor pathway (Hsieh et al., 2011). However, there is some conflicting evidence in the literature, with a recent study reporting no effect of the CB₂ agonists GW-405,833 and JWH-133 on mechanical allodynia in CCI model of neuropathy (Brownjohn & Ashton, 2012). This study also reported no elevation of CB₂ at either the protein or mRNA level, probably due to the choice of neuropathic pain model (SNL or CCI).

The endocannabinoids AEA and 2-AG are also increased in the spinal cord following induction of a neuropathic pain state in a CCI model (Petrosino et al., 2007; Starowicz et al., 2012), suggesting that pharmacological manipulation of endocannabinoid accumulation or breakdown may suppress neuropathic nociception in rodents. Both FAAH and MAGL represent potential therapeutic targets for the development of pharmacological agents to treat chronic pain resulting from nerve injury. A significant reduction of neuropathic pain symptoms following inhibition of the AEA hydrolytic enzyme with URB597 in a rat CCI model was reported (Starowicz et al., 2013, 2012). Depending on the dose of URB597 used, and on the consequent lesser or higher elevation of endogenous AEA levels, analgesia was mediated via CB₁ or TRPV1 receptors, respectively. These data suggest that indirect modulation of TRPV1 function, as well as strengthening endogenous AEA signaling by inhibiting its enzymatic degradation, together

hold promise for the development of novel multitarget pharmacological treatments. These studies highlight the importance of the endocannabinoid system as a potential therapeutic target for treatment of neuropathic pain.



5. PERIPHERAL SITES AND MECHANISMS OF ACTION

5.1 Evidence From Acute Pain Models

In behavioral experiments, administration of the endogenous CB₁R agonist, AEA, into the ipsilateral hind paw of the rat reduced formalin-induced nociception (Calignano et al., 1998), indicating that activation of peripheral CB₁R produces antinociception. PEA produced a similar effect by activating peripheral CB₂R. Furthermore, PEA was administered together with AEA, the two compounds acted synergistically. The peripheral actions of CB₁R agonists are attributed to an inhibition of both the sensitizing effects of NGF and CGRP release (Rice, Farquhar-Smith, & Nagy, 2002; Richardson, Kilo, & Hargreaves, 1998).

In 2001, it was demonstrated that selective activation of peripheral CB₂R results in antinociception (Malan et al., 2001). AM1241, the CB₂R-selective agonist, administered both locally and systemically (i.p.) produced thermal hypoalgesia, which was absent when the compound was coadministered with AM630, a CB₂R antagonist, but not AM251, the CB₁R antagonist. AM1241 administered locally to the contralateral paw did not elicit antinociception, which suggests a local site of action. Moreover, local administration of AM630 blocked the antinociceptive effect of AM1241 injected i.p., further implicating peripheral CB₂R as the main site of action. Ibrahim et al. (2005) reported that CB₂R activation produces antinociception by stimulating the release of β -endorphin from keratinocytes, which in turn acts at μ -opioid receptors on primary afferent neurons. Furthermore, it was also suggested that other mediators might be released from local cells after activation of CB₂R, contributing to its antinociceptive effects. Nonetheless, β -endorphin release was suggested to be critical for CB₂R-mediated antinociception because the effects of AM1241 were completely prevented by a β -endorphin-sequestering antiserum (Ibrahim et al., 2005).

Inhibition of endocannabinoid metabolism is considered a promising therapeutic target on its own. It has been demonstrated that blocking AEA degradation results in antinociceptive effects in the mouse hot plate test (Kathuria et al., 2003). The carbamate compound URB597 reduces pain-related behavior in the rat produced by prior i.p. injection of CFA in a manner blocked by a CB₁R but not a CB₂R antagonist (Wilson,

Clayton, Medhurst, Bountra, & Chessell, 2004). Also global deletion of FAAH results in lower inflammatory response to local administration of carrageenan (Lichtman et al., 2004). There is good evidence in the literature that CB₂R may regulate oedema and hyperalgesia in response to carrageenan (Holt, Comelli, Costa, & Fowler, 2005). Antioedemic effect of the CB₂R agonists, AM1241 and JTE-907, was demonstrated (Iwamura, Suzuki, Ueda, Kaya, & Inaba, 2001; Quartilho et al., 2003). Moreover, URB597 reduced oedema formation in a CB₂R-dependent manner (Holt et al., 2005).

5.2 Evidence From Inflammatory Pain Models

Studies have demonstrated that administration of the endogenous CB₁R agonist, AEA, into the ipsilateral hind paw of the rat reduces carrageenan-induced hyperalgesia (Richardson et al., 1998) and that administration of the PEA reduced oedema and inflammatory hyperalgesia (Mazzari, Canella, Petrelli, Marcolongo, & Leon, 1996). It was demonstrated that activation of CB₂R suppresses the development of inflammatory pain (Nackley, Makriyannis, & Hohmann, 2003). AM1241, when injected i.p., suppressed the development of carrageenan-evoked thermal and mechanical hyperalgesia as well as allodynia in a CB₂-dependent manner. Furthermore, intraplantar administration suppressed hyperalgesia and allodynia only on the inflamed paw and was inactive following administration in the contralateral (noninflamed) paw (Nackley et al., 2003).

As a result of systemic administration of the selective FAAH inhibitor, URB597, elevation in endogenous AEA levels reduced the mechanical allodynia and thermal hyperalgesia in an inflammatory pain model in both CB₁R- and CB₂R-dependent manner (Jayamanne et al., 2006). Moreover, two distinct inhibitors of MAGL (JZL184 and URB602) elicited local analgesia in the formalin-induced pain model that involved both CB₁R and CB₂R. URB602 produced regionally restricted increases in 2-AG levels in rat hind paw skin without altering AEA levels (Guindon, Guijarro, Piomelli, & Hohmann, 2011). The earlier findings indicate that increase in endocannabinoid tone blocks the development of inflammatory pain.

5.3 Evidence From Neuropathic Pain Models

Studies by Fox et al. (2001) and Elmes et al. (2004) showed that antinociceptive effects in the partial sciatic nerve ligation and SNL models were produced by the activation of peripheral CB₁R and CB₂R, respectively.

In particular, WIN55,212-2 reversed mechanical hyperalgesia following intraplantar administration into the ipsilateral hind paw (Fox et al., 2001). CB₁ mRNA is localized in DRG neurons, and CB₁R has been shown to undergo peripheral axonal flow in the sciatic nerve (Hohmann & Herkenham, 1999a, 1999b). Moreover, data from Hargreaves' group indicate that CB₁R activation inhibits sensory neuropeptide release from the skin of rat hind paws, demonstrating a functional inhibitory activity on peripheral sensory nerves (Richardson et al., 1998). JWH-133, a cannabinoid CB₂R agonist, also significantly reduced noxious mechanically evoked responses of wide dynamic range dorsal horn neurons following intraplantar injections (Elmes et al., 2004). Indeed CB₂ agonists offer promise in neuropathic pain management. CCI of the sciatic nerve-induced neuropathic pain behavior and bilateral elevation of both CB₂R protein and mRNA in lumbar L4–L5 as well as cervical C7–C8 DRG when compared with naive animals. CB₂R protein and mRNA were increased not only in DRG neurons but also in satellite glial cells. Such changes suggest propagation of neuroinflammation alongside the neuraxis and the neuroprotective effects of CB₂R (Svizenska et al., 2013). Work of Leichsenring et al. analyzed the effect of repeated i.p. administration of the CB₂R agonist GW-405,833 on mechanical allodynia, compared with the potent cannabinoid receptor agonist WIN55,212-2 (Leichsenring, Andriske, Bäcker, Stichel, & Lübbert, 2009). Both drugs, applied daily at a low nonpsychotropic dose, were equally effective in reducing mechanical allodynia induced by SNL. A reappearance of glial activation was also associated with return of neuropathic pain-related behavior in this study (Leichsenring et al., 2009). The involvement of peripheral CB₂R in neuropathic pain symptoms alleviation was also a subject of studies by Kinsey, Mahadevan, et al. (2011) and Kinsey, Naidu, et al. (2011). An ethyl sulfonamide THC analogue, O-3223, a selective CB₂ agonist, was reported to reduce thermal hyperalgesia in the CCI-induced neuropathic pain model. Its antihyperalgesic effects were blocked by pretreatment with the CB₂R-selective antagonist SR144528, but not by the CB₁R antagonist, rimonabant. In addition, O-3223 (unlike CP-55,940, CB₁R and CB₂R agonist) did not elicit hypothermia or motor disturbances, indicating it has significant antiinflammatory and antinociceptive effects in vivo, but does not cause CB₁R-mediated side effects.

The therapeutic utility of locally administered AEA for neuropathic pain was proven by Guindon and Beaulieu (2006). However, surprising data on the lack of antiallodynic and antihyperalgesic effects of URB597 in a neuropathic pain model were published by Jayamanne et al. (2006). In animals subjected to partial ligation of the sciatic nerve, i.p. administration of the

selective FAAH inhibitor, URB597, produced no significant change in mechanical paw withdrawal latency. It has been suggested that repeated administration of URB597 may prove to be more efficacious in neuropathic pain models, as observed previously for exogenous cannabinoid receptor agonists (Costa et al., 2004). Moreover, acute administration of the irreversible FAAH inhibitor, URB597 and of the reversible FAAH inhibitor, OL-135, decreases allodynia in mouse CCI model of neuropathic pain (Kinsey et al., 2009). This attenuation was completely blocked by pretreatment with either CB₁ or CB₂R antagonists. Given the neuro-inflammatory nature of the nerve injury in the CCI model, it is not surprising that both cannabinoid receptors play a role in modulating neuropathic pain.

Another FAAH inhibitor, PF-3845, characterized by an increased FAAH specificity and longer duration of in vivo activity (Kinsey, Long, Cravatt, & Lichtman, 2010) also showed an attenuation of CCI-induced mechanical and cold allodynia in wild-type mice (Kinsey et al., 2009). Subsequent work from the Lichtman group explored the contribution of CB₁R and/or CB₂R for the antiallodynic effects of the FAAH and the MAGL inhibitors in a mouse model of neuropathic pain (Kinsey et al., 2010) and further confirmed that both CB₁ and CB₂R are necessary for the antiallodynic effects of FAAH inhibitors, while only CB₁R is necessary for the antiallodynic effects caused by MAGL inhibition. These data indicate that the endocannabinoids may affect different levels of the nociceptive and inflammatory pathways involved in neuropathic pain.



6. CONCLUSION

Cannabinoids exert a direct antinociceptive effect on pain of different origins. The CB₁R-mediated analgesic effects of cannabinoid ligands are well established, but limited by their side-effect profile. The observation that CB₂R activation produces desirable actions in a range of preclinical models (Han, Thatte, Buzard, & Jones, 2013; Leleu-Chavain et al., 2012) attracted considerable interest. However, despite very favorable efficacy in a range of preclinical models, CB₂ agonists have fared poorly in the clinic (Dhopeshwarkar & Mackie, 2014). The targeted manipulation of the endocannabinoid system might also be beneficial in the face of inflammation and chronic pain conditions. Interestingly investigations into the endocannabinoids and their effector sites, along with other noncannabinoid receptors, have exploded in recent years, and insights reveal this area of

pharmacology to be highly complex and dynamic (Piscitelli & Di Marzo, 2012; Starowicz & Di Marzo, 2013). Data derived from complex and clinically relevant animal models highlight the question of effectiveness of dual-acting compounds (Aiello, Carullo, Badolato, & Brizzi, 2016; Ligresti et al., 2014; Malek & Starowicz, 2016) and support the case for multitarget pharmacological intervention for effective pain treatment.

CONFLICT OF INTEREST

The authors have no conflicts of interest to declare.

ACKNOWLEDGMENTS

This work was supported by grants from the National Science Center, Poland (SONATA BIS NCN/2012/07/E/NZ7/01269 and OPUS NCN/2014/13/B/NZ7/02311) to K.S. and a grant from Science Foundation Ireland (10/IN.1/B2976) to D.P.F. The authors wish to thank Natalia Malek for her assistance in preparing the figure and table.

REFERENCES

- Ahluwalia, J., Urban, L., Bevan, S., Capogna, M., & Nagy, I. (2002). Cannabinoid 1 receptors are expressed by nerve growth factor- and glial cell-derived neurotrophic factor-responsive primary sensory neurones. *Neuroscience*, 110(4), 747–753.
- Aiello, F., Carullo, G., Badolato, M., & Brizzi, A. (2016). TRPV1-FAAH-COX: The couples game in pain treatment. *ChemMedChem*, 11(16), 1686–1694.
- Alexander, S. P., & Kendall, D. A. (2007). The complications of promiscuity: Endocannabinoid action and metabolism. *British Journal of Pharmacology*, 152(5), 602–623.
- Anand, U., Otto, W. R., Sanchez-Herrera, D., Facer, P., Yiangou, Y., Korchev, Y., et al. (2008). Cannabinoid receptor CB2 localisation and agonist-mediated inhibition of capsaicin responses in human sensory neurons. *Pain*, 138(3), 667–680.
- Ashton, J. C., Friberg, D., Darlington, C. L., & Smith, P. F. (2006). Expression of the cannabinoid CB2 receptor in the rat cerebellum: An immunohistochemical study. *Neuroscience Letters*, 396(2), 113–116.
- Baek, J. H., Zheng, Y., Darlington, C. L., & Smith, P. F. (2008). Cannabinoid CB2 receptor expression in the rat brainstem cochlear and vestibular nuclei. *Acta Oto-Laryngologica*, 128(9), 961–967.
- Barnes, M. P. (2006). Sativex: Clinical efficacy and tolerability in the treatment of symptoms of multiple sclerosis and neuropathic pain. *Expert Opinion on Pharmacotherapy*, 7(5), 607–615.
- Beltramo, M., Bernardini, N., Bertorelli, R., Campanella, M., Nicolussi, E., Fredduzzi, S., et al. (2006). CB2 receptor-mediated antihyperalgesia: Possible direct involvement of neural mechanisms. *The European Journal of Neuroscience*, 23(6), 1530–1538.
- Berdyshev, E. V. (2000). Cannabinoid receptors and the regulation of immune response. *Chemistry and Physics of Lipids*, 108(1–2), 169–190.
- Bicher, H. I., & Mechoulam, R. (1968). Pharmacological effects of two active constituents of marijuana. *Archives Internationales de Pharmacodynamie et de Thérapie*, 172, 24–31.
- Bjursell, M., Ryberg, E., Wu, T., Greasley, P. J., Bohlooly, Y. M., & Hjorth, S. (2016). Deletion of Gpr55 results in subtle effects on energy metabolism, motor activity and thermal pain sensation. *PLoS One*, 11(12), e0167965.

- Bloom, A. S., & Dewey, W. L. (1978). A comparison of some pharmacological actions of morphine and delta9-tetrahydrocannabinol in the mouse. *Psychopharmacology*, 57(3), 243–248.
- Boychuk, D. G., Goddard, G., Mauro, G., & Orellana, M. F. (2015). The effectiveness of cannabinoids in the management of chronic nonmalignant neuropathic pain: A systematic review. *Journal of Oral & Facial Pain and Headache*, 29(1), 7–14.
- Brownjohn, P. W., & Ashton, J. C. (2012). Spinal cannabinoid CB2 receptors as a target for neuropathic pain: An investigation using chronic constriction injury. *Neuroscience*, 203, 180–193.
- Brusco, A., Tagliaferro, P. A., Saez, T., & Onaivi, E. S. (2008). Ultrastructural localization of neuronal brain CB2 cannabinoid receptors. *Annals of the New York Academy of Sciences*, 1139, 450–457.
- Buckley, N. E. (2008). The peripheral cannabinoid receptor knockout mice: An update. *British Journal of Pharmacology*, 153(2), 309–318.
- Buckley, N. E., McCoy, K. L., Mezey, E., Bonner, T., Zimmer, A., Felder, C. C., et al. (2000). Immunomodulation by cannabinoids is absent in mice deficient for the cannabinoid CB(2) receptor. *European Journal of Pharmacology*, 396(2–3), 141–149.
- Calignano, A., La Rana, G., Giuffrida, A., & Piomelli, D. (1998). Control of pain initiation by endogenous cannabinoids. *Nature*, 394(6690), 277–281.
- Canadian Agency for Drugs and Technologies in Health. (2016). *Cannabinoid buccal spray for chronic non-cancer or neuropathic pain: A review of clinical effectiveness, safety, and guidelines*. Ottawa, ON.
- Carey, L. M., Gutierrez, T., Deng, L., Lee, W. H., Mackie, K., & Hohmann, A. G. (2017). Inflammatory and neuropathic nociception is preserved in GPR55 knockout mice. *Scientific Reports*, 7(1), 944. <http://dx.doi.org/10.1038/s41598-017-01062-2>.
- Carey, L. M., Slivicki, R. A., Leishman, E., Cornett, B., Mackie, K., Bradshaw, H., et al. (2016). A pro-nociceptive phenotype unmasked in mice lacking fatty-acid amide hydrolase. *Molecular Pain*, 12, 1–23. <http://dx.doi.org/10.1177/1744806916649192>.
- Castane, A., Celerier, E., Martin, M., Ledent, C., Parmentier, M., Maldonado, R., et al. (2006). Development and expression of neuropathic pain in CB1 knockout mice. *Neuropharmacology*, 50(1), 111–122.
- Concannon, R. M., Okine, B. N., Finn, D. P., & Dowd, E. (2015). Differential upregulation of the cannabinoid CB(2) receptor in neurotoxic and inflammation-driven rat models of Parkinson's disease. *Experimental Neurology*, 269, 133–141.
- Connell, K., Bolton, N., Olsen, D., Piomelli, D., & Hohmann, A. G. (2006). Role of the basolateral nucleus of the amygdala in endocannabinoid-mediated stress-induced analgesia. *Neuroscience Letters*, 397(3), 180–184.
- Corcoran, L., Roche, M., & Finn, D. P. (2015). The role of the brain's endocannabinoid system in pain and its modulation by stress. *International Review of Neurobiology*, 125, 203–255.
- Costa, B., Colleoni, M., Conti, S., Trovato, A. E., Bianchi, M., Sotgiu, M. L., et al. (2004). Repeated treatment with the synthetic cannabinoid WIN 55,212-2 reduces both hyperalgesia and production of pronociceptive mediators in a rat model of neuropathic pain. *British Journal of Pharmacology*, 141(1), 4–8.
- Costa, B., Trovato, A. E., Colleoni, M., Giagnoni, G., Zarini, E., & Croci, T. (2005). Effect of the cannabinoid CB1 receptor antagonist, SR141716, on nociceptive response and nerve demyelination in rodents with chronic constriction injury of the sciatic nerve. *Pain*, 116(1–2), 52–61.
- Cravatt, B. F., Demarest, K., Patricelli, M. P., Bracey, M. H., Giang, D. K., Martin, B. R., et al. (2001). Supersensitivity to anandamide and enhanced endogenous cannabinoid signaling in mice lacking fatty acid amide hydrolase. *Proceedings of the National Academy of Sciences of the United States of America*, 98(16), 9371–9376.

- de Novellis, V., Palazzo, E., Rossi, F., De Petrocellis, L., Petrosino, S., Guida, F., et al. (2008). The analgesic effect of N-arachidonoyl-serotonin, a FAAH inhibitor and TRPV1 receptor antagonist, associated with changes in rostral ventromedial medulla and locus coeruleus cell activity in rats. *Neuropharmacology*, 55(7), 1105–1113.
- de Novellis, V., Vita, D., Gatta, L., Luongo, L., Bellini, G., De Chiaro, M., et al. (2011). The blockade of the transient receptor potential vanilloid type 1 and fatty acid amide hydrolase decreases symptoms and central sequelae in the medial prefrontal cortex of neuropathic rats. *Molecular Pain*, 7, 7.
- De Vry, J., Jentzsch, K. R., Kuhl, E., & Eckel, G. (2004). Behavioral effects of cannabinoids show differential sensitivity to cannabinoid receptor blockade and tolerance development. *Behavioural Pharmacology*, 15(1), 1–12.
- Deliu, E., Sperow, M., Console-Bram, L., Carter, R. L., Tilley, D. G., Kalamarides, D. J., et al. (2015). The lysophosphatidylinositol receptor GPR55 modulates pain perception in the periaqueductal gray. *Molecular Pharmacology*, 88(2), 265–272.
- Deng, L., Guindon, J., Cornett, B. L., Makriyannis, A., Mackie, K., & Hohmann, A. G. (2015). Chronic cannabinoid receptor 2 activation reverses paclitaxel neuropathy without tolerance or cannabinoid receptor 1-dependent withdrawal. *Biological Psychiatry*, 77(5), 475–487.
- Derbenev, A. V., Stuart, T. C., & Smith, B. N. (2004). Cannabinoids suppress synaptic input to neurones of the rat dorsal motor nucleus of the vagus nerve. *The Journal of Physiology*, 559(Pt. 3), 923–938.
- Devane, W. A., Dysarz, F. A., 3rd, Johnson, M. R., Melvin, L. S., & Howlett, A. C. (1988). Determination and characterization of a cannabinoid receptor in rat brain. *Molecular Pharmacology*, 34(5), 605–613.
- Devane, W. A., Hanus, L., Breuer, A., Pertwee, R. G., Stevenson, L. A., Griffin, G., et al. (1992). Isolation and structure of a brain constituent that binds to the cannabinoid receptor. *Science*, 258(5090), 1946–1949.
- Dhopeswarkar, A., & Mackie, K. (2014). CB2 cannabinoid receptors as a therapeutic target—What does the future hold? *Molecular Pharmacology*, 86, 430–437.
- Di Marzo, V., Breivogel, C. S., Tao, Q., Bridgen, D. T., Razdan, R. K., Zimmer, A. M., et al. (2000). Levels, metabolism, and pharmacological activity of anandamide in CB(1) cannabinoid receptor knockout mice: Evidence for non-CB(1), non-CB(2) receptor-mediated actions of anandamide in mouse brain. *Journal of Neurochemistry*, 75(6), 2434–2444.
- Egertova, M., Giang, D. K., Cravatt, B. F., & Elphick, M. R. (1998). A new perspective on cannabinoid signalling: Complementary localization of fatty acid amide hydrolase and the CB1 receptor in rat brain. *Proceedings. Biological Sciences*, 265(1410), 2081–2085.
- Elmes, S. J. R., Jhaveri, M. D., Smart, D., Kendall, D. A., & Chapman, V. (2004). Cannabinoid CB2 receptor activation inhibits mechanically evoked responses of wide dynamic range dorsal horn neurons in naïve rats and in rat models of inflammatory and neuropathic pain. *European Journal of Neuroscience*, 20(9), 2311–2320.
- Escobar, W., Ramirez, K., Avila, C., Limongi, R., Vanegas, H., & Vazquez, E. (2012). Metamizol, a non-opioid analgesic, acts via endocannabinoids in the PAG-RVM axis during inflammation in rats. *European Journal of Pain*, 16(5), 676–689.
- Facci, L., Dal Toso, R., Romanello, S., Buriani, A., Skaper, S. D., & Leon, A. (1995). Mast cells express a peripheral cannabinoid receptor with differential sensitivity to anandamide and palmitoylethanolamide. *Proceedings of the National Academy of Sciences of the United States of America*, 92(8), 3376–3380.
- Fang, Q., Han, Z. L., Li, N., Wang, Z. L., He, N., & Wang, R. (2012). Effects of neuropeptide FF system on CB(1) and CB(2) receptors mediated antinociception in mice. *Neuropharmacology*, 62(2), 855–864.

- Farquhar-Smith, W. P., Egertova, M., Bradbury, E. J., McMahon, S. B., Rice, A. S., & Elphick, M. R. (2000). Cannabinoid CB(1) receptor expression in rat spinal cord. *Molecular and Cellular Neurosciences*, 15(6), 510–521.
- Felder, C. C., Nielsen, A., Briley, E. M., Palkovits, M., Priller, J., Axelrod, J., et al. (1996). Isolation and measurement of the endogenous cannabinoid receptor agonist, anandamide, in brain and peripheral tissues of human and rat. *FEBS Letters*, 393(2–3), 231–235.
- Finn, D. P., & Chapman, V. (2004). Cannabinoids as analgesic agents: Evidence from in vivo studies. *Current Neuropharmacology*, 2(1), 75–89.
- Finn, D. P., Jhaveri, M. D., Beckett, S. R., Kendall, D. A., Marsden, C. A., & Chapman, V. (2004). Cannabinoids modulate ultrasound-induced aversive responses in rats. *Psychopharmacology*, 172(1), 41–51.
- Finn, D. P., Jhaveri, M. D., Beckett, S. R., Roe, C. H., Kendall, D. A., Marsden, C. A., et al. (2003). Effects of direct periaqueductal grey administration of a cannabinoid receptor agonist on nociceptive and aversive responses in rats. *Neuropharmacology*, 45(5), 594–604.
- Ford, G. K., Kieran, S., Dolan, K., Harhen, B., & Finn, D. P. (2011). A role for the ventral hippocampal endocannabinoid system in fear-conditioned analgesia and fear responding in the presence of nociceptive tone in rats. *Pain*, 152(11), 2495–2504.
- Fox, A., Kessingland, A., Gentry, C., McNair, K., Patel, S., Urban, L., et al. (2001). The role of central and peripheral cannabinoid 1 receptors in the antihyperalgesic activity of cannabinoids in a model of neuropathic pain. *Pain*, 92(1–2), 91–100.
- Freitas, R. L., Salgado-Rohner, C. J., Hallak, J. E., Crippa, J. A., & Coimbra, N. C. (2013). Involvement of prelimbic medial prefrontal cortex in panic-like elaborated defensive behaviour and innate fear-induced antinociception elicited by GABAA receptor blockade in the dorsomedial and ventromedial hypothalamic nuclei: Role of the endocannabinoid CB1 receptor. *The International Journal of Neuropsychopharmacology*, 16(8), 1781–1798.
- Giordano, C., Cristino, L., Luongo, L., Siniscalco, D., Petrosino, S., Piscitelli, F., et al. (2012). TRPV1-dependent and -independent alterations in the limbic cortex of neuropathic mice: Impact on glial caspases and pain perception. *Cerebral Cortex*, 22(11), 2495–2518.
- Glass, M., Dragunow, M., & Faull, R. L. M. (1997). Cannabinoid receptors in the human brain: A detailed anatomical and quantitative autoradiographic study in the fetal, neonatal and adult human brain. *Neuroscience*, 77(2), 299–318.
- Gong, J. P., Onaivi, E. S., Ishiguro, H., Liu, Q. R., Tagliaferro, P. A., Brusco, A., et al. (2006). Cannabinoid CB2 receptors: Immunohistochemical localization in rat brain. *Brain Research*, 1071(1), 10–23.
- Gonzalez, S., Cebeira, M., & Fernandez-Ruiz, J. (2005). Cannabinoid tolerance and dependence: A review of studies in laboratory animals. *Pharmacology, Biochemistry, and Behavior*, 81(2), 300–318.
- Griffin, G., Wray, E. J., Tao, Q., McAllister, S. D., Rorrer, W. K., Aung, M. M., et al. (1999). Evaluation of the cannabinoid CB2 receptor-selective antagonist, SR144528: Further evidence for cannabinoid CB2 receptor absence in the rat central nervous system. *European Journal of Pharmacology*, 377(1), 117–125.
- Grinspoon, L., & Bakalar, J. B. (1993). *Marijuana, the forbidden medicine*. New Haven: Yale University Press.
- Guindon, J., & Beaulieu, P. (2006). Antihyperalgesic effects of local injections of anandamide, ibuprofen, rofecoxib and their combinations in a model of neuropathic pain. *Neuropharmacology*, 50(7), 814–823.
- Guindon, J., Guijarro, A., Piomelli, D., & Hohmann, A. G. (2011). Peripheral antinociceptive effects of inhibitors of monoacylglycerol lipase in a rat model of inflammatory pain. *British Journal of Pharmacology*, 163(7), 1464–1478.

- Han, S., Thatte, J., Buzard, D. J., & Jones, R. M. (2013). Therapeutic utility of cannabinoid receptor type 2 (CB2) selective agonists. *Journal of Medicinal Chemistry*, 56, 8224–8256.
- Hanus, L., Abu-Lafi, S., Fride, E., Breuer, A., Vogel, Z., Shalev, D. E., et al. (2001). 2-Arachidonoyl glyceryl ether, an endogenous agonist of the cannabinoid CB1 receptor. *Proceedings of the National Academy of Sciences of the United States of America*, 98(7), 3662–3665.
- Hasanein, P., Parviz, M., Keshavarz, M., & Javanmardi, K. (2007). CB1 receptor activation in the basolateral amygdala produces antinociception in animal models of acute and tonic nociception. *Clinical and Experimental Pharmacology & Physiology*, 34(5–6), 439–449.
- Herkenham, M., Lynn, A. B., Johnson, M. R., Melvin, L. S., de Costa, B. R., & Rice, K. C. (1991). Characterization and localization of cannabinoid receptors in rat brain: A quantitative in vitro autoradiographic study. *The Journal of Neuroscience*, 11(2), 563–583.
- Herkenham, M., Lynn, A. B., Little, M. D., Johnson, M. R., Melvin, L. S., de Costa, B. R., et al. (1990). Cannabinoid receptor localization in brain. *Proceedings of the National Academy of Sciences of the United States of America*, 87(5), 1932–1936.
- Hohmann, A. G., Briley, E. M., & Herkenham, M. (1999a). Pre- and postsynaptic distribution of cannabinoid and mu opioid receptors in rat spinal cord. *Brain Research*, 822(1–2), 17–25.
- Hohmann, A. G., & Herkenham, M. (1998). Regulation of cannabinoid and mu opioid receptors in rat lumbar spinal cord following neonatal capsaicin treatment. *Neuroscience Letters*, 252(1), 13–16.
- Hohmann, A. G., & Herkenham, M. (1999a). Cannabinoid receptors undergo axonal flow in sensory nerves. *Neuroscience*, 92(4), 1171–1175.
- Hohmann, A. G., & Herkenham, M. (1999b). Localization of central cannabinoid CB1 receptor messenger RNA in neuronal subpopulations of rat dorsal root ganglia: A double-label in situ hybridization study. *Neuroscience*, 90(3), 923–931.
- Hohmann, A. G., Suplita, R. L., Bolton, N. M., Neely, M. H., Fegley, D., Mangieri, R., et al. (2005). An endocannabinoid mechanism for stress-induced analgesia. *Nature*, 435(7045), 1108–1112.
- Hohmann, A. G., Tsou, K., & Michael Walker, J. (1998). Cannabinoid modulation of wide dynamic range neurons in the lumbar dorsal horn of the rat by spinally administered WIN55,212-2. *Neuroscience Letters*, 257(3), 119–122.
- Hohmann, A. G., Tsou, K., & Walker, J. M. (1999a). Cannabinoid suppression of noxious heat-evoked activity in wide dynamic range neurons in the lumbar dorsal horn of the rat. *Journal of Neurophysiology*, 81(2), 575–583.
- Hohmann, A. G., Tsou, K., & Walker, J. M. (1999b). Intrathecal cannabinoid administration suppresses noxious stimulus-evoked Fos protein-like immunoreactivity in rat spinal cord: Comparison with morphine. *Zhongguo Yao Li Xue Bao*, 20(12), 1132–1136.
- Holt, S., Comelli, F., Costa, B., & Fowler, C. J. (2005). Inhibitors of fatty acid amide hydro-lase reduce carrageenan-induced hind paw inflammation in pentobarbital-treated mice: Comparison with indomethacin and possible involvement of cannabinoid receptors. *British Journal of Pharmacology*, 146(3), 467–476.
- Howlett, A. C., Barth, F., Bonner, T. I., Cabral, G., Casellas, P., Devane, W. A., et al. (2002). International union of pharmacology. XXVII. Classification of cannabinoid receptors. *Pharmacological Reviews*, 54(2), 161–202.
- Hsieh, G. C., Pai, M., Chandran, P., Hooker, B. A., Zhu, C. Z., Salyers, A. K., et al. (2011). Central and peripheral sites of action for CB(2) receptor mediated analgesic activity in chronic inflammatory and neuropathic pain models in rats. *British Journal of Pharmacology*, 162(2), 428–440.
- Huang, S. M., Bisogno, T., Trevisani, M., Al-Hayani, A., De Petrocellis, L., Fezza, F., et al. (2002). An endogenous capsaicin-like substance with high potency at recombinant and

- native vanilloid VR1 receptors. *Proceedings of the National Academy of Sciences of the United States of America*, 99(12), 8400–8405.
- Ibrahim, M. M., Porreca, F., Lai, J., Albrecht, P. J., Rice, F. L., Khodorova, A., et al. (2005). CB2 cannabinoid receptor activation produces antinociception by stimulating peripheral release of endogenous opioids. *Proceedings of the National Academy of Sciences of the United States of America*, 102(8), 3093–3098.
- Iskedjian, M., Bereza, B., Gordon, A., Piwko, C., & Einarson, T. R. (2007). Meta-analysis of cannabis based treatments for neuropathic and multiple sclerosis-related pain. *Current Medical Research and Opinion*, 23(1), 17–24.
- Iwamura, H., Suzuki, H., Ueda, Y., Kaya, T., & Inaba, T. (2001). In vitro and in vivo pharmacological characterization of JTE-907, a novel selective ligand for cannabinoid CB2 receptor. *The Journal of Pharmacology and Experimental Therapeutics*, 296(2), 420–425.
- Jayamanne, A., Greenwood, R., Mitchell, V. A., Aslan, S., Piomelli, D., & Vaughan, C. W. (2006). Actions of the FAAH inhibitor URB597 in neuropathic and inflammatory chronic pain models. *British Journal of Pharmacology*, 147(3), 281–288.
- Jhaveri, M. D., Elmes, S. J., Richardson, D., Barrett, D. A., Kendall, D. A., Mason, R., et al. (2008). Evidence for a novel functional role of cannabinoid CB(2) receptors in the thalamus of neuropathic rats. *The European Journal of Neuroscience*, 27(7), 1722–1730. <http://dx.doi.org/10.1111/j.1460-9568.2008.06162.x>.
- Ji, G., & Neugebauer, V. (2014). CB1 augments mGluR5 function in medial prefrontal cortical neurons to inhibit amygdala hyperactivity in an arthritis pain model. *The European Journal of Neuroscience*, 39(3), 455–466.
- Johanek, L. M., Simone, D. A., & Lisa, M. (2005). Cannabinoid agonist, CP 55, 940, prevents capsaicin-induced sensitization of spinal cord dorsal horn neurons. *Spinal Cord*, 93, 989–997.
- Kathuria, S., Gaetani, S., Fegley, D., Valiño, F., Duranti, A., Tontini, A., et al. (2003). Modulation of anxiety through blockade of anandamide hydrolysis. *Nature Medicine*, 9(1), 76–81.
- Kinsey, S. G., Long, J. Z., Cravatt, B. F., & Lichtman, A. H. (2010). Fatty acid amide hydrolase and monoacylglycerol lipase inhibitors produce anti-allodynic effects in mice through distinct cannabinoid receptor mechanisms. *Journal of Pain*, 11(12), 1420–1428.
- Kinsey, S. G., Long, J. Z., O'Neal, S. T., Abdullah, R. A., Poklis, J. L., Boger, D. L., et al. (2009). Blockade of endocannabinoid-degrading enzymes attenuates neuropathic pain. *The Journal of Pharmacology and Experimental Therapeutics*, 330(3), 902–910.
- Kinsey, S. G., Mahadevan, A., Zhao, B., Sun, H., Naidu, P. S., Razdan, R. K., et al. (2011a). The CB2 cannabinoid receptor-selective agonist O-3223 reduces pain and inflammation without apparent cannabinoid behavioral effects. *Neuropharmacology*, 60(2–3), 244–251.
- Kinsey, S. G., Naidu, P. S., Cravatt, B. F., Dudley, D. T., & Lichtman, A. H. (2011b). Fatty acid amide hydrolase blockade attenuates the development of collagen-induced arthritis and related thermal hyperalgesia in mice. *Pharmacology, Biochemistry, and Behavior*, 99(4), 718–725.
- Klein, T. W., Newton, C. A., & Friedman, H. (2001). Cannabinoids and the immune system. *Pain Research & Management*, 6(2), 95–101.
- La Porta, C., Bura, S. A., Aracil-Fernandez, A., Manzanares, J., & Maldonado, R. (2013). Role of CB1 and CB2 cannabinoid receptors in the development of joint pain induced by monosodium iodoacetate. *Pain*, 154(1), 160–174.
- La Porta, C., Bura, A. S., Llorente-Onaíndia, J., Pastor, A., Navarrete, F., García-Gutiérrez, M. S., et al. (2015). Role of the endocannabinoid system in the emotional manifestations of osteoarthritis pain. *Pain*, 156(1), 2001–2012.
- Ledent, C., Valverde, O., Cossu, G., Petitot, F., Aubert, J. F., Beslot, F., et al. (1999). Unresponsiveness to cannabinoids and reduced addictive effects of opiates in CB1 receptor knockout mice. *Science*, 283, 401–404.

- Lee, M. C., Ploner, M., Wiech, K., Bingel, U., Wanigasekera, V., Brooks, J., et al. (2013). Amygdala activity contributes to the dissociative effect of cannabis on pain perception. *Pain*, 154(1), 124–134.
- Leichsenring, A., Andriske, M., Bäcker, I., Stichel, C. C., & Lübbert, H. (2009). Analgesic and antiinflammatory effects of cannabinoid receptor agonists in a rat model of neuropathic pain. *Naunyn-Schmiedeberg's Archives of Pharmacology*, 379(6), 627–636.
- Leleu-Chavain, N., Body-Malapel, M., Spencer, J., Chavatte, P., Desreumaux, P., & Millet, R. (2012). Recent advances in the development of selective CB2 agonists as promising anti-inflammatory agents. *Current Medicinal Chemistry*, 19(21), 3457–3474.
- Li, M. H., Suchland, K. L., & Ingram, S. L. (2017). Compensatory activation of cannabinoid CB2 receptor inhibition of GABA release in the rostral ventromedial medulla in inflammatory pain. *The Journal of Neuroscience*, 37(3), 626–636.
- Lichtman, A. H., Cook, S. A., & Martin, B. R. (1996). Investigation of brain sites mediating cannabinoid-induced antinociception in rats: Evidence supporting periaqueductal gray involvement. *The Journal of Pharmacology and Experimental Therapeutics*, 276(2), 585–593.
- Lichtman, A. H., & Martin, B. R. (1991). Spinal and supraspinal components of cannabinoid-induced antinociception. *The Journal of Pharmacology and Experimental Therapeutics*, 258(2), 517–523.
- Lichtman, A. H., & Martin, B. R. (1997). The selective cannabinoid antagonist SR 141716A blocks cannabinoid-induced antinociception in rats. *Pharmacology, Biochemistry, and Behavior*, 57(1–2), 7–12.
- Lichtman, A. H., Shelton, C. C., Advani, T., & Cravatt, B. F. (2004). Mice lacking fatty acid amide hydrolase exhibit a cannabinoid receptor-mediated phenotypic hypoalgesia. *Pain*, 109(3), 319–327.
- Ligresti, A., Martos, J., Wang, J., Guida, F., Allarà, M., Palmieri, V., et al. (2014). Prostamide F2 α receptor antagonism combined with inhibition of FAAH may block the pro-inflammatory mediators formed following selective FAAH inhibition. *British Journal of Pharmacology*, 171(6), 1408–1419.
- Lim, G., Sung, B., Ji, R. R., & Mao, J. (2003). Upregulation of spinal cannabinoid-1-receptors following nerve injury enhances the effects of Win 55,212-2 on neuropathic pain behaviors in rats. *Pain*, 105(1–2), 275–283.
- Luongo, L., Palazzo, E., Tambaro, S., Giordano, C., Gatta, L., Scafuro, M. A., et al. (2010). 1-(2',4'-Dichlorophenyl)-6-methyl-N-cyclohexylamine-1,4-dihydroindeno[1,2-c]pyrazole-3-carboxamide, a novel CB2 agonist, alleviates neuropathic pain through functional microglial changes in mice. *Neurobiology of Disease*, 37(1), 177–185.
- Lynch, M. E., & Ware, M. A. (2015). Cannabinoids for the treatment of chronic non-cancer pain: An updated systematic review of randomized controlled trials. *Journal of Neuroimmune Pharmacology*, 10(2), 293–301.
- Madasu, M. K., Roche, M., & Finn, D. P. (2015). Supraspinal TRPV1 in pain and psychiatric disorders. *Modern Trends in Pharmacopsychiatry*, 30, 80–93.
- Mailleux, P., Parmentier, M., & Vanderhaeghen, J. J. (1992). Distribution of cannabinoid receptor messenger RNA in the human brain: An in situ hybridization histochemistry with oligonucleotides. *Neuroscience Letters*, 143(1–2), 200–204.
- Maione, S., Bisogno, T., de Novellis, V., Palazzo, E., Cristino, L., Valenti, M., et al. (2006). Elevation of endocannabinoid levels in the ventrolateral periaqueductal grey through inhibition of fatty acid amide hydrolase affects descending nociceptive pathways via both cannabinoid receptor type 1 and transient receptor potential vanilloid type-1 receptors. *The Journal of Pharmacology and Experimental Therapeutics*, 316(3), 969–982.
- Malan, T. P., Jr., Ibrahim, M. M., Deng, H., Liu, Q., Mata, H. P., Vanderah, T., et al. (2001). CB2 cannabinoid receptor-mediated peripheral antinociception. *Pain*, 93(3), 239–245.

- Malek, N., & Starowicz, K. (2016). Dual-acting compounds targeting endocannabinoid and endovanilloid systems—A novel treatment option for chronic pain management. *Frontiers in Pharmacology*, 7, 257.
- Manning, B. H., Martin, W. J., & Meng, I. D. (2003). The rodent amygdala contributes to the production of cannabinoid-induced antinociception. *Neuroscience*, 120(4), 1157–1170.
- Manning, B. H., Merin, N. M., Meng, I. D., & Amaral, D. G. (2001). Reduction in opioid- and cannabinoid-induced antinociception in rhesus monkeys after bilateral lesions of the amygdaloid complex. *Journal of Neuroscience*, 21(20), 8238–8246.
- Martin, W. J., Coffin, P. O., Attias, E., Balinsky, M., Tsou, K., & Walker, J. M. (1999a). Anatomical basis for cannabinoid-induced antinociception as revealed by intracerebral microinjections. *Brain Research*, 822(1–2), 237–242.
- Martin, W. J., Lai, N. K., Patrick, S. L., Tsou, K., & Walker, J. M. (1993). Antinociceptive actions of cannabinoids following intraventricular administration in rats. *Brain Research*, 629(2), 300–304.
- Martin, W. J., Loo, C. M., & Basbaum, A. I. (1999b). Spinal cannabinoids are anti-allodynic in rats with persistent inflammation. *Pain*, 82(2), 199–205.
- Martin, W. J., Patrick, S. L., Coffin, P. O., Tsou, K., & Walker, J. M. (1995). An examination of the central sites of action of cannabinoid-induced antinociception in the rat. *Life Sciences*, 56(23–24), 2103–2109.
- Martin, W. J., Tsou, K., & Walker, J. M. (1998). Cannabinoid receptor-mediated inhibition of the rat tail-flick reflex after microinjection into the rostral ventromedial medulla. *Neuroscience Letters*, 242(1), 33–36.
- Matsuda, L. A., Lolait, S. J., Brownstein, M. J., Young, A. C., & Bonner, T. I. (1990). Structure of a cannabinoid receptor and functional expression of the cloned cDNA. *Nature*, 346(6284), 561–564.
- Mazzari, S., Canella, R., Petrelli, L., Marcolongo, G., & Leon, A. (1996). N-(2-hydroxyethyl) hexadecanamide is orally active in reducing edema formation and inflammatory hyperalgesia by down-modulating mast cell activation. *European Journal of Pharmacology*, 300(3), 227–236.
- McCormick, M., et al. (2017). *The health effects of cannabis and cannabinoids: The current state of evidence and recommendations for research*. Washington, DC: The National Academies Press.
- McGaraughty, S., Chu, K. L., Bitner, R. S., Martino, B., El Kouhen, R., Han, P., et al. (2003). Capsaicin infused into the PAG affects rat tail flick responses to noxious heat and alters neuronal firing in the RVM. *Journal of Neurophysiology*, 90(4), 2702–2710.
- Mechoulam, R., Ben-Shabat, S., Hanus, L., Ligumsky, M., Kaminski, N. E., Schatz, A. R., et al. (1995). Identification of an endogenous 2-monoglyceride, present in canine gut, that binds to cannabinoid receptors. *Biochemical Pharmacology*, 50(1), 83–90.
- Mechoulam, R., & Gaoni, Y. (1967). Isolation, structure and partial synthesis of an active constituent of hashish. *Journal of the American Chemical Society*, 86, 1646–1647.
- Meng, I. D., & Johansen, J. P. (2004). Antinociception and modulation of rostral ventromedial medulla neuronal activity by local microinfusion of a cannabinoid receptor agonist. *Neuroscience*, 124(3), 685–693.
- Meng, I. D., Manning, B. H., Martin, W. J., & Fields, H. L. (1998). An analgesia circuit activated by cannabinoids. *Nature*, 395(6700), 381–383.
- Millan, M. J. (2002). Descending control of pain. *Progress in Neurobiology*, 66(6), 355–474.
- Milligan, E. D., Twining, C., Chacur, M., Biedenkapp, J., O'Connor, K., Poole, S., et al. (2003). Spinal glia and proinflammatory cytokines mediate mirror-image neuropathic pain in rats. *The Journal of Neuroscience*, 23(3), 1026–1040.
- Molina-Holgado, F., Rubio-Araiz, A., Garcia-Ovejero, D., Williams, R. J., Moore, J. D., Arevalo-Martin, A., et al. (2007). CB2 cannabinoid receptors promote mouse neural stem cell proliferation. *The European Journal of Neuroscience*, 25(3), 629–634.

- Monhemius, R., Azami, J., Green, D. L., & Roberts, M. H. (2001). CB1 receptor mediated analgesia from the nucleus reticularis gigantocellularis pars alpha is activated in an animal model of neuropathic pain. *Brain Research*, 908(1), 67–74.
- Munro, S., Thomas, K. L., & Abu-Shaar, M. (1993). Molecular characterization of a peripheral receptor for cannabinoids. *Nature*, 365(6441), 61–65.
- Nackley, A. G., Makriyannis, A., & Hohmann, A. G. (2003). Selective activation of cannabinoid CB2 receptors suppresses spinal Fos protein expression and pain behavior in a rat model of inflammation. *Neuroscience*, 119(3), 747–757.
- Neugebauer, V., Galhardo, V., Maione, S., & Mackey, S. C. (2009). Forebrain pain mechanisms. *Brain Research Reviews*, 60(1), 226–242.
- Neugebauer, V., Li, W., Bird, G. C., & Han, J. S. (2004). The amygdala and persistent pain. *The Neuroscientist*, 10(3), 221–234.
- Nozaki, C., Markert, A., & Zimmer, A. (2015). Inhibition of FAAH reduces nitroglycerin-induced migraine-like pain and trigeminal neuronal hyperactivity in mice. *European Neuropsychopharmacology*, 25(8), 1388–1396.
- Okine, B. N., Madasu, M. K., McGowan, F., Prendergast, C., Gaspar, J. C., Harhen, B., et al. (2016). N-palmitoylethanolamide in the anterior cingulate cortex attenuates inflammatory pain behaviour indirectly via a CB1 receptor-mediated mechanism. *Pain*, 157(12), 2687–2696.
- Okine, B. N., Rea, K., Olango, W. M., Price, J., Herdman, S., Madasu, M. K., et al. (2014). A role for PPARalpha in the medial prefrontal cortex in formalin-evoked nociceptive responding in rats. *British Journal of Pharmacology*, 171(6), 1462–1471.
- Olango, W. M., Roche, M., Ford, G. K., Harhen, B., & Finn, D. P. (2012). The endocannabinoid system in the rat dorsolateral periaqueductal grey mediates fear-conditioned analgesia and controls fear expression in the presence of nociceptive tone. *British Journal of Pharmacology*, 165(8), 2549–2560.
- Onaivi, E. S., Ishiguro, H., Gong, J. P., Patel, S., Meozzi, P. A., Myers, L., et al. (2008). Brain neuronal CB2 cannabinoid receptors in drug abuse and depression: From mice to human subjects. *PLoS One*, 3(2), e1640.
- Onaivi, E. S., Ishiguro, H., Gong, J. P., Patel, S., Perchuk, A., Meozzi, P. A., et al. (2006a). Discovery of the presence and functional expression of cannabinoid CB2 receptors in brain. *Annals of the New York Academy of Sciences*, 1074, 514–536.
- Onaivi, E. S., Ishiguro, H., Sejal, P., Meozzi, P. A., Myers, L., Tagliaferro, P., et al. (2006b). Methods to study the behavioral effects and expression of CB2 cannabinoid receptor and its gene transcripts in the chronic mild stress model of depression. *Methods in Molecular Medicine*, 123, 291–298.
- Ossipov, M. H., Morimura, K., & Porreca, F. (2014). Descending pain modulation and chronification of pain. *Current Opinion in Supportive and Palliative Care*, 8(2), 143–151.
- Palazuelos, J., Aguado, T., Egia, A., Mechoulam, R., Guzman, M., & Galve-Roperh, I. (2006). Non-psychoactive CB2 cannabinoid agonists stimulate neural progenitor proliferation. *The FASEB Journal*, 20(13), 2405–2407.
- Palazzo, E., de Novellis, V., Marabese, I., Cuomo, D., Rossi, F., Berrino, L., et al. (2002). Interaction between vanilloid and glutamate receptors in the central modulation of nociception. *European Journal of Pharmacology*, 439(1–3), 69–75.
- Palazzo, E., Marabese, I., de Novellis, V., Oliva, P., Rossi, F., Berrino, L., et al. (2001). Metabotropic and NMDA glutamate receptors participate in the cannabinoid-induced antinociception. *Neuropharmacology*, 40(3), 319–326.
- Pan, J. X., Wang, Z. L., Li, N., Han, Z. L., Li, X. H., Tang, H. H., et al. (2014). Analgesic tolerance and cross-tolerance to the cannabinoid receptors ligands hemopressin, VD-hemopressin(alpha) and WIN55,212-2 at the supraspinal level in mice. *Neuroscience Letters*, 578, 187–191.

- Pertwee, R. G. (2001). Cannabinoid receptors and pain. *Progress in Neurobiology*, 63(5), 569–611.
- Petrenko, A. B., Yamazaki, M., Sakimura, K., Kano, M., & Baba, H. (2014). Augmented tonic pain-related behavior in knockout mice lacking monoacylglycerol lipase, a major degrading enzyme for the endocannabinoid 2-arachidonoylglycerol. *Behavioural Brain Research*, 271, 51–58.
- Petrosino, S., Palazzo, E., de Novellis, V., Bisogno, T., Rossi, F., Maione, S., et al. (2007). Changes in spinal and supraspinal endocannabinoid levels in neuropathic rats. *Neuropharmacology*, 52(2), 415–422.
- Piscitelli, F., & Di Marzo, V. (2012). “Redundancy” of endocannabinoid inactivation: New challenges and opportunities for pain control. *ACS Chemical Neuroscience*, 3, 356–363.
- Porter, A. C., Sauer, J. M., Knierman, M. D., Becker, G. W., Berna, M. J., Bao, J., et al. (2002). Characterization of a novel endocannabinoid, virodhamine, with antagonist activity at the CB1 receptor. *The Journal of Pharmacology and Experimental Therapeutics*, 301(3), 1020–1024.
- Quartilho, A., Mata, H. P., Ibrahim, M. M., Vanderah, T. W., Porreca, F., Makriyannis, A., et al. (2003). Inhibition of inflammatory hyperalgesia by activation of peripheral CB2 cannabinoid receptors. *Anesthesiology*, 99(4), 955–960.
- Racz, I., Nadal, X., Alferink, J., Baños, J. E., Rehnelt, J., Martín, M., et al. (2008). Interferon- γ is a critical modulator of CB(2) cannabinoid receptor signaling during neuropathic pain. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, 28(46), 12136–12145.
- Racz, I., Nent, E., Erxlebe, E., & Zimmer, A. (2015). CB1 receptors modulate affective behaviour induced by neuropathic pain. *Brain Research Bulletin*, 114, 42–48.
- Raffa, R. B., Stone, D. J., Jr., & Hipp, S. J. (1999). Differential cholera-toxin sensitivity of supraspinal antinociception induced by the cannabinoid agonists delta9-THC, WIN 55,212-2 and anandamide in mice. *Neuroscience Letters*, 263(1), 29–32.
- Rea, K., Olango, W. M., Harhen, B., Kerr, D. M., Galligan, R., Fitzgerald, S., et al. (2013). Evidence for a role of GABAergic and glutamatergic signalling in the basolateral amygdala in endocannabinoid-mediated fear-conditioned analgesia in rats. *Pain*, 154(4), 576–585.
- Rea, K., Olango, W. M., Okine, B. N., Madasu, M. K., McGuire, I. C., Coyle, K., et al. (2014). Impaired endocannabinoid signalling in the rostral ventromedial medulla underpins genotype-dependent hyper-responsivity to noxious stimuli. *Pain*, 155(1), 69–79.
- Rea, K., Roche, M., & Finn, D. P. (2007). Supraspinal modulation of pain by cannabinoids: The role of GABA and glutamate. *British Journal of Pharmacology*, 152(5), 633–648.
- Rice, A. S., Farquhar-Smith, W. P., & Nagy, I. (2002). Endocannabinoids and pain: Spinal and peripheral analgesia in inflammation and neuropathy. *Prostaglandins, Leukotrienes, and Essential Fatty Acids*, 66(2–3), 243–256.
- Richardson, J., Kilo, S., & Hargreaves, K. (1998). Cannabinoids reduce hyperalgesia and inflammation via interaction with peripheral CB1 receptors. *Pain*, 75(1), 111–119.
- Roche, M., & Finn, D. P. (2010). Brain CB(2) receptors: Implications for neuropsychiatric disorders. *Pharmaceuticals (Basel)*, 3(8), 2517–2553.
- Roche, M., Johnston, P., Mhuircheartaigh, O. N., Olango, W. M., Mackie, K., & Finn, D. P. (2010). Effects of intra-basolateral amygdala administration of rimonabant on nociceptive behaviour and neuronal activity in the presence or absence of contextual fear. *European Journal of Pain*, 14(5), 487–495.
- Roche, M., O'Connor, E., Diskin, C., & Finn, D. P. (2007). The effect of CB(1) receptor antagonism in the right basolateral amygdala on conditioned fear and associated analgesia in rats. *The European Journal of Neuroscience*, 26(9), 2643–2653.

- Romero-Sandoval, A., & Eisenach, J. C. (2007). Spinal cannabinoid receptor type 2 activation reduces hypersensitivity and spinal cord glial activation after paw incision. *Anesthesiology*, 106(4), 787–794.
- Romero-Sandoval, A., Natile-McMenemy, N., & DeLeo, J. A. (2008). Spinal microglial and perivascular cell cannabinoid receptor type 2 activation reduces behavioral hypersensitivity without tolerance after peripheral nerve injury. *Anesthesiology*, 108(4), 722–734.
- Ross, R. A., Coutts, A. A., McFarlane, S. M., Anavi-Goffer, S., Irving, A. J., Pertwee, R. G., et al. (2001). Actions of cannabinoid receptor ligands on rat cultured sensory neurones: Implications for antinociception. *Neuropharmacology*, 40(2), 221–232.
- Russo, E. B. (2016). Current therapeutic cannabis controversies and clinical trial design issues. *Frontiers in Pharmacology*, 7, 309.
- Sanudo-Pena, M. C., Strangman, N. M., Mackie, K., Walker, J. M., & Tsou, K. (1999). CB1 receptor localization in rat spinal cord and roots, dorsal root ganglion, and peripheral nerve. *Zhongguo Yao Li Xue Bao*, 20(12), 1115–1120.
- Schlosburg, J. E., Blankman, J. L., Long, J. Z., Nomura, D. K., Pan, B., Kinsey, S. G., et al. (2010). Chronic monoacylglycerol lipase blockade causes functional antagonism of the endocannabinoid system. *Nature Neuroscience*, 13(9), 1113–1119.
- Sideris, A., Piskoun, B., Russo, L., Norcini, M., Blanck, T., & Recio-Pinto, E. (2016). Cannabinoid 1 receptor knockout mice display cold allodynia, but enhanced recovery from spared-nerve injury-induced mechanical hypersensitivity. *Molecular Pain*, 12, 1–12. pii:1744806916649191.
- Siegling, A., Hofmann, H. A., Denzer, D., Mauler, F., & De Vry, J. (2001). Cannabinoid CB1 receptor upregulation in a rat model of chronic neuropathic pain. *European Journal of Pharmacology*, 415(1), R5–R7.
- Smith, F. L., Cichewicz, D., Martin, Z. L., & Welch, S. P. (1998). The enhancement of morphine antinociception in mice by delta9-tetrahydrocannabinol. *Pharmacology, Biochemistry, and Behavior*, 60(2), 559–566.
- Smith, P. B., & Martin, B. R. (1992). Spinal mechanisms of Δ^9 -tetrahydrocannabinol-induced analgesia. *Brain Research*, 578(1–2), 8–12.
- Sofia, R. D., Vassar, H. B., & Knobloch, L. C. (1975). Comparative analgesic activity of various naturally occurring cannabinoids in mice and rats. *Psychopharmacologia*, 40(4), 285–295.
- Sokal, D. M., Elmes, S. J. R., Kendall, D. A., & Chapman, V. (2003). Intraplantar injection of anandamide inhibits mechanically-evoked responses of spinal neurones via activation of CB2 receptors in anaesthetised rats. *Neuropharmacology*, 45(3), 404–411.
- Starowicz, K., & Di Marzo, V. (2013). Non-psychoactive analgesic drugs from the endocannabinoid system: “magic bullet” or “multiple-target” strategies? *European Journal of Pharmacology*, 716(1–3), 41–53.
- Starowicz, K., Maione, S., Cristino, L., Palazzo, E., Marabese, I., Rossi, F., et al. (2007). Tonic endovanilloid facilitation of glutamate release in brainstem descending antinociceptive pathways. *The Journal of Neuroscience*, 27(50), 13739–13749.
- Starowicz, K., Makuch, W., Korostynski, M., Malek, N., Slezak, M., Zychowska, M., et al. (2013). Full inhibition of spinal FAAH leads to TRPV1-mediated analgesic effects in neuropathic rats and possible lipoxygenase-mediated remodeling of anandamide metabolism. *PLoS One*, 8(4), e60040.
- Starowicz, K., Makuch, W., Osikowicz, M., Piscitelli, F., Petrosino, S., Di Marzo, V., et al. (2012). Spinal anandamide produces analgesia in neuropathic rats: Possible CB 1- and TRPV1-mediated mechanisms. *Neuropharmacology*, 62(4), 1746–1755.
- Staton, P. C., Hatcher, J. P., Walker, D. J., Morrison, A. D., Shapland, E. M., Hughes, J. P., et al. (2008). The putative cannabinoid receptor GPR55 plays a role in mechanical hyperalgesia associated with inflammatory and neuropathic pain. *Pain*, 139(1), 225–236.

- Stella, N., Schweitzer, P., & Piomelli, D. (1997). A second endogenous cannabinoid that modulates long-term potentiation. *Nature*, 388(6644), 773–778.
- Strangman, N. M., Walker, J. M., & Strangman, N. M. (1999). Cannabinoid WIN 55,212-2 inhibits the activity-dependent facilitation of spinal nociceptive responses. *Journal of Neurophysiology*, 82(1), 472–477.
- Suarez, J., Bermudez-Silva, F. J., Mackie, K., Ledent, C., Zimmer, A., Cravatt, B. F., et al. (2008). Immunohistochemical description of the endogenous cannabinoid system in the rat cerebellum and functionally related nuclei. *The Journal of Comparative Neurology*, 509(4), 400–421.
- Sugiura, T., Kondo, S., Sukagawa, A., Nakane, S., Shinoda, A., Itoh, K., et al. (1995). 2-Arachidonoylglycerol: A possible endogenous cannabinoid receptor ligand in brain. *Biochemical and Biophysical Research Communications*, 215(1), 89–97.
- Suplita, R. L., 2nd, Farthing, J. N., Gutierrez, T., & Hohmann, A. G. (2005). Inhibition of fatty-acid amide hydrolase enhances cannabinoid stress-induced analgesia: Sites of action in the dorsolateral periaqueductal gray and rostral ventromedial medulla. *Neuropharmacology*, 49(8), 1201–1209.
- Suzuki, R., & Dickenson, A. (2005). Spinal and supraspinal contributions to central sensitization in peripheral neuropathy. *Neurosignals*, 14(4), 175–181.
- Suzuki, R., Rygh, L. J., & Dickenson, A. H. (2004). Bad news from the brain: Descending 5-HT pathways that control spinal pain processing. *Trends in Pharmacological Sciences*, 25(12), 613–617.
- Svizenska, I. H., Brazda, V., Klusakova, I., & Dubovy, P. (2013). Bilateral changes of cannabinoid receptor type 2 protein and mRNA in the dorsal root ganglia of a rat neuropathic pain model. *The Journal of Histochemistry and Cytochemistry*, 61(7), 529–547.
- Thomas, B. F., Wei, X., & Martin, B. R. (1992). Characterization and autoradiographic localization of the cannabinoid binding site in rat brain using [³H]11-OH-delta 9-THC-DMH. *The Journal of Pharmacology and Experimental Therapeutics*, 263(3), 1383–1390.
- Thorat, S. N., & Bhargava, H. N. (1994). Evidence for a bidirectional cross-tolerance between morphine and delta 9-tetrahydrocannabinol in mice. *European Journal of Pharmacology*, 260(1), 5–13.
- Toniolo, E. F., Maique, E. T., Ferreira, W. A., Heimann, A. S., Ferro, E. S., Ramos-Ortolaza, D. L., et al. (2014). Hemopressin, an inverse agonist of cannabinoid receptors, inhibits neuropathic pain in rats. *Peptides*, 56, 125–131.
- Tsou, K., Brown, S., Sanudo-Pena, M. C., Mackie, K., & Walker, J. M. (1998a). Immunohistochemical distribution of cannabinoid CB1 receptors in the rat central nervous system. *Neuroscience*, 83(2), 393–411.
- Tsou, K., Martin, W. J., & Bereiter, D. A. (1996). Suppression stimulus-evoked expression of Fos protein-like immunoreactivity in rat spinal cord by a selective agonist. *Neuroscience*, 4522(3), 791–798.
- Tsou, K., Nogueron, M. I., Muthian, S., Sanudo-Pena, M. C., Hillard, C. J., Deutsch, D. G., et al. (1998b). Fatty acid amide hydrolase is located preferentially in large neurons in the rat central nervous system as revealed by immunohistochemistry. *Neuroscience Letters*, 254(3), 137–140.
- Ueda, M., Iwasaki, H., Wang, S., Murata, E., Poon, K. Y. T., Mao, J., et al. (2014). Cannabinoid receptor type 1 antagonist, AM251, attenuates mechanical allodynia and thermal hyperalgesia after burn injury. *Anesthesiology*, 121(6), 1311–1319.
- Valverde, O., Karsak, M., & Zimmer, A. (2005). Analysis of the endocannabinoid system by using CB1 cannabinoid receptor knockout mice. *Handbook of Experimental Pharmacology*, 168, 117–145.

- Van Sickle, M. D., Duncan, M., Kingsley, P. J., Mouihate, A., Urbani, P., Mackie, K., et al. (2005). Identification and functional characterization of brainstem cannabinoid CB2 receptors. *Science*, 310(5746), 329–332.
- Vaughan, C. W., Connor, M., Bagley, E. E., & Christie, M. J. (2000). Actions of cannabinoids on membrane properties and synaptic transmission in rat periaqueductal gray neurons in vitro. *Molecular Pharmacology*, 57(2), 288–295.
- Vermersch, P. (2011). Sativex((R)) (tetrahydrocannabinol + cannabidiol), an endo-cannabinoid system modulator: Basic features and main clinical data. *Expert Review of Neurotherapeutics*, 11(Suppl. 4), 15–19.
- Viscomi, M. T., Oddi, S., Latini, L., Pasquariello, N., Florenzano, F., Bernardi, G., et al. (2009). Selective CB2 receptor agonism protects central neurons from remote axotomy-induced apoptosis through the PI3K/Akt pathway. *The Journal of Neuroscience*, 29(14), 4564–4570.
- Walker, J. M., Huang, S. M., Strangman, N. M., Tsou, K., & Sanudo-Pena, M. C. (1999). Pain modulation by release of the endogenous cannabinoid anandamide. *Proceedings of the National Academy of Sciences of the United States of America*, 96(21), 12198–12203.
- Watkins, L. R., Milligan, E. D., & Maier, S. F. (2003). Glial proinflammatory cytokines mediate exaggerated pain states: Implications for clinical pain. *Advances in Experimental Medicine and Biology*, 521, 1–21.
- Welch, S. P. (1994). Blockade of cannabinoid-induced antinociception by naloxone benzoylhydrazone (NalBZH). *Pharmacology, Biochemistry, and Behavior*, 49(4), 929–934.
- Welch, S. P., Huffman, J. W., & Lowe, J. (1998). Differential blockade of the antinociceptive effects of centrally administered cannabinoids by SR141716A. *The Journal of Pharmacology and Experimental Therapeutics*, 287(3), 1301–1308.
- Welch, S. P., Thomas, C., & Patrick, G. S. (1995). Modulation of cannabinoid-induced antinociception after intracerebroventricular versus intrathecal administration to mice: Possible mechanisms for interaction with morphine. *The Journal of Pharmacology and Experimental Therapeutics*, 272(1), 310–321.
- Whiting, P. F., Wolff, R. F., Deshpande, S., Di Nisio, M., Duffy, S., Hernandez, A. V., et al. (2015). Cannabinoids for medical use: A systematic review and meta-analysis. *JAMA*, 313(24), 2456–2473.
- Wiley, J. L., & Martin, B. R. (2002). Cannabinoid pharmacology: Implications for additional cannabinoid receptor subtypes. *Chemistry and Physics of Lipids*, 121(1–2), 57–63.
- Wilkerson, J. L., Gentry, K. R., Dengler, E. C., Wallace, J. A., Kerwin, A. A., Armijo, L. M., et al. (2012). Intrathecal cannabidiol CB2R agonist, AM1710, controls pathological pain and restores basal cytokine levels. *Pain*, 153(5), 1091–1106.
- Wilson, A. W., Clayton, N. M., Medhurst, S. J., Bountra, C., & Chessell, I. P. (2004). The FAAH inhibitor URB597 reverses inflammatory pain through a CB1 receptor mediated mechanism. In *049P University of Newcastle Winter Meeting, December 2004, Copyright 2004. The British Pharmacological Society*.
- Wilson, A. R., Maher, L., & Morgan, M. M. (2008). Repeated cannabinoid injections into the rat periaqueductal gray enhance subsequent morphine antinociception. *Neuropharmacology*, 55(7), 1219–1225.
- Wilson-Poe, A. R., Pocius, E., Herschbach, M., & Morgan, M. M. (2013). The periaqueductal gray contributes to bidirectional enhancement of antinociception between morphine and cannabinoids. *Pharmacology, Biochemistry, and Behavior*, 103(3), 444–449.
- Wotherspoon, G., Fox, A., McIntyre, P., Colley, S., Bevan, S., & Winter, J. (2005). Peripheral nerve injury induces cannabinoid receptor 2 protein expression in rat sensory neurons. *Neuroscience*, 135(1), 235–245.
- Yaksh, T. L. (1981). The antinociceptive effects of intrathecally administered levonantradol and desacetyllevonantradol in the rat. *The Journal of Clinical Pharmacology*, 21(S1), 334S–340S.

- Zhang, H. Y., Gao, M., Liu, Q. R., Bi, G. H., Li, X., Yang, H. J., et al. (2014). Cannabinoid CB2 receptors modulate midbrain dopamine neuronal activity and dopamine-related behavior in mice. *Proceedings of the National Academy of Sciences of the United States of America*, 111(46), E5007–E5015.
- Zhang, J., Hoffert, C., Vu, H. K., Groblewski, T., Ahmad, S., & O'Donnell, D. (2003). Induction of CB2 receptor expression in the rat spinal cord of neuropathic but not inflammatory chronic pain models. *The European Journal of Neuroscience*, 17(12), 2750–2754.
- Zheng, T., Zhang, T., Zhang, R., Wang, Z. L., Han, Z. L., Li, N., et al. (2017). Pharmacological characterization of rat VD-hemopressin(alpha), an alpha-hemoglobin-derived peptide exhibiting cannabinoid agonist-like effects in mice. *Neuropeptides*, 63, 83–90.
- Zimmer, A., Zimmer, A. M., Hohmann, A. G., Herkenham, M., & Bonner, T. I. (1999). Increased mortality, hypoactivity, and hypoalgesia in cannabinoid CB1 receptor knock-out mice. *Proceedings of the National Academy of Sciences of the United States of America*, 96(10), 5780–5785.

FURTHER READING

- Di Marzo, V., Bifulco, M., & De Petrocellis, L. (2004). The endocannabinoid system and its therapeutic exploitation. *Nature Reviews Drug Discovery*, 3(9), 771–784.
- Richardson, J. (2001). Cannabinoids modulate pain by multiple mechanisms of action. *The Journal of Pain*, 1, 2–14.
- Salio, C., Doly, S., Fischer, J., Franzoni, M. F., & Conrath, M. (2002). Neuronal and astrocytic localization of the cannabinoid receptor-1 in the dorsal horn of the rat spinal cord. *Neuroscience Letters*, 329(1), 13–16.