

(R)-(-)-[⁷⁷Br]4-Bromo-2,5-dimethoxyamphetamine Labels a Novel 5-Hydroxytryptamine Binding Site in Brain Membranes

STEPHEN J. PEROUTKA, ANNE HAMIK, MELISSA A. HARRINGTON, ANDREW J. HOFFMAN, CHESTER A. MATHIS, PAMELA A. PIERCE, and SAMUEL S.-H. WANG

Department of Neurology, Stanford University, Stanford, California 94305 (S.J.P., A.H., M.A.H., P.A.P., S.S.-H.W.), Department of Medicinal Chemistry and Pharmacognosy, Purdue University, West Lafayette, Indiana 47907 (A.J.H.), and Donner Laboratory, Lawrence Berkeley Laboratory, University of California, Berkeley, California 94720 (C.A.H.)

Received May 5, 1988; Accepted July 13, 1988

SUMMARY

(R)-(-)-[⁷⁷Br]4-Bromo-2,5-dimethoxyamphetamine [(R)-(-)-[⁷⁷Br]DOB] was synthesized to a high specific activity (1875 ± 50 Ci/mmol) and used to label membrane-associated recognition sites in rat brain. (R)-(-)-[⁷⁷Br]DOB displayed high affinity ($K_D = 0.60 \pm 0.08$ nM) for a relatively low density of binding sites ($B_{max} = 1.2 \pm 0.08$ pmol/g of tissue) in rat cortical membranes as compared with [³H]ketanserin ($K_D = 0.65 \pm 0.1$ nM; $B_{max} = 6.2 \pm 0.6$ pmol/g of tissue). Guanine, but not adenine, nucleotides were found to inhibit specific (R)-(-)-[⁷⁷Br]DOB binding. GTP (10^{-4} M) did not eliminate specific (R)-(-)-[⁷⁷Br]DOB binding but caused a competitive inhibition of the radioligand. Drug competition studies of 5-hydroxytryptamine (5-HT) and related agents indicate that both putative agonists and antagonists display

nanomolar potency for these sites. A significant correlation ($p < 0.01$) exists between drug potencies for (R)-(-)-[⁷⁷Br]DOB-labeled sites and both 5-HT₂ ($r = 0.64$) and 5-HT_{1c} ($r = 0.68$) binding sites. However, the sites do not appear to be identical. Moreover, a significant correlation exists between drug potencies for (R)-(-)-[⁷⁷Br]DOB-labeled sites and human hallucinogenic drug potencies ($r = 0.89$; $p < 0.01$). We conclude that (R)-(-)-[⁷⁷Br]DOB labels a unique 5-HT recognition site in rat brain that does not coincide with previously described 5-HT binding site subtypes. The (R)-(-)-[⁷⁷Br]DOB site does not appear to be a high affinity "state" of the 5-HT₂ receptor but may label a subset of heterogeneous 5-HT₂ recognition sites.

(R)-(-)DOB is a potent human hallucinogen that selectively interacts with 5-HT receptors (1). Recently, (±)-[³H]DOB was synthesized and reported to label a high affinity "state" of the 5-HT₂ receptor in the central nervous system (1-3). This conclusion was based on the fact that the affinity of a variety of 5-HT₂ antagonists for the (±)-[³H]DOB site correlated with the affinity for the 5-HT₂ receptor labeled by [³H]ketanserin. However, the site displayed nanomolar affinity for 5-HT, in contrast to the micromolar affinity displayed by 5-HT and 5-HT₂ receptors, as defined in both radioligand studies using ³H-antagonists (4, 5) and functional studies (6). In addition, the density of (±)-[³H]DOB sites represented approximately 5% of the density of 5-HT₂ sites labeled by [³H]ketanserin in rat frontal cortex brain membranes. Unfortunately, the low specific activity of commercially available (±)-[³H]DOB (16 Ci/mmol) has hampered the extensive characterization of this potentially important binding site.

This work was supported in part by the John A. and George L. Hartford Foundation, Inc., the Alfred P. Sloan Foundation, and National Institutes of Health Grants NS 12151-13, NS 23560-02 (S.J.P.), MH 17047-06 (M.A.H.), and NS 22899-02 (C.A.H.).

In order to further investigate this site, we have developed a novel brominated radioligand of extremely high specific activity, (R)-(-)-[⁷⁷Br]DOB. Our preliminary analysis of this non-racemic compound revealed that it labeled a relatively low density of binding sites that did not appear to coincide with previously defined 5-HT binding site subtypes (7). We now report on the detailed characterization of this novel 5-HT binding site.

Materials and Methods

Radioligand binding assays were performed according to the methods of Titeler and colleagues (1-3) and Wang *et al.* (7). In brief, frozen rat brains were thawed from -70° and brain regions were dissected. Tissues were homogenized in 20 volumes of 50 mM Tris·HCl (pH 7.7 at 25°) and centrifuged at 45,000 × g for 10 min. The pellet was resuspended in Tris·HCl buffer and incubated for 10 min at 37°. After a second centrifugation, the pellet was resuspended in 80 volumes of a buffer consisting of 50 mM Tris·HCl, 0.5 mM EDTA, 10 mM MgCl₂, 0.1% ascorbate, and 10⁻⁵ M pargyline. The homogenates were immediately used in the binding assay.

Assays consisted of 100 μl of a solution of (R)-(-)-[⁷⁷Br]DOB (final

ABBREVIATIONS: DOB, 1-(4-bromo-2,5-dimethoxy 4-bromoamphetamine; 5-HT, 5-hydroxytryptamine; d-LSD; d-lysergic acid diethylamide; GppNhp, guanosine 5'-β,γ-imidotriphosphate; GTPγS, guanosine 5'-O-(3-thiotriphosphate; 8-OH-DPAT, 8-hydroxy-2-di-N-propylamino-tetralin; DOI, (±)-(2,5-dimethoxy-4-iodoamphetamine; 5-CT, 5-carboxyamidotryptamine.

concentrations, 0.052–9.6 nM), 800 μ l of tissue suspension, and 100 μ l of buffer or displacing drug. Drugs were dissolved in buffer, except for (+)-butaclamol, which was dissolved to 10^{-3} M in ethanol and then diluted in buffer. After incubation at 25° for 30 min, the assay mixtures were rapidly filtered through no. 32 glass fiber filters (Schleicher and Schuell; Keene, NH) and washed two times with 5 ml of 50 mM Tris-HCl buffer. Specific binding was stable after incubation periods of approximately 15–60 min. Radioactivity was measured by liquid scintillation counting in 5 ml of Aquasol (New England Nuclear; Boston MA) at an estimated 80% efficiency. Specific binding was defined as the excess over blanks taken in the presence of 10^{-8} M DOI, a concentration of nonradioactive drug that displaces greater than 99% of specific radioligand binding. Competition experiments using (R)-(-)-[⁷⁷Br]DOB were performed using 0.08–2.7 nM of radioactive ligand. At these concentrations, 80–90% of the total binding was specific.

⁷⁷Br was synthesized by a high energy spallation reaction at Los Alamos National Laboratories (Los Alamos, NM) (8). (R)-(-)-[⁷⁷Br]DOB was synthesized to a radiochemical purity of greater than 99% by a modified Coenen synthesis (9, 10). The initial specific activity was determined by high pressure liquid chromatography ultraviolet quantitation. The specific activity was 375 Ci/mmol in the first synthesis and 1875 Ci/mmol in the second synthesis. The specific activity was corrected to allow for the decay of ⁷⁷Br ($t_{1/2} = 57$ hr) for each experiment.

Drugs were obtained from commercial sources except for (+)-DOB, (-)-DOB, *d*-LSD, and (-)-dimethoxymethyl amphetamine, which were obtained from the National Institute on Drug Abuse (Bethesda, MD). 5-CT was the generous gift of Dr. Roger Whiting of Syntex Corporation (Palo Alto, CA).

Results

Saturation analysis of specific (R)-(-)-[⁷⁷Br]DOB binding in rat brain membranes. Specific (R)-(-)-[⁷⁷Br]DOB binding to rat brain membranes was saturable and of high affinity. The experiment shown in Fig. 1A was performed using (R)-(-)-[⁷⁷Br]DOB with a specific activity of 450 Ci/mmol. At a concentration of 0.12 nM, specific binding accounted for 88% of total binding. Half-maximal specific binding was achieved at approximately 0.80 nM (R)-(-)-[⁷⁷Br]DOB, at which specific binding accounted for 84% of total binding. At the highest concentration of (R)-(-)-[⁷⁷Br]DOB analyzed in this single experiment (9.6 nM), specific binding represented 50% of total binding. The Scatchard analysis of these data (Fig. 1B) is consistent with a homogeneous population of binding sites.

A total of eight saturation studies was performed in the present study. Scatchard analysis of the saturation data reveals a K_D value of 0.60 ± 0.08 nM with a B_{max} value of 1.2 ± 0.2 pmol/g in rat cortical membranes. By contrast, saturation studies using [³H]ketanserin in the same tissue (six experiments) reveal a K_D value of 0.65 ± 0.1 nM and a B_{max} value of 6.2 ± 0.6 pmol/g of tissue. Thus, the density of (R)-(-)-[⁷⁷Br]DOB-labeled sites represents only 19% of the density of 5-HT₂ sites labeled by [³H]ketanserin.

Regional studies were also performed using approximately 0.3 nM (R)-(-)-[⁷⁷Br]DOB. As shown in Table 1, the cortex displays the highest amount of specific (R)-(-)-[⁷⁷Br]DOB binding (100% = 0.52 ± 0.4 pmol/g) whereas the caudate has slightly less specific binding. The midbrain and hippocampus display intermediate amounts of specific (R)-(-)-[⁷⁷Br]DOB binding whereas the cerebellum has the least amount of specific binding. These findings are similar to the regional differences in (±)-[³H]DOB binding that have been observed previously (3).

Nucleotide effects on specific (R)-(-)-[⁷⁷Br]DOB bind-

ing. The results of studies examining 10^{-4} M nucleotide effects on (R)-(-)-[⁷⁷Br]DOB binding are given in Table 2. GTP and its nonhydrolyzable analogues GppNHp and GTP γ S have the greatest effect on the specific binding of (R)-(-)-[⁷⁷Br]DOB. These nucleotides (10^{-4} M) significantly inhibit specific (R)-(-)-[⁷⁷Br]DOB binding by 63–75%. The inhibitory effect of GDP is slightly less pronounced (49%) but is still significantly different from control values ($p < 0.01$). By contrast, GMP and the adenine nucleotides are markedly less effective in modulating specific (R)-(-)-[⁷⁷Br]DOB binding. These nucleotides have no significant effect on the specific binding of (R)-(-)-[⁷⁷Br]DOB at a concentration of 10^{-4} M.

This decrease in binding appears to be due to an affinity shift with no change in site density, as shown by Scatchard analysis. For example, the effect of GTP on the affinity and density of (R)-(-)-[⁷⁷Br]DOB-labeled binding sites was determined in rat cortex. Under control conditions in this series of experiments (three experiments), the K_D value of (R)-(-)-[⁷⁷Br]DOB was 0.60 ± 0.08 nM with a B_{max} value of 1.4 ± 0.1 pmol/g of tissue. In the presence of 10^{-5} M GTP, the K_D is slightly increased to 0.92 ± 0.1 nM whereas the B_{max} is essentially unchanged at 1.1 ± 0.1 pmol/g of tissue. The ability of GTP to alter the K_D of (R)-(-)-[⁷⁷Br]DOB is best observed in the presence of 10^{-4} M GTP. Under this condition, the K_D for (R)-(-)-[⁷⁷Br]DOB is 1.6 ± 0.4 nM with a B_{max} value of 1.2 ± 0.2 pmol/g of tissue. A representative experiment is shown in Fig. 2.

Drug competition studies with (R)-(-)-[⁷⁷Br]DOB-labeled binding sites in rat cortex. The results of drug competition studies using (R)-(-)-[⁷⁷Br]DOB are listed in Table 3. (-)-DOB is the most potent displacing agent ($K_i = 0.62 \pm 0.1$ nM) and is an order of magnitude more potent than (+)-DOB ($K_i = 18 \pm 9$ nM). Specific (R)-(-)-[⁷⁷Br]DOB binding is also displaced by nanomolar concentrations of 5-HT, metergoline, and a number of putative 5-HT₂-selective antagonists. By contrast, the 5-HT_{1A}-selective agent 8-OH-DPAT, the 5-HT_{1B} agent RU 24969, the 5-HT_{1D} agent 5-CT, and the 5-HT₃-selective agent ICS 205-930 were significantly less potent at (R)-(-)-[⁷⁷Br]DOB-labeled sites.

Representative drug competition curves are shown in Fig. 3. 5-HT, *d*-LSD, and ketanserin are inactive in competing for specific (R)-(-)-[⁷⁷Br]DOB binding at concentrations below 10^{-9} M. Monophasic competition is apparent at drug concentrations between 10^{-9} and 10^{-7} M. In the experiment shown in Fig. 3, 5-HT is slightly more potent than *d*-LSD and ketanserin. The slopes of the three competition curves are nearly identical and represent Hill slopes of approximately unity.

Correlation of drug affinities for (R)-(-)-[⁷⁷Br]DOB-labeled sites and drug affinities for 5-HT₂ and 5-HT_{1C} binding sites. Drug affinities for (R)-(-)-[⁷⁷Br]DOB-labeled sites were compared with drug affinities for 5-HT₂ and 5-HT_{1C} binding sites as defined in previous studies. As shown in Fig. 4A, a significant correlation exists between drug K_i values for (R)-(-)-[⁷⁷Br]DOB-labeled sites in rat cortex and 5-HT₂ binding sites ($r = 0.64$; $p < 0.01$; slope = 0.98; $n = 19$). Despite the statistical significance of this correlation, a number of apparent discrepancies should be noted. For example, the affinity of 5-HT for (R)-(-)-[⁷⁷Br]DOB-labeled sites (3.5 nM) is 2 to 3 orders of magnitude more potent than its affinity (562–30,000 nM) for [³H]ketanserin-labeled 5-HT₂ sites in rat cortex as reported in multiple laboratories (11, 12). Conversely, the affinity of chlor-




Fig. 1. Saturation analysis of (R)-(-)-[⁷⁷Br]DOB binding to rat cortical membranes. Binding assays were carried out as described in Materials and Methods. A, Data from a typical saturation experiment. Nonspecific binding was determined in the presence of 1 μ M ($\pm$)-DOI. B, Scatchard analysis of the data shown in A.

TABLE 1
Regional variations in specific (R)-(-)-[⁷⁷Br]DOB binding

Various rat brain regions were analyzed for specific (R)-(-)-[⁷⁷Br]DOB binding using 0.25 $\pm$ 0.1 nM radioligand as described in Materials and Methods. All experiments were performed in triplicate and repeated two to six times. The amount of specific (R)-(-)-[⁷⁷Br]DOB binding in cortex (0.52 $\pm$ 0.4 pmol/g) was defined as 100%.

Region	Specific (R)-(-)-[⁷⁷ Br]DOB binding % cortical values
Cortex	100 $\pm$ 6
Caudate	89 $\pm$ 10
Midbrain	50 $\pm$ 2
Hippocampus	42 $\pm$ 0
Cerebellum	17 $\pm$ 4

promazine for (R)-(-)-[⁷⁷Br]DOB-labeled sites (70 nM) is more than an order of magnitude less potent than its affinity for [³H]ketanserin-labeled 5-HT₂ sites (2.8 nM) in rat cortex (3).

A slightly higher correlation coefficient ($r = 0.68$; $p < 0.01$; slope = 0.80; $n = 14$) is observed between drug affinities for (R)-(-)-[⁷⁷Br]DOB-labeled sites in rat cortex and 5-HT_{1C} binding sites (Fig. 4B). Again, however, the two sites are not identical, inasmuch as drugs such as spiperone (1200 nM) (11) and ketanserin (98 nM) (13) are significantly less potent at 5-HT_{1C} sites than at (R)-(-)-[⁷⁷Br]DOB-labeled sites. Metergoline (0.51 nM) (11), on the other hand, is more potent at 5-

TABLE 2
Nucleotide effects on (R)-(-)-[⁷⁷Br]DOB binding in rat cortical membranes

Radioligand binding assays were performed as described in Materials and Methods. Values given are the means $\pm$ standard errors of two to six experiments, each performed in triplicate.

Nucleotide	Specific binding of (R)-(-)-[⁷⁷ Br] DOB
10^{-4} M	% of control
GppNHp	25 $\pm$ 3 ^a
GTP γ S	37 $\pm$ 8 ^b
GTP	35 $\pm$ 6 ^c
GDP	51 $\pm$ 3 ^c
ADP	83 $\pm$ 9
ATP	84 $\pm$ 9
AMP	92 $\pm$ 0
GMP	100 $\pm$ 20

^a $p < 0.02$.

^b $p < 0.05$.

^c $p < 0.01$.

HT_{1C} than at (R)-(-)-[⁷⁷Br]DOB binding sites. In addition, DOB has been reported to have an affinity of 69 nM for 5-HT_{1C} binding sites (14). Moreover, 5-HT_{1C} sites are found in extremely high density in the choroid plexus, whereas we failed




Fig. 2. GTP effects of (R)-(-)[⁷⁷Br]DOB saturation studies in rat cortical membranes. Radioligand binding assays were performed as described in Materials and Methods. Nonspecific binding was determined in the presence of 1 μ M ($\pm$)-DOI. The Scatchard data are taken from a single experiment, performed in triplicate. The experiment was repeated three times, with similar results.

TABLE 3

Drug affinities for (R)-(-)[⁷⁷Br]DOB-labeled sites in rat cortex

Radioligand binding assays were performed as described in Materials and Methods. Values given are the means $\pm$ standard errors of three to six experiments, each performed in triplicate.

	K_i
	nM
1. (-)-DOB	0.62 $\pm$ 0.1
2. ($\pm$)-DOI	1.3 $\pm$ 0.3
3. Metergoline	2.7 $\pm$ 0.5
4. (-)-dimethoxyethyl amphetamine	3.1 $\pm$ 0.6
5. 5-HT	3.5 $\pm$ 0.8
6. Spiperone	3.8 $\pm$ 2
7. Mesulergine	4.5 $\pm$ 0.9
8. Ketanserin	4.8 $\pm$ 0.7
9. α -LSD	4.8 $\pm$ 0.6
10. (-)-dimethoxymethyl amphetamine	6.4 $\pm$ 1
11. Methiothepin	6.5 $\pm$ 3
12. Cinanserin	7.2 $\pm$ 0.6
13. (+)-DOB	18 $\pm$ 9
14. Psilocin	19 $\pm$ 6
15. <i>m</i> -chlorophenylpiperazine	22 $\pm$ 3
16. Psilocybin	25 $\pm$ 4
17. (+)-Butaclamol	29 $\pm$ 10
18. Quipazine	31 $\pm$ 1
19. Tryptamine	49 $\pm$ 10
20. Chlorpromazine	70 $\pm$ 40
21. RU 24969	91 $\pm$ 3
22. 5-CT	320 $\pm$ 80
23. Mescaline	440 $\pm$ 100
24. ($\pm$)-3,4,5-trimethoxyamphetamine	450 $\pm$ 100
25. ($\pm$)-2,5-dimethoxyamphetamine	580 $\pm$ 100
26. 8-OH-DPAT	2,700 $\pm$ 300
27. (-)-Butaclamol	10,000 $\pm$ 2,000
28. ICS 205-930	17,000 $\pm$ 6,000

to identify significant densities of (R)-(-)[⁷⁷Br]DOB binding in the rat choroid plexus in preliminary autoradiographic studies.¹

¹ Unpublished observations.




Fig. 3. Drug competition studies versus specific (R)-(-)[⁷⁷Br]DOB binding in rat cortical membranes. Radioligand binding assays were performed as described in Materials and Methods. Nonspecific binding was determined in the presence of 1 μ M ($\pm$)-DOI. Data shown are the results of a single experiment, performed in triplicate. Each experiment was repeated 3–6 times. Drugs analyzed are 5-HT (●), α -LSD (▲), and ketanserin (○).




Fig. 4. Correlation of drug potencies for (R)-(-)[⁷⁷Br]DOB-labeled sites and drug potencies for 5-HT₂ and 5-HT_{1C} binding sites. Drug affinities (K_i , nM) for (R)-(-)[⁷⁷Br]DOB-labeled sites were obtained from Table 3. A, Drug affinities for 5-HT₂ binding sites were obtained from Lyon *et al.* (3), Hoyer *et al.* (11), and Glennon *et al.* (20). B, Drug affinities for 5-HT_{1C} binding sites were obtained from Lyon *et al.* (3), Hoyer *et al.* (11), and Pazos *et al.* (13).

Correlation of drug affinities for (*R*)-(-)-[⁷⁷Br]DOB-labeled sites and human hallucinogenic dosages. As shown in Table 3, a number of hallucinogenic agents display high affinity for (*R*)-(-)-[⁷⁷Br]DOB-labeled binding sites. A significant correlation exists ($r = 0.89$; $p < 0.01$; slope = 0.93; $n = 11$) between drug affinities for these sites in rat cortex and the average dosage that has been reported to induce hallucinations in humans (15–17). As illustrated in Fig. 5, 10 of the 11 hallucinogenic agents lie near the linear regression line. The single exception is *d*-LSD; the drug is significantly less potent at (*R*)-(-)-[⁷⁷Br]DOB binding sites than might have been predicted on the basis of the affinities displayed by the other hallucinogenic agents.

Discussion

The major finding of the present study is that (*R*)-(-)-[⁷⁷Br]DOB specifically labels a distinct and unique population of 5-HT recognition sites in rat brain. The sites labeled by (*R*)-(-)-[⁷⁷Br]DOB do not appear to correspond to previously defined 5-HT₁, 5-HT₂, or 5-HT₃ binding subtypes because, for example, they display nanomolar affinity for both 5-HT and spiperone (17). 8-OH-DPAT, RU 24969, 5-CT, and ICS 205-930 (which display high affinity for 5-HT_{1A}, 5-HT_{1B}, 5-HT_{1D}, and 5-HT₃ sites, respectively) all display relatively low affinities for the sites labeled by (*R*)-(-)-[⁷⁷Br]DOB. Although a significant correlation is observed between drug affinities for (*R*)-(-)-[⁷⁷Br]DOB binding sites and drug affinities for both 5-HT₂ and 5-HT_{1C} binding sites, the correlation appears to be based more on the similarities between these sites than on their exact correspondence.

The pharmacological characteristics of the (*R*)-(-)-[⁷⁷Br]DOB binding sites appear to be identical to those of the sites labeled by (±)-[³H]DOB (1–3). In contrast to our interpretation of the data, Lyon *et al.* (3) concluded that (±)-[³H]DOB labels a “high affinity agonist state” of the 5-HT₂ receptor. A number of facts were cited by these authors to support this hypothesis. For example, presumed agonists, but not antagonists, discriminate two subpopulations of 5-HT₂ sites labeled by [³H]ketanserin based on the analysis of drug competition curves (18). However, the ability to discriminate two subpopulations of [³H]

ketanserin-labeled 5-HT₂ recognition sites is not agonist specific. For example, trifluoromethylpiperazine displays a 10-fold selectivity for the “high” versus “low” affinity component of [³H]ketanserin binding, yet trifluoromethylpiperazine is actually a pure antagonist of the 5-HT₂ receptor when analyzed using biochemical models of 5-HT₂ receptor function such as phosphoinositide turnover (19).

A second major observation that Titeler *et al.* (2) used to support their hypothesis that [³H]DOB labeled a “high affinity agonist state” of the 5-HT₂ receptor was the effect of 10⁻⁴ M GTP on [³H]ketanserin-labeled binding sites. Although 10⁻⁴ M GTP had no significant effect on [³H]ketanserin binding, it did cause a “shift” of putative agonist potencies. In addition, an increase was observed in the Hill slope of the putative agonist competition curves versus [³H]ketanserin. They suggested that 10⁻⁴ M GTP effectively eliminated the “high affinity agonist state” of the 5-HT₂ receptor (18) with a shift of these sites into a “low affinity agonist state.” By contrast, the present data demonstrate that 10⁻⁴ M GTP does not eliminate (*R*)-(-)-[⁷⁷Br]DOB binding in rat cortex. This concentration of nucleotide appears to induce an approximately 3- to 5-fold decrease in the affinity of (*R*)-(-)-[⁷⁷Br]DOB without changing the density of the binding sites (Fig. 2).

A third fact cited by Titeler and colleagues (1–3) to support their hypothesis was that antagonist potencies at 5-HT₂ sites correlate significantly with their affinity for (*R*)-(-)-[⁷⁷Br]DOB-labeled sites. They did not include drugs that they considered to be “agonists” at the 5-HT₂ receptor. Because the agonist versus antagonist properties of drugs are difficult, and perhaps impossible, to determine from radioligand studies, all drugs were included in the correlations reported in the present report. As noted above, the *K_i* values for the drugs analyzed in the present study were found to significantly “correlate” with their affinity for both the 5-HT₂ and 5-HT_{1C} binding sites. In contrast to the conclusion of Titeler and colleagues, we suggest that this correlation implies a similarity, rather than an identity, between the binding sites.

Therefore, we would like to offer an alternative hypothesis to the suggestion of Titeler and colleagues (i.e. that DOB labels a “high affinity agonist state” of the 5-HT₂ receptor). Our conclusion is based on an interpretation of both our own data using (*R*)-(-)-[⁷⁷Br]DOB and the extensive and important work of Titeler and colleagues using (±)-[³H]DOB. We believe that the available data concerning the unique pharmacological properties of these sites indicate that, at the present time, they should be considered a novel and distinct subset of “5-HT binding sites” (17). Furthermore, the pharmacological similarity of these sites to both the 5-HT_{1C} and 5-HT₂ receptors suggests that the sites labeled by (*R*)-(-)-[⁷⁷Br]DOB may be structurally similar to these other 5-HT receptor subtypes. Additional radioligand, functional, and molecular biological studies are needed before the sites labeled by (*R*)-(-)-[⁷⁷Br]DOB can be more definitively categorized. Given the controversies surrounding the nomenclature of 5-HT recognition sites, it may be best to refer to this site as the “DOB binding site” until such studies are completed.

Another potentially important observation in the present study is the significant correlation that exists between drug affinities for the (*R*)-(-)-[⁷⁷Br]DOB binding site and human hallucinogenic doses. This correlation has also been reported recently by Titeler *et al.* (12). In addition, Glennon and Rose-




Fig. 5. Correlation of drug potencies for (*R*)-(-)-[⁷⁷Br]DOB-labeled sites and human hallucinogenic doses. Drug affinities (*K_d*, nM) for (*R*)-(-)-[⁷⁷Br]DOB-labeled sites were obtained from Table 3. Human hallucinogenic drug dosages were obtained from Shulgin (14), Jaffe (15) and Glennon and Rosecrans (16).

crans (16) have previously reported a significant correlation between human hallucinogenic doses and drug effects on the 5-HT receptor in rat fundus (a 5-HT receptor that does not coincide with the "DOB binding site" described in the present report nor with the 5-HT₁, 5-HT₂, or 5-HT₃ classification system). More recently, Glennon *et al.* (20) reported a significant correlation between human hallucinogenic doses and 5-HT₂ receptor affinity. Therefore, the (*R*)-(-)-[⁷⁷Br]DOB binding site represents the third putative 5-HT receptor for which a significant correlation with hallucinogenic doses has been documented. Although the mechanism of action of hallucinogenic drugs in humans remains unknown, these data suggest that each of the three sites may play a role in the pathophysiology of hallucinosis.

Finally, a central feature of this study was the use of a ⁷⁷Br-labeled ligand. High specific activity ⁷⁷Br should facilitate the analysis of radioligand binding in low density regions and/or tissues. In addition, short-lived positron-emitting isotopes of bromine may also be useful in positron emission tomographic scanning techniques. ⁷⁷Br itself is not appropriate for such studies due to its low positron emission abundance, but the isotope ⁷⁶Br may be useful as an *in vivo* radiolabel in human positron emission tomograph studies of hallucinogens. The use of such radiolabeled hallucinogens holds great promise for further study of the mechanism of hallucinogenic drug actions in man.

Acknowledgments

We thank Cynthia M. Rosewicz for her assistance in the preparation of the manuscript.

References

1. Titeler, M., R. A. Lyon, K. H. Davis, and R. A. Glennon. Role of [³H]-4-bromo-2,5-dimethoxyphenylisopropylamine ([³H]DOB), a 5-HT₂ agonist radioligand. *Biochem. Pharmacol.* **36**:3265-3271 (1985).
2. Titeler, M., K. Herrick, R. A. Lyon, J. D. McKenney, and R. A. Glennon. [³H]DOB: a specific agonist radioligand for 5-HT₂ serotonin receptors. *Eur. J. Pharmacol.* **117**:145-146 (1987).
3. Lyon, R. A., K. H. Davis, and M. Titeler. ³H-DOB (4-bromo-2,5-dimethoxyphenylisopropylamine) labels a guanyl nucleotide-sensitive state of cortical 5-HT₂ receptors. *Mol. Pharmacol.* **31**:194-199 (1987).
4. Leysen, J. E., C. J. E. Niemegeers, J. P. Tollenaere, and P. M. Laduron. Serotonergic component of neuroleptic receptors. *Nature (Lond.)* **272**:166-171 (1978).
5. Peroutka, S. J., and S. H. Snyder. Multiple serotonin receptors: differential binding of [³H]5-hydroxytryptamine, [³H]lysergic acid diethylamide, and [³H]spiroperidol. *Mol. Pharmacol.* **16**:687-699 (1979).
6. Leysen, J. E., D. De Chaffoy De Courcelles, F. De Clerck, C. J. E. Niemegeers, and J. M. Van Nueten. Serotonin-S₂ receptor binding sites and functional correlates. *Neuropharmacology* **23**:1493-1501 (1984).
7. Wang, S. H., C. A. Mathis, and S. J. Peroutka. (*R*)-(-)-2,5-Dimethoxy-4-⁷⁷bromoamphetamine [⁷⁷Br-*R*(-)-DOB]: a novel radioligand which labels a 5-HT binding site subtype. *Psychopharmacology* **94**:431-432 (1988).
8. Grant, P. M., R. E. Whipple, J. W. Barnes, G. E. Bentley, P. M. Wanek, and H. A. O'Brien, Jr. The production and recovery of bromine-77 at Los Alamos for nuclear medicine studies. *J. Inorg. Nucl. Chem.* **43**:2217-2222 (1981).
9. Coenen, H. H. Synthesis of ⁷⁷Br-labeled 2-(4-bromo-2,5-dimethoxyphenyl)-isopropylamine with high specific activity. *J. Labelled Compd. Radiopharm.* **18**:739-746 (1981).
10. Mathis, C. A., A. J. Hoffman, D. E. Nichols, and A. T. Shulgin. Synthesis of high specific activity ¹²⁵I- and ¹²³I-labelled enantiomers of 2,5-dimethoxy-4-iodophenylisopropylamine (DOI). *J. Labelled Compd. Radiopharm.*, in press.
11. Hoyer, D., G. Engel, and H. O. Kalkman. Molecular pharmacology of 5-HT₁ and 5-HT₂ recognition sites in rat and pig brain membranes: radioligand binding studies with [³H]5-HT, [³H]8-OH-DPAT, (-)-[¹²⁵I]iodocyanopindolol, [³H]mesulergine and [³H]ketanserin. *Eur. J. Pharmacol.* **118**:13-23 (1985).
12. Titeler, M., R. A. Lyon, and R. A. Glennon. Radioligand binding evidence implicates the brain 5-HT₂ receptor as a site of action for LSD and phenylisopropylamine hallucinogens. *Psychopharmacology* **94**:213-216 (1988).
13. Pazos, A., D. Hoyer, and J. M. Palacios. The binding of serotonergic ligands to the porcine choroid plexus: characterization of a new type of serotonin recognition site. *Eur. J. Pharmacol.* **106**:539-546 (1984).
14. Shulgin, A. T. Psychotomimetic drugs: structure-activity relationships, in *Handbook of Psychopharmacology* (S. H. Snyder, L. L. Iversen, and S. D. Iversen, eds.), Vol. 2. Plenum, New York, 243-333 (1978).
15. Jaffe, J. H. Drug addiction and drug abuse, in *The Pharmacological Basis of Therapeutics* (A. G. Gilman, L. S. Goodman, and A. Gilman, eds.), MacMillan, New York, 535-584 (1980).
16. Glennon, R. A., and J. A. Rosecrans. Indolealkylamine and phenalkylamine hallucinogens: a brief overview. *Neurosci. Biobehav. Rev.* **6**:489-497 (1982).
17. Peroutka, S. J. 5-Hydroxytryptamine receptor subtypes. *Annu. Rev. Neurosci.* **11**:45-60 (1988).
18. Battaglia, G., M. Shannon, and M. Titeler. Guanyl nucleotide and divalent cation regulation of cortical S₂ serotonin receptors. *J. Neurochem.* **43**:1213-1219 (1984).
19. Conn, P. J., and E. Sanders-Bush. Relative efficacies of piperazines at the phosphoinositide hydrolysis-linked serotonergic 5-HT₂ and 5-HT_{1C} receptors. *J. Pharmacol. Exp. Ther.* **242**:552-557 (1987).
20. Glennon, R. A., M. Titeler, and J. D. McKenney. Evidence for 5-HT₂ involvement in the mechanism of action of hallucinogenic agents. *Life Sci.* **35**:2505-2511 (1984).

Send reprint requests to: Stephen J. Peroutka, M.D., Ph.D., Department of Neurology, C-338, Stanford University Medical Center, Stanford, CA 94305.
