

Psychedelics and hypnosis: Commonalities and therapeutic implications

Clément E Lemerrier¹  and Devin B Terhune²

Abstract

Background: Recent research on psychedelics and hypnosis demonstrates the value of both methods in the treatment of a range of psychopathologies with overlapping applications and neurophenomenological features. The potential of harnessing the power of suggestion to influence the phenomenological response to psychedelics toward more therapeutic action has remained unexplored in recent research and thereby warrants empirical attention.

Aims: Here we aim to elucidate the phenomenological and neurophysiological similarities and dissimilarities between psychedelic states and hypnosis in order to revisit how contemporary knowledge may inform their conjunct usage in psychotherapy.

Methods: We review recent advances in phenomenological and neurophysiological research on psychedelics and hypnosis, and we summarize early investigations on the coupling of psychedelics and hypnosis in scientific and therapeutic contexts.

Results/outcomes: We highlight commonalities and differences between psychedelics and hypnosis that point to the potential efficacy of combining the two in psychotherapy. We propose multiple research paths for coupling these two phenomena at different stages in the preparation, acute phase and follow-up of psychedelic-assisted psychotherapy in order to prepare, guide and integrate the psychedelic experience with the aim of enhancing therapeutic outcomes.

Conclusions/interpretation: Harnessing the power of suggestion to modulate response to psychedelics could enhance their therapeutic efficacy by helping to increase the likelihood of positive responses, including mystical-type experiences.

Keywords

Consciousness, hypnosis, psychedelics, psychotherapy, suggestion

Introduction

Alterations of consciousness have aided therapeutic processes in a variety of practices from ancient shamanism to approaches used in contemporary psychotherapy (Buckley and Galanter, 1979). Psychedelic-induced states of consciousness have been used as a central feature of many religious, sacramental and medicinal practices in a variety of cultures (Schultes and Hofmann, 1979). Psychedelics induce profound changes in perception, thought, affect and self-awareness (Preller and Vollenweider, 2018) and are recognized as agents capable of eliciting deeply personal and spiritually meaningful experiences (so-called mystical-type experiences) associated with sustained beneficial effects on personality and on psychological well-being (Griffiths et al., 2008, 2011; MacLean et al., 2011; Schmid and Liechti, 2017). In the last decade, we have witnessed a surge in research exploring the impact of psychedelics, such as lysergic acid diethylamide (LSD) and psilocybin, on psychological and brain functions (Carhart-Harris et al., 2012, 2016b; De Araujo et al., 2012; Komater et al., 2015; Kraehenmann et al., 2015; Tagliazucchi et al., 2016) and in psychotherapy (Bogenschutz et al., 2015; Carhart-Harris et al., 2016a; Garcia-Romeu et al., 2015; Griffiths et al., 2016; Johnson et al., 2014; Ross et al., 2016; for reviews, see Carhart-Harris and Goodwin, 2017; Liechti, 2017). It is increasingly being recognized that research on psychedelics has the potential to inform our understanding of the neurobiology of psychiatric disorders and consciousness in general and to provide a platform upon which the therapeutic benefits of alterations in consciousness can be safely harnessed and studied.

Hypnosis began as a therapeutic discipline in the eighteenth century and is considered as the first western conception of psychotherapy: the first time a spoken interaction between a doctor and a patient was thought to have therapeutic potential (Ellenberger, 1970). Hypnosis is a mind–body intervention that consists of a set of procedures comprising the administration of verbal suggestions for alterations in affect, cognition and perception (Elkins, 2017). The therapeutic potential of hypnosis has been documented since its origins, but it is only recently that the method has experienced a renewed interest for clinicians with accumulating evidence for its efficacy for the treatment of multiple symptoms and conditions (Elkins et al., 2013; Hasan et al., 2014; Patterson and Jensen, 2003; Schaefer et al., 2014; for reviews, see Elkins, 2017; Terhune et al., 2017). Recent cognitive and neurophysiological studies have begun to elucidate the underlying bases of hypnosis and have shown that manipulation of subjective awareness with hypnotic suggestion can provide

¹Association AlternatiMed, Bordeaux, France

²Department of Psychology, Goldsmiths, University of London, UK

Corresponding author:

Clément E Lemerrier, Association AlternatiMed, 26 rue Leydet, Bordeaux 33800, France.

Email: clement.lemerrier@alternatimed.com

insights into brain mechanisms involved in pain, perception, attention, motor control, sense of agency and executive control (for reviews, see Oakley and Halligan, 2013; Terhune et al., 2017). Moreover, mounting evidence points to the promise of using hypnotic suggestion to model and experimentally modulate conscious states that are otherwise challenging to control, such as pathological symptoms (Oakley and Halligan, 2013; Woody and Szechtman, 2011) or anomalous experiences (Hastings, 2006; Lynn and Evans, 2017; Zeev-Wolf et al., 2017).

Despite multiple phenomenological parallels, recent research on psychedelics and hypnosis has occurred largely in isolation. By contrast, earlier research highlighted overlapping characteristics between these phenomena (Barber, 1970) and the potential for joint use in psychotherapy (Levine and Ludwig, 1966; Ludwig and Levine, 1965). In particular, the potential of harnessing the power of suggestion to influence the response to psychedelics may have implications for both clinical and basic research. In this article, we review recent advances that suggest commonalities between these two phenomena. Against this backdrop, we revisit early investigations on the coupling of psychedelics and hypnosis in scientific and therapeutic contexts and conclude by describing potential applications of hypnosis in psychedelic-assisted psychotherapy.

Psychedelics

Classic psychedelics, such as LSD, mescaline, psilocybin, and N,N-dimethyltryptamine (DMT), exert their hallucinatory effects primarily through activation of serotonin 2A receptors (5-HT_{2A}) (González-Maeso et al., 2007; for a comprehensive review, see Nichols, 2016). There is broad consensus that response to psychedelics depends on a confluence of factors referred to as ‘drug, set, and setting’ (Zinberg, 1986). These include the drug dosage and the user’s psychological profile, current mood state, pre-drug history, response expectancies, and social and environmental context. Psychedelics are generally regarded as non-addictive (Fantegrossi et al., 2004) and as safe when used in an adapted setting and with well-prepared participants (Johnson et al., 2008; Schmid et al., 2015; Studerus et al., 2011). Moreover, large population studies suggest that their long-term use is not associated with increased risk for mental health problems (Johansen and Krebs, 2015), but rather with reduced psychological distress and suicidality (Hendricks et al., 2015).

Neurophysiology

The neurophysiology of the psychedelic experience has been shown to be characterized by changes in the dynamics and connectivity patterns of resting state networks (RSNs; for reviews, see Barrett and Griffiths, 2017; Dos Santos et al., 2016). Psychedelics induce a global increase in cerebral metabolic rate in frontal regions of the brain (Hermle et al., 1992; Vollenweider et al., 1997). EEG and MEG research suggests that psychedelic states are characterized by a broadband desynchronization of cortical oscillations (Kometer et al., 2015; Muthukumaraswamy et al., 2013), including in several brain regions comprising well-established RSNs such as the default mode network (DMN) (Carhart-Harris et al., 2012, 2016b; Kometer et al., 2015; Muthukumaraswamy et al., 2013; Palhano-Fontes et al., 2015). Alteration of the efficiency of long-range communication

between nodes of the DMN, decreased integrity and increased desegregation of RSNs (Carhart-Harris et al., 2016b; Kometer et al., 2015; Muthukumaraswamy et al., 2013) have been proposed as the mechanisms underlying psychedelic experiences (for reviews, see Barrett and Griffiths, 2017; Dos Santos et al., 2016). Despite some mixed and potentially conflicting evidence, this growing body of research reveals that classic psychedelics produce their effects by modulating cortical and subcortical regions involved in self-awareness, perception, affect processing, and executive and higher cognitive functions (Barrett and Griffiths, 2017; Dos Santos et al., 2016).

Therapeutic potential

Psychedelic-assisted psychotherapy generally consists of one or more sessions during which the patient or participant ingests a predetermined dose of a psychedelic drug in a supportive environment. During the session, the therapist provides generally non-directive support for the subsequent experience of the individual. The individual may experience a range of psychological effects including extreme positive and negative emotions. Over the last decade, older clinical studies have been revisited with modern study designs, highlighting the therapeutic potential of psychedelics (for reviews, see Carhart-Harris and Goodwin, 2017; Liechti, 2017). In particular, psychedelic-assisted psychotherapy seems to hold promise in the treatment of depression (Carhart-Harris et al., 2016a), anxiety disorders (Griffiths et al., 2016; Ross et al., 2016), and nicotine and alcohol addiction (Bogenschutz et al., 2015; Garcia-Romeu et al., 2015; Johnson et al., 2014). Critically, multiple studies have reported an association between therapeutic outcome and measurable mystical-type experiences sometimes induced by the drug, suggesting that the nature and quality of one’s response plays a significant role in therapeutic outcome (Bogenschutz et al., 2015; Garcia-Romeu et al., 2015; Johnson et al., 2014; Roseman et al., 2018). Despite the promising results of these preliminary studies, they suffer from methodological limitations such as small sample sizes, lack of comparison against conventional treatments and the significant challenge of the inability to properly control for the placebo response.

Hypnosis

Hypnosis is a unique interpersonal method in which a therapist or experimenter uses verbal suggestion to modulate the conscious states of a patient or participant. A session typically consists of three phases: an *induction*, in which instructions and suggestions for reduced meta-awareness and absorption in the words of the experimenter are administered alongside the general suggestions to enter hypnosis (Terhune and Cardeña, 2016); a *suggestion* phase, in which one or more suggestions are given to modulate the contents of consciousness; and finally a *de-induction* phase, in which instructions and suggestions are administered in order to elicit a return to normal alertness with the potential administration of posthypnotic suggestions intended to take effect following termination of the session (e.g., Barnier and McConkey, 1999). Particularly among highly suggestible individuals, an induction is frequently accompanied by spontaneous alterations in self-related processing and perception (Pekala and Kumar,

2007). However, the evidence for the importance of hypnotic inductions in responsiveness to suggestions is mixed (Terhune and Cardeña, 2016), thereby highlighting the potential utility of applying suggestion in the absence of a formal induction procedure. In response to specific suggestions, highly suggestible individuals are capable of experiencing pronounced alterations in consciousness including hallucinations, amnesia and cognitive deficits (Barnier et al., 2014). Responses to suggestions are typically accompanied by a lack of authorship over the response, as measured by both self-report and implicit perceptual indices (e.g., Lush et al., 2017; Polito et al., 2015), with reductions in the sense of agency being comparable in magnitude in highly suggestible individuals to phenomenological aberrations reported by patients with schizophrenia during passivity symptoms (Polito et al., 2015). Hypnosis can be conceptualized as a manifestation of the broader phenomena of suggestion and suggestibility (Halligan and Oakley, 2014) and has even been referred to as a 'non-deceptive placebo' (Kirsch, 1994). Response expectancies are known to be important contributing factors to responsiveness to both hypnotic suggestions and placebos (De Pascalis et al., 2002; Lynn et al., 2008). However, the available evidence indicates that a relationship between hypnosis and placebo, if one exists, is potentially complex and moderated by multiple factors (De Pascalis et al., 2002; De Pascalis and Scacchia, 2016; McGlashan et al., 1969; Woody et al., 1997).

As measured by standard, and well-validated, behavioural scales (for a review, see Woody and Barnier, 2008), hypnotic suggestibility is normally distributed with approximately 10–15% of the population displaying low and high suggestibility and the remainder exhibiting moderate responsiveness (Laurence et al., 2008). Hypnotic suggestibility exhibits trait-like stability over long periods of time (Piccione et al., 1989) and is at least partly hereditary (Morgan et al., 1970; Rominger et al., 2014). Nevertheless, hypnotic suggestibility has few known personality correlates, typically pertaining to absorption in activities, heightened responsiveness to social-emotional cues and a propensity for self-transcendent experiences, although these effects are typically small in magnitude (Cardeña et al., 2009; Cardeña and Terhune, 2014; Lynn et al., 2015; Wickramasekera and Szlyk, 2003). Similarly, there are no known robust cognitive correlates of hypnotic suggestibility (Parris, 2017), although highly suggestible individuals seem to display heightened automaticity (Braffman and Kirsch, 2001; Dixon and Laurence, 1992), poorer working and short-term memory (Farvolden and Woody, 2004; Khodaverdi-Khani and Laurence, 2016; Terhune et al., 2011) and selectively impaired metacognition (Lush et al., 2016; Terhune and Hedman, 2017). Although hypnosis shares superficial similarity to meditation, recent research suggests that they are actually opposing metacognitive phenomena (Lush et al., 2016).

Neurophysiology

During the past two decades, researchers have utilized neuroimaging techniques to study the neurophysiological correlates of response to hypnotic inductions and suggestions and individual differences in hypnotic suggestibility (Oakley and Halligan, 2013; Terhune et al., 2017). Although the field was traditionally occupied with the question of whether hypnosis constitutes an altered state of consciousness, contemporary research has directed attention toward more substantive questions pertaining

to the correlates, characteristics, and mechanisms of response to hypnotic suggestions and individual differences in hypnotic suggestibility (Jensen et al., 2017). Accumulating data suggest that a hypnotic induction may produce a reduction in global, frontal or frontal-parietal functional connectivity, as measured by resting state EEG (for a review, see Terhune et al., 2017). Functional neuroimaging research has further shown that hypnosis seems to produce a selective reduction in activity of the anterior medial prefrontal cortex (McGeown et al., 2009; see also Deeley et al., 2012), which corresponds to the anterior node of the DMN (Greicius et al., 2009). Other research has similarly observed that the perception of being deeply hypnotized following an induction was associated with reduced functional connectivity between posterior cingulate cortex (PCC) and dorsolateral prefrontal cortex (DLPFC), decreased anterior cingulate cortex (ACC) activation and increased connectivity between DLPFC and insula (Jiang et al., 2017). These results suggest that a hypnotic induction produces in highly suggestible individuals a reduction in self-related or metacognitive processing (Cardeña et al., 2013; Pekala and Kumar, 2007), coupled with atypical connectivity of the executive control network, which may reflect cognitive control with reduced awareness (Dienes and Perner, 2007). Most imaging studies indicate that hypnotic suggestions for altered perception engage neural systems that overlap with those of the genuine experience: for instance, suggestions for colour hallucinations modulate fusiform areas including V4 (Kosslyn et al., 2000; McGeown et al., 2012) and suggestions for altered pain perception alter somatosensory areas and ACC (Derbyshire et al., 2004; Rainville et al., 1997), although the neurophysiological correlates shared by responses to different hypnotic suggestions remain elusive (Oakley and Halligan, 2013; Terhune et al., 2017).

Therapeutic potential

In a clinical setting, hypnosis involves the administration of verbal suggestions and metaphors to guide individuals into dynamic, multimodal experiences with the aim of promoting emotional catharsis and desirable changes in perceptual experiences, self-image, behaviours, habits and general health as appropriate to the presenting symptom or condition (Lynn et al., 2010). Treatment strategies with hypnosis can include direct suggestion, symptom substitution, ego-strengthening and hypnoanalytic therapies (Elkins, 2017). The clinical efficacy of hypnosis is only weakly predicted by hypnotic suggestibility (Montgomery et al., 2011), plausibly because responsiveness to therapeutic suggestions is influenced by a range of non-hypnotic factors, such as patient motivation and response expectancies (e.g., Lynn et al., 2008). Moreover, many suggestions utilized in the therapeutic application of hypnosis do not require a high level of hypnotic suggestibility (Elkins, 2017). Accordingly, medium suggestible individuals are capable of responding, and potentially benefiting from, suggestions administered within clinical contexts (Lynn et al., 2010). Verbal suggestion can also be applied in the absence of hypnosis in a variety of contexts (Amigó and Ferrández, 2015; Peerdeman et al., 2016), thereby potentially removing the need for the explicit use of hypnosis *per se* (see also Terhune and Cardeña, 2016). Although the most recognized use of hypnosis is in the treatment or management of pain, such as in surgical contexts (Faymonville et al., 1995), there is reliable evidence for its efficacy in treating a range of conditions including acute and

chronic pain (Patterson and Jensen, 2003; Tome-Pires and Miro, 2012), post-menopausal hot flashes (Elkins et al., 2013), irritable bowel syndrome (Schaefer et al., 2014) and enhancement of immunological functions (Miller and Cohen, 2001). Although less robust, promising, albeit preliminary, evidence suggests that hypnosis constitutes a potentially valuable option for treating depression (Alladin and Alibhai, 2007), anxiety (Hammond, 2010), nicotine addiction (Hasan et al., 2014), symptoms inherent to neurodegenerative disorders such as amyotrophic lateral sclerosis (Kleinbub et al., 2015) and dermatological problems (Shenefelt, 2000; for reviews, see Elkins, 2017; Terhune et al., 2017). Moreover, when hypnosis is used as an adjunct to non-hypnotic methods, such as cognitive behavioural therapy, therapeutic outcomes are often strongly amplified (Kirsch et al., 1995). In addition, preliminary evidence suggests that suggestibility may influence response to non-suggestion-based treatments (e.g., Nitzan et al., 2015), thus further highlighting the widespread presence and influence of suggestions in a variety of therapeutic contexts. Cumulatively, the available evidence indicates that hypnosis and suggestion-based interventions constitute safe (Bollinger, 2018) and efficacious interventions for treating a range of psychopathological and somatic symptoms.

Psychedelics and hypnosis

During the 1960–1970s, several studies compared phenomenological parallels between psychedelics and hypnosis (Barber, 1970; Grünholz, 1971; Gubel, 1962; Halpern, 1961; Krippner, 1964) and explored the potential benefits of combining them in psychotherapy (Levine and Ludwig, 1966; Ludwig and Levine, 1965; Ludwig et al., 1969). Particularly relevant is the finding that psychedelics seem to enhance suggestibility (Middlefell, 1967; Netz and Engstam, 1968; Sjöberg and Hollister, 1965; Solursh and Rae, 1966; Ulett et al., 1972; Van Nuys, 1972). Other research reported on the use of hypnosis to recreate psychedelic-like experiences (Aaronson, 1970; Baumann, 1970; Erickson, 1965; Tart, 1967; Fogel and Hoffer, 1962) or to control, guide and deepen LSD-induced psychedelic experiences (Fogel and Hoffer, 1962; Levine and Ludwig, 1965; Levine et al., 1963) and to influence the subjects' experiential response to psychedelics with explicit suggestions (Levis and Mehlman, 1964). Here we revisit this seminal work and relate it to recent advances in our knowledge of these phenomena.

Hypnodelic treatment technique

Based on previous research on the potential of hypnosis in modulating the effects of LSD (Fogel and Hoffer, 1962), Ludwig and Levine hypothesised that hypnosis could be used in conjunction with LSD to enhance therapeutic outcome, an approach known as the *hypnodelic treatment technique* (Levine and Ludwig, 1966; Ludwig et al., 1969; Ludwig and Levine, 1965). They evaluated the coupling of these approaches in a randomized controlled trial in which 70 participants with drug addictions were randomly assigned to one of five treatment conditions: (a) LSD + hypnosis + psychotherapy (hypnodelic treatment); (b) LSD + psychotherapy; (c) LSD alone; (d) psychotherapy; or (e) hypnosis + psychotherapy. In the hypnodelic treatment, a 45-minute induction was administered just after drug administration in order to temporally coincide the onset of the therapy with the onset of the

psychedelic state. They reported that participants who received hypnodelic treatment showed greater improvement than those in the other conditions both at 2 weeks and 2 months post-treatment (Ludwig and Levine, 1965). All patients had responded positively on simple suggestibility assessments that ensured a moderate level of hypnotic suggestibility. Hypnotic suggestibility did not differ across treatment groups although it did seem to contribute to therapeutic outcomes. Ludwig and Levine proposed that the enhanced efficacy of hypnodelic treatment technique might be attributable to the mental calmness arising from hypnosis that may have better prepared patients to experience the effects of LSD, patients' enhanced acceptance of, and control over, the psychedelic experiences, greater participation in the therapeutic process, and deeper and more intense experiences afforded with this approach (Levine and Ludwig, 1966; Ludwig and Levine, 1965). A second clinical trial on alcoholism failed to report any beneficial effects with conditions (a) to (c) (Ludwig et al., 1969), with the reason for this failure being unclear. Although preliminary and limited from a methodological perspective, this line of research points to the potential utility of coupling psychedelics and hypnosis, an approach that is likely to benefit considerably from recent advances in our understanding of both phenomena. In the following section, we review recent advances in neuropsychological and clinical research pointing to overlapping characteristics of psychedelics and hypnosis.

Phenomenology

Levine and Ludwig (1965) reported that LSD and hypnotic experiences share overlapping phenomenology and this observation is borne out in recent research. In the context of neutral hypnosis, with the only suggestion being to go as deeply into hypnosis as possible, participants frequently report experiential responses strikingly similar to the phenomenology of psychedelic experiences, such as alterations in perception, body image, imagery, self-awareness, affect, time perception and meaning (Cardeña, 2005; Cardeña et al., 2013). Beyond the spontaneous experiential response to an induction, it has been shown that hypnotic suggestion can be used to experimentally induce psychological states that closely resemble those experienced in response to psychoactive substances including psychedelics (Baumann, 1970; Fogel and Hoffer, 1962), narcotics (Ludwig and Lyle, 1964) and MDMA (Hastings, 2006). Similarly, hypnotic suggestions can be used to elicit mystical-type experiences in highly suggestible individuals (Lynn and Evans, 2017; Sacerdote, 1977; Spanos and Moretti, 1988). Lynn and Evans (2017) reported that 22% of participants (undergraduate volunteers) experienced a 'complete' mystical-type experience, based on the criterion proposed by Barrett et al. (2015). In the same way as absorption predicts mystical-type experience with psychedelics, hypnotic suggestibility correlates with the propensity to experience hypnotically suggested mystical-type experiences (Lynn and Evans, 2017; Spanos and Moretti, 1988). Although a recent investigation suggests that non-psychedelic-induced mystical-type experiences tend to be less intense and associated with lower positive existential impact than psychedelic-induced experiences (Yaden et al., 2017), it has not yet been determined whether and to what extent hypnotically induced mystical-type experiences exert long-term beneficial effects on individuals.

LSD and mescaline, but not psilocybin, seem to enhance suggestibility to a similar extent as a hypnotic induction (Carhart-Harris et al., 2015; Netz and Engstam, 1968; Sjöberg and Hollister, 1965; Solursh and Rae, 1966; Ulett et al., 1972). The mechanisms underlying this augmentation remain unknown, but it may be due to a reduction in reality testing, metacognition or modulation of executive functioning. Nonpharmacological factors such as expectation, preparation, intention, and physical and social environment all influence response to hypnosis and psychedelics (Kirsch, 2000; Zinberg, 1986). In addition, there is evidence for a shared psychological propensity for response to both phenomena: absorption, the individual tendency to experience episodes of intense attentional involvement (Tellegen and Atkinson, 1974), appears to be a reliable predictor of phenomenal response to psychedelics (Studerus et al., 2012) as well as hypnotic suggestibility (Cardeña and Terhune, 2014; but see Council et al., 1986; Roche and McConkey, 1990). Despite these commonalities, psychedelics and hypnosis clearly differ in their mode of induction: the former being psychopharmacological and the latter being suggestion-based and occurring typically in an interpersonal context. The two phenomena are also temporally different: a psychedelic experience is limited in time relative to the metabolism and clearance of the drug from the body whereas a hypnotic experience can be easily terminated by the participant or therapist/experimenter at any time (although see Perry, 1977).

Neurophysiology

Neuroimaging studies have shown that both psychedelics and hypnosis affect the activity of several brain areas and networks, including prefrontal regions, cingulate cortex and the DMN (Carhart-Harris et al., 2012, 2016b; Deeley et al., 2012; Hermle et al., 1992; Jiang et al., 2017; McGeown et al., 2009; Palhano-Fontes et al., 2015; Pekala and Kumar, 2007; Vollenweider et al., 1997). Psychedelics seem to affect a larger set of brain regions and networks than hypnosis and induce broader changes specifically in frontal regions, medial temporal lobe, occipital cortex, hippocampus and amygdala (Barrett and Griffiths, 2017; Dos Santos et al., 2016). In addition, spontaneous phenomenological effects shared by the two phenomena seem to be associated with converse global functional connectivity patterns. For example, ego-dissolution under the influence of psychedelics was associated with increased connectivity (Tagliazucchi et al., 2016) whereas transcendent experiences following a hypnotic induction were associated with lower connectivity (Cardeña et al., 2013). One possibility for these discrepancies is that the relation between anomalous self-awareness and global functional connectivity adheres to an inverted-U shape with greater aberrations occurring with both lower and higher connectivity patterns. Alternatively, these differences may reflect distinct alterations in the experience of the self in these different conditions. Similarities in spontaneous experiential response to psychedelics and hypnosis may be explained in part by the modulation of the DMN and global functional connectivity. Prepulse inhibition (PPI) of the acoustic startle response, a parameter thought to reflect the early, pre-attentive stage of automatic sensory processing used to assess efficiency of the sensorimotor gating system, is disrupted in response to psilocybin (Vollenweider et al., 2007) and LSD (Schmid et al., 2015) and seems to be lower in highly suggestible individuals (Levin et al., 2011; Lichtenberg et al., 2008; Storozheva et al., 2018; but see De Pascalis and Russo,

2013). These results suggest a potentially overlapping mechanism underlying individual differences in hypnotic suggestibility and responsiveness to psychedelics, which may relate to the psychedelic enhancement of suggestibility (Carhart-Harris et al., 2015). Although the neurophysiological basis of hypnotically induced mystical-type experiences is unknown (Lynn and Evans, 2017), hypnotic suggestion often, but not always, produces a similar neurophysiological pattern as the actual phenomenon (Oakley and Halligan, 2013; Terhune et al., 2017), suggesting the viability of using hypnotic suggestion to reproduce psychedelics-occasioned mystical-type experiences (Barrett and Griffiths, 2017). Neurophenomenological parallels between hypnosis and psychedelics seem to offer a valuable platform where these two phenomena could be mutually beneficial to each other.

Future directions

Recent research on psychedelics and hypnosis allows for revisiting their combined use in psychotherapy. Enhanced suggestibility under psychedelics and hypnosis may have implications for their use as adjuncts in psychotherapy, where suggestibility plays a significant role in therapeutic outcomes (Montgomery et al., 2011). Researchers and clinicians may already be harnessing the power of suggestion to modulate set and setting and, in turn, to influence the phenomenological response to psychedelics. Harnessing the effects of suggestion more explicitly may facilitate the development of a more precise understanding of the pharmacological and contextual factors influencing the psychedelic experience and a way to further enhance their therapeutic efficacy. In what follows, we propose several pathways concerning the combined use of hypnosis and psychedelics to prepare, guide and integrate the psychedelic experience in order to enhance treatment efficacy.

Hypnosis-based training for participant preparation

Experience of hypnotically suggested alterations in consciousness, including phenomenal states resembling psychedelic experiences, could be used as a preliminary training regimen to familiarize naïve participants with the experiential effects of psychedelics in a controlled setting. Such training may better prepare individuals to participate in psychedelic research or therapy and could potentially reduce pre- and/or post-treatment anxiety, and participant attrition, whilst augmenting positive response expectancies and involvement in the psychedelic experience. These complementary effects could increase the likelihood of safe and profound experiences and, concomitantly, successful therapeutic outcomes. Moreover, we expect that similarly to absorption (Studerus et al., 2012), hypnotic suggestibility could constitute a valuable predictor of perceptual, cognitive, affective and self-related phenomenological features of the psychedelic experience.

Hypnodelic treatment technique: Using suggestion to guide psychedelic states

Ludwig and Levine reported that the conjunct use of hypnosis and psychedelics enabled them to enhance participants' experiential responses to LSD, rendering the experience more intense and malleable to control with potentially increased therapeutic efficacy

(Levine and Ludwig, 1965; Ludwig and Levine, 1965). Moreover, as psychedelics seem to increase suggestibility (Carhart-Harris et al., 2015; Middlefell, 1967; Netz and Engstam, 1968; Sjöberg and Hollister, 1965; Solursh and Rae, 1966; Ulett et al., 1972; Van Nuys, 1972), therapeutically coupling the two could further potentiate the impact of suggestions. Harnessing the power of suggestion to modulate response to psychedelics could exert considerable effects in enhancing their therapeutic efficacy including by guiding the psychedelic experience to increase the likelihood of positive responses, promote, or enhance, mystical-type experiences (e.g., Lynn and Evans, 2017) and avoid unwanted experiences. Moreover, posthypnotic suggestions, including suggestions for how the experience could be processed afterward in terms of meaning, values and future life directions, may further aid integration of the experience and carry over to everyday life. For these reasons, we maintain that a renewed attention should be directed to the development of a formalized suggestion-based psychedelic-assisted psychotherapy.

Hypnosis and mild dosages of psychedelics: Transversal applications

Drug dosage reduction for equivalent treatment outcome is an important goal in psychopharmacology. At present, little is known about the neurophenomenological effects (Greiner et al., 1958), and/or therapeutic potential, of psychedelics at low dosages and the dose level required to enhance suggestibility is unknown. Insofar as hypnotic suggestion may be useful in augmenting classical psychedelic experiences (Levine and Ludwig, 1965), it would be valuable to determine whether such an approach could be extended to mild dosages in order to evaluate whether suggestion can be used to similarly enhance weak psychedelic experiences.

Hypnosis for drug-free re-experiencing of psychedelic states

Several studies have shown that hypnosis can be used to induce mystical-type experiences (Lynn and Evans, 2017; Sacerdote, 1977; Spanos and Moretti, 1988). A potential benefit of combining psychedelics and hypnosis could be to use suggestions to reproduce such experiences in the days after administration of psychedelics (e.g., Fogel and Hoffer, 1962; Hastings, 2006). This approach may afford the potential for patients to re-experience the associated affective states, with greater control over the post-administration response to psychedelic-assisted therapy, thereby providing additional opportunities to harness the positive features of psychedelic experiences for therapeutic impact, with the potential to further enhance outcomes over a sustained period.

Conclusion

Recent clinical research on psychedelic-assisted psychotherapy and hypnosis demonstrates the value of both methods in the treatment of a range of conditions with overlapping applications and neurophenomenological features. Older research on the overlap of these two phenomena considered against the backdrop of recent advances clearly points to the potential of harnessing the power of suggestion to influence the phenomenological response to psychedelics. The coupling of psychedelics and hypnosis represents a potentially highly fertile

platform for both basic research on alterations in consciousness as well as their exploitation in therapy and thereby warrants renewed empirical attention.

Acknowledgements

The authors would like to thank Neiloufar Family and anonymous reviewers for constructive comments on an earlier draft of this manuscript.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: DBT acknowledges the support of bursary 70/16 from the Bial Foundation.

ORCID iD

Clément E. Lemerrier  <https://orcid.org/0000-0002-7113-3543>

References

- Aaronson BS (1970) Some hypnotic analogues to the psychedelic state. In: Aaronson BS and Osmond H (eds) *Psychedelics: The Uses and Implications of Hallucinogenic Drugs*. New York: Anchor, pp. 279–295.
- Alladin A and Alibhai A (2007) Cognitive hypnotherapy for depression: An empirical investigation. *Int J Clin Exp Hypn* 55: 147–166.
- Amigó S and Ferrández C (2015) Experiencing effects of cocaine and speed with self-regulation therapy. *Span J Psychol* 18: E49.
- Barber TX (1970) *LSD, Marihuana, Yoga, and Hypnosis*. Chicago: Aldine.
- Barnier AJ and McConkey KM (1999) Hypnotic and posthypnotic suggestion: Finding meaning in the message of the hypnotist. *Int J Clin Exp Hypn* 47: 192–208.
- Barnier AJ, Cox RE and McConkey KM (2014) The province of ‘highs’: The high hypnotizable person in the science of hypnosis and in psychological science. *Psychol Conscious* 1: 168–183.
- Barrett FS, Johnson MW and Griffiths RR (2015) Validation of the revised Mystical Experience Questionnaire in experimental sessions with psilocybin. *J Psychopharmacol* 29: 1182–1190.
- Barrett FS and Griffiths RR (2017) Classic hallucinogens and mystical experiences: Phenomenology and neural correlates. *Curr Top Behav Neurosci* 36: 393–430.
- Baumann F (1970) Hypnosis and the adolescent drug abuser. *Am J Clin Hypn* 13: 17–21.
- Bogenschutz MP, Forcehimes AA, Pommy JA, et al. (2015) Psilocybin-assisted treatment for alcohol dependence: A proof-of-concept study. *J Psychopharmacol* 29: 289–299.
- Bollinger JW (2018) The rate of adverse events related to hypnosis during clinical trials. *Am J Clin Hypn* 60: 357–366.
- Braffman W and Kirsch I (2001) Reaction time as a predictor of imaginative suggestibility and hypnotizability. *Contemp Hypn* 18: 107–120.
- Buckley P and Galanter M (1979) Altered states of consciousness during psychotherapy: A historical and cultural perspective. *Int J Soc Psychiatry* 25: 118–124.
- Cardeña E (2005) The phenomenology of deep hypnosis: Quiescent and physically active. *Int J Clin Exp Hypn* 53: 37–59.
- Cardeña E and Terhune DB (2014) Hypnotizability, personality traits, and the propensity to experience alterations of consciousness. *Psychol Conscious* 1: 292–307.

- Cardeña E, Jonsson P, Terhune DB, et al. (2013) The neurophenomenology of neutral hypnosis. *Cortex* 49: 375–385.
- Cardeña E, Terhune DB, Loof A, et al. (2009) Hypnotic experience is related to emotional contagion. *Int J Clin Exp Hypn* 57: 33–46.
- Carhart-Harris RL and Goodwin GM (2017) The therapeutic potential of psychedelic drugs: Past, present, and future. *Neuropsychopharmacology* 42: 2105–2113.
- Carhart-Harris RL, Bolstridge M, Rucker J, et al. (2016a) Psilocybin with psychological support for treatment-resistant depression: An open-label feasibility study. *Lancet Psychiatry* 3: 619–627.
- Carhart-Harris RL, Erritzoe D, Williams T, et al. (2012) Neural correlates of the psychedelic state as determined by fMRI studies with psilocybin. *Proc Natl Acad Sci USA* 109: 2138–2143.
- Carhart-Harris RL, Kaelen M, Whalley MG, et al. (2015) LSD enhances suggestibility in healthy volunteers. *Psychopharmacology (Berl)* 232: 785–794.
- Carhart-Harris RL, Muthukumaraswamy S, Roseman L, et al. (2016b) Neural correlates of the LSD experience revealed by multimodal neuroimaging. *Proc Natl Acad Sci USA* 113: 4853–4858.
- Council JR, Kirsch I and Hafner LP (1986) Expectancy versus absorption in the prediction of hypnotic responding. *J Pers Soc Psychol* 50: 182–189.
- De Araujo DB, Ribeiro S, Cecchi GA, et al. (2012) Seeing with the eyes shut: Neural basis of enhanced imagery following Ayahuasca ingestion. *Hum Brain Mapp* 33: 2550–2560.
- Deeley Q, Oakley DA, Toone B, et al. (2012) Modulating the default mode network using hypnosis. *Int J Clin Exp Hypn* 60: 206–228.
- De Pascalis V and Russo E (2013) Hypnotizability, hypnosis and pre-pulse inhibition of the startle reflex in healthy women: An ERP analysis. *PLoS One* 8: e79605.
- De Pascalis V and Scacchia P (2016) Hypnotizability and placebo analgesia in waking and hypnosis as modulators of auditory startle responses in healthy women: An ERP study. *PLoS One* 11: e0159135.
- De Pascalis V, Chiaradia C and Carotenuto E (2002) The contribution of suggestibility and expectation to placebo analgesia phenomenon in an experimental setting. *Pain* 96: 393–402.
- Derbyshire SW, Whalley MG, Stenger VA, et al. (2004) Cerebral activation during hypnotically induced and imagined pain. *Neuroimage* 23: 392–401.
- Dienes Z and Perner J (2007) Executive control without conscious awareness: The cold control theory of hypnosis. In: Jamieson GA (ed.) *Hypnosis and Conscious States: The Cognitive Neuroscience Perspective*. Oxford: Oxford University Press, pp. 293–314.
- Dixon M and Laurence JR (1992) Hypnotic susceptibility and verbal automaticity: Automatic and strategic processing differences in the Stroop color-naming task. *J Abnorm Psychol* 101: 344–347.
- Dos Santos RG, Osorio FL, Crippa JA, et al. (2016) Classical hallucinogens and neuroimaging: A systematic review of human studies: Hallucinogens and neuroimaging. *Neurosci Biobehav Rev* 71: 715–728.
- Elkins G (2017) *Handbook of Medical and Psychological Hypnosis: Foundations, Applications, and Professional Issues*. New York, NY: Springer.
- Elkins GR, Fisher WI, Johnson AK, et al. (2013) Clinical hypnosis in the treatment of postmenopausal hot flashes: A randomized controlled trial. *Menopause* 20: 291–298.
- Ellenberger HF (1970) *The Discovery of the Unconscious: The History and Evolution of Dynamic Psychiatry*. New York: Basic Books, Inc.
- Erickson MH (1965) A special inquiry with Aldous Huxley into the nature and character of various altered states of consciousness. *Am J Clin Hypn* 8: 17–33.
- Fantegrossi WE, Woods JH and Winger G (2004) Transient reinforcing effects of phenylisopropylamine and indolealkylamine hallucinogens in rhesus monkeys. *Behav Pharmacol* 15: 149–157.
- Farvolden P and Woody EZ (2004) Hypnosis, memory, and frontal executive functioning. *Int J Clin Exp Hypn* 52: 3–26.
- Faymonville ME, Fissette J, Mambourg PH, et al. (1995) Hypnosis as adjunct therapy in conscious sedation for plastic surgery. *Reg Anesth* 20: 145–151.
- Fogel S and Hoffer A (1962) The use of hypnosis to interrupt and to reproduce an LSD-25 experience. *J Clin Exp Psychopathol Q Rev Psychiatry Neurol* 23: 11–16.
- Garcia-Romeu A, Griffiths RR and Johnson MW (2015) Psilocybin-occasioned mystical experiences in the treatment of tobacco addiction. *Curr Drug Abuse Rev* 7: 157–164.
- González-Maeso J, Weisstaub NV, Zhou M, et al. (2007) Hallucinogens recruit specific cortical 5-HT(2A) receptor-mediated signaling pathways to affect behavior. *Neuron* 53: 439–452.
- Greicius MD, Supekar K, Menon V, et al. (2009) Resting-state functional connectivity reflects structural connectivity in the default mode network. *Cereb Cortex* 19: 72–78.
- Greiner T, Burch NR and Edelberg R (1958) Psychopathology and psychophysiology of minimal LSD-25 dosage; a preliminary dosage-response spectrum. *AMA Arch Neurol Psychiatry* 79: 208–210.
- Griffiths RR, Johnson MW, Carducci MA, et al. (2016) Psilocybin produces substantial and sustained decreases in depression and anxiety in patients with life-threatening cancer: A randomized double-blind trial. *J Psychopharmacol* 30: 1181–1197.
- Griffiths RR, Johnson MW, Richards WA, et al. (2011) Psilocybin occasioned mystical-type experiences: Immediate and persisting dose-related effects. *Psychopharmacology (Berl)* 218: 649–665.
- Griffiths R, Richards W, Johnson M, et al. (2008) Mystical-type experiences occasioned by psilocybin mediate the attribution of personal meaning and spiritual significance 14 months later. *J Psychopharmacol* 22: 621–632.
- Grünholz G (1971) [Autohypnosis replaces LSD]. *Z Allgemeinmed* 47: 856–859.
- Gubel I (1962) Hallucinogenic drugs and hypnosis in psychotherapy. *Am J Clin Hypn* 4: 169–173.
- Halligan PW and Oakley DA (2014) Hypnosis and beyond: Exploring the broader domain of suggestion. *Psychol Conscious* 1: 105–122.
- Halpern S (1961) On the similarity between hypnotic and mescaline hallucinations. *Int J Clin Exp Hypn* 9: 139–149.
- Hammond DC (2010) Hypnosis in the treatment of anxiety- and stress-related disorders. *Expert Rev Neurother* 10: 263–273.
- Hasan FM, Zagarins SE, Pischke KM, et al. (2014) Hypnotherapy is more effective than nicotine replacement therapy for smoking cessation: Results of a randomized controlled trial. *Complement Ther Med* 22: 1–8.
- Hastings A (2006) An extended nondrug MDMA-like experience evoked through posthypnotic suggestion. *J Psychoactive Drugs* 38: 273–283.
- Hendricks PS, Thorne CB, Clark CB, et al. (2015) Classic psychedelic use is associated with reduced psychological distress and suicidality in the United States adult population. *J Psychopharmacol* 29: 280–288.
- Hermle L, Funfgeld M, Oepen G, et al. (1992) Mescaline-induced psychopathological, neuropsychological, and neurometabolic effects in normal subjects: Experimental psychosis as a tool for psychiatric research. *Biol Psychiatry* 32: 976–991.
- Jensen MP, Jamieson GA, Lutz A, et al. (2017) New directions in hypnosis research: Strategies for advancing the cognitive and clinical neuroscience of hypnosis. *Neurosci Conscious* 3: pii: nix004.
- Jiang H, White MP, Greicius MD, et al. (2017) Brain activity and functional connectivity associated with hypnosis. *Cereb Cortex* 27: 4083–4093.
- Johansen PO and Krebs TS (2015) Psychedelics not linked to mental health problems or suicidal behavior: A population study. *J Psychopharmacol* 29: 270–279.

- Johnson MW, Garcia-Romeu A, Cosimano MP, et al. (2014) Pilot study of the 5-HT_{2A}R agonist psilocybin in the treatment of tobacco addiction. *J Psychopharmacol* 28: 983–992.
- Johnson M, Richards W and Griffiths R (2008) Human hallucinogen research: Guidelines for safety. *J Psychopharmacol* 22: 603–620.
- Khodaverdi-Khani M and Laurence J-R (2016) Working memory and hypnotizability. *Psychol Conscious* 3: 80–92.
- Kirsch I (1994) Clinical hypnosis as a nondeceptive placebo: Empirically derived techniques. *Am J Clin Hypn* 37: 95–106.
- Kirsch I (2000) The response set theory of hypnosis. *Am J Clin Hypn* 42: 274–292.
- Kirsch I, Montgomery G and Sapirstein G (1995) Hypnosis as an adjunct to cognitive-behavioral psychotherapy: A meta-analysis. *J Consult Clin Psychol* 63: 214–220.
- Kleinbub JR, Palmieri A, Broggio A, et al. (2015) Hypnosis-based psychodynamic treatment in ALS: A longitudinal study on patients and their caregivers. *Front Psychol* 6: 822.
- Kometer M, Pokorny T, Seifritz E, et al. (2015) Psilocybin-induced spiritual experiences and insightfulness are associated with synchronization of neuronal oscillations. *Psychopharmacology (Berl)* 232: 3663–3676.
- Kosslyn SM, Thompson WL, Costantini-Ferrando MF, et al. (2000) Hypnotic visual illusion alters color processing in the brain. *Am J Psychiatry* 157: 1279–1284.
- Kraehenmann R, Preller KH, Scheidegger M, et al. (2015) Psilocybin-induced decrease in amygdala reactivity correlates with enhanced positive mood in healthy volunteers. *Biol Psychiatry* 78: 572–581.
- Krippner S (1964) The hypnotic trance, the psychedelic experience, and the creative act. *Am J Clin Hypn* 7: 140–147.
- Laurence JR, Beaulieu-Prévost D and Du Chéné T (2008) Measuring and understanding individual differences in hypnotizability. In: Nash M and Barnier A (eds) *The Oxford Handbook of Hypnosis: Theory, Research, and Practice*. Oxford: Oxford University Press, pp. 225–253.
- Levin R, Heresco-Levy U, Edelman S, et al. (2011) Hypnotizability and sensorimotor gating: A dopaminergic mechanism of hypnosis. *Int J Clin Exp Hypn* 59: 399–405.
- Levine J and Ludwig AM (1965) Alterations in consciousness produced by combinations of LSD, hypnosis and psychotherapy. *Psychopharmacologia* 7: 123–137.
- Levine J and Ludwig AM (1966) The hypnodelic treatment technique. *Int J Clin Exp Hypn* 14: 207–213.
- Levine J, Ludwig AM and Lyle WH Jr (1963) The controlled psychedelic state. *Am J Clin Hypn* 6: 163–164.
- Levis DJ and Mehlman B (1964) Suggestion and mescaline sulphate. *J Neuropsychiatr* 5: 197–200.
- Lichtenberg P, Even-Or E, Bar G, et al. (2008) Reduced prepulse inhibition is associated with increased hypnotizability. *Int J Neuropsychopharmacol* 11: 541–545.
- Liechti ME (2017) Modern clinical research on LSD. *Neuropsychopharmacology* 42: 2114–2127.
- Ludwig AM and Levine J (1965) A controlled comparison of five brief treatment techniques employing LSD, hypnosis, and psychotherapy. *Am J Psychother* 19: 417–435.
- Ludwig AM and Lyle WH Jr (1964) The experimental production of narcotic drug effects and withdrawal symptoms through hypnosis. *Int J Clin Exp Hypn* 12: 1–17.
- Ludwig A, Levine J, Stark L, et al. (1969) A clinical study of LSD treatment in alcoholism. *Am J Psychiatry* 126: 59–69.
- Lush P, Caspar EA, Cleeremans A, et al. (2017) The power of suggestion: Posthypnotically induced changes in the temporal binding of intentional action outcomes. *Psychol Sci* 28: 661–669.
- Lush P, Naish P and Dienes Z (2016) Metacognition of intentions in mindfulness and hypnosis. *Neurosci Conscious* 2016: 1–10.
- Lynn SJ and Evans J (2017) Hypnotic suggestion produces mystical-type experiences in the laboratory: A demonstration proof. *Psychol Conscious* 4: 23–37.
- Lynn SJ, Kirsch I and Hallquist M (2008) Social cognitive theories of hypnosis. In: Nash M and Barnier A (eds) *The Oxford Handbook of Hypnosis: Theory, Research, and Practice*. Oxford: Oxford University Press, pp. 111–140.
- Lynn SJ, Laurence JR and Kirsch I (2015) Hypnosis, suggestion, and suggestibility: An integrative model. *Am J Clin Hypn* 57: 314–329.
- Lynn SJ, Rhue JW and Kirsch I (2010) *Handbook of Clinical Hypnosis*. 2nd ed. Washington, DC: American Psychological Association.
- MacLean KA, Johnson MW and Griffiths RR (2011) Mystical experiences occasioned by the hallucinogen psilocybin lead to increases in the personality domain of openness. *J Psychopharmacol* 25: 1453–1461.
- McGeown WJ, Mazzoni G, Venneri A, et al. (2009) Hypnotic induction decreases anterior default mode activity. *Conscious Cogn* 18: 848–855.
- McGeown WJ, Venneri A, Kirsch I, et al. (2012) Suggested visual hallucination without hypnosis enhances activity in visual areas of the brain. *Conscious Cogn* 21: 100–116.
- McGlashan TH, Evans FJ and Orne MT (1969) The nature of hypnotic analgesia and placebo response to experimental pain. *Psychosom Med* 31: 227–246.
- Middlefell R (1967) The effects of LSD on body sway suggestibility in a group of hospital patients. *Br J Psychiatry* 113: 277–280.
- Miller GE and Cohen S (2001) Psychological interventions and the immune system: A meta-analytic review and critique. *Health Psychol* 20: 47–63.
- Montgomery GH, Schnur JB and David D (2011) The impact of hypnotic suggestibility in clinical care settings. *Int J Clin Exp Hypn* 59: 294–309.
- Morgan AH, Hilgard ER and Davert EC (1970) The heritability of hypnotic susceptibility of twins: A preliminary report. *Behav Genet* 1: 213–224.
- Muthukumaraswamy SD, Carhart-Harris RL, Moran RJ, et al. (2013) Broadband cortical desynchronization underlies the human psychedelic state. *J Neurosci* 33: 15171–15183.
- Netz B and Engstam PO (1968) *Lysergic Acid Diethylamide (LSD-25) and Suggestibility. Part II: Effects of a Threshold Dosage of LSD-25 on Hypnotic Susceptibility*. Stockholm: Military Psychological Institute, MPI B-rapport nr 17.
- Nichols DE (2016) Psychedelics. *Pharmacol Rev* 68: 264–355.
- Nitzan U, Chalamish Y, Krieger I, et al. (2015) Suggestibility as a predictor of response to antidepressants: A preliminary prospective trial. *J Affect Disord* 185: 8–11.
- Oakley DA and Halligan PW (2013) Hypnotic suggestion: Opportunities for cognitive neuroscience. *Nat Rev Neurosci* 14: 565–576.
- Palhano-Fontes F, Andrade KC, Tofoli LF, et al. (2015) The psychedelic state induced by ayahuasca modulates the activity and connectivity of the default mode network. *PLoS One* 10: e0118143.
- Parris B (2017) The role of frontal executive functions in hypnosis and hypnotic suggestibility. *Psychol Conscious* 4: 211–229.
- Patterson DR and Jensen MP (2003) Hypnosis and clinical pain. *Psychol Bull* 129: 495–521.
- Peerdeman KJ, van Laarhoven AI, Keij SM, et al. (2016) Relieving patients' pain with expectation interventions: A meta-analysis. *Pain* 157: 1179–1191.
- Pekala RJ and Kumar VK (2007) An empirical-phenomenological approach to quantifying consciousness and states of consciousness: With particular reference to understanding the nature of hypnosis. In: Jamieson GA (ed.) *Hypnosis and Conscious States: The Cognitive Neuroscience Perspective*. Oxford: Oxford University Press, pp. 167–194.
- Perry C (1977) Variables influencing the posthypnotic persistence of an uncanceled hypnotic suggestion. *Ann N Y Acad Sci* 296: 264–273.

- Piccione C, Hilgard ER and Zimbardo PG (1989) On the degree of stability of measured hypnotizability over a 25-year period. *J Pers Soc Psychol* 56: 289–295.
- Polito V, Langdon R and Barnier AJ (2015) Sense of agency across contexts: Insights from schizophrenia and hypnosis. *Psychol Conscious* 2: 301–314.
- Preller KH and Vollenweider FX (2018) Phenomenology, structure, and dynamic of psychedelic states. *Curr Top Behav Neurosci* 36: 221–256.
- Rainville P, Duncan GH, Price DD, et al. (1997) Pain affect encoded in human anterior cingulate but not somatosensory cortex. *Science* 277: 968–971.
- Roche SM and McConkey KM (1990) Absorption: Nature, assessment, and correlates. *J Pers Soc Psychol* 59: 91–101.
- Rominger C, Weiss EM, Nagl S, et al. (2014) Carriers of the COMT Met/Met allele have higher degrees of hypnotizability, provided that they have good attentional control: A case of gene-trait interaction. *Int J Clin Exp Hypn* 62: 455–482.
- Roseman L, Nutt D and Carhart-Harris RL (2018) Quality of acute psychedelic experience predicts therapeutic efficacy of psilocybin for treatment-resistant depression. *Front Pharmacol* 8: 974.
- Ross S, Bossis A, Guss J, et al. (2016) Rapid and sustained symptom reduction following psilocybin treatment for anxiety and depression in patients with life-threatening cancer: A randomized controlled trial. *J Psychopharmacol* 30: 1165–1180.
- Sacerdote P (1977) Applications of hypnotically elicited mystical states to the treatment of physical and emotional pain. *Int J Clin Exp Hypn* 25: 309–324.
- Schaefer R, Klose P, Moser G, et al. (2014) Efficacy, tolerability, and safety of hypnosis in adult irritable bowel syndrome: Systematic review and meta-analysis. *Psychosom Med* 76: 389–398.
- Schmid Y and Liechti ME (2017) Long-lasting subjective effects of LSD in normal subjects. *Psychopharmacology (Berl)* 235: 535–545.
- Schmid Y, Enzler F, Gasser P, et al. (2015) Acute effects of lysergic acid diethylamide in healthy subjects. *Biol Psychiatry* 78: 544–553.
- Schultes RE and Hofmann A (1979) *Plants of the Gods: Origins of Hallucinogenic Use*. New York: McGraw-Hill.
- Shenefelt PD (2000) Hypnosis in dermatology. *Arch Dermatol* 136: 393–399.
- Sjoberg BM Jr and Hollister LE (1965) The effects of psychotomimetic drugs on primary suggestibility. *Psychopharmacologia* 8: 251–262.
- Solursh LP and Rae JM (1966) LSD, suggestion and hypnosis. *Int J Neuropsychiatry* 2: 60–64.
- Spanos NP and Moretti P (1988) Correlates of mystical and diabolical experiences in a sample of female university students. *J Sci Study Relig* 27: 105–116.
- Storozheva ZI, Kirenskaya AV, Gordeev MN, et al. (2018) COMT genotype and sensory and sensorimotor gating in high and low hypnotizable subjects. *Int J Clin Exp Hypn* 66: 83–105.
- Studerus E, Gamma A, Komater M, et al. (2012) Prediction of psilocybin response in healthy volunteers. *PLoS One* 7: e30800.
- Studerus E, Komater M, Hasler F, et al. (2011) Acute, subacute and long-term subjective effects of psilocybin in healthy humans: A pooled analysis of experimental studies. *J Psychopharmacol* 25: 1434–1452.
- Tagliazucchi E, Roseman L, Kaelen M, et al. (2016) Increased global functional connectivity correlates with LSD-induced ego dissolution. *Curr Biol* 26: 1043–1050.
- Tart CT (1967) Psychedelic experiences associated with a novel hypnotic procedure, mutual hypnosis. *Am J Clin Hypn* 10: 65–78.
- Tellegen A and Atkinson G (1974) Openness to absorbing and self-altering experiences (“absorption”), a trait related to hypnotic susceptibility. *J Abnorm Psychol* 83: 268–277.
- Terhune DB and Cardeña E (2016) Nuances and uncertainties regarding hypnotic inductions: Toward a theoretically informed praxis. *Am J Clin Hypn* 59: 155–174.
- Terhune DB and Hedman LRA (2017) Metacognition of agency is reduced in high hypnotic suggestibility. *Cognition* 168: 176–181.
- Terhune DB, Cardeña E and Lindgren M (2011) Dissociative tendencies and individual differences in high hypnotic suggestibility. *Cogn Neuropsychiatry* 16: 113–135.
- Terhune DB, Cleeremans A, Raz A, et al. (2017) Hypnosis and top-down regulation of consciousness. *Neurosci Biobehav Rev* 81: 59–74.
- Tome-Pires C and Miro J (2012) Hypnosis for the management of chronic and cancer procedure-related pain in children. *Int J Clin Exp Hypn* 60: 432–457.
- Ulett GA, Akpinar S and Itil TM (1972) Hypnosis: Physiological, pharmacological reality. *Am J Psychiatry* 128: 799–805.
- Van Nuys DW (1972) Drug use and hypnotic susceptibility. *Int J Clin Exp Hypn* 20: 31–37.
- Vollenweider FX, Csomor PA, Knappe B, et al. (2007) The effects of the preferential 5-HT_{2A} agonist psilocybin on prepulse inhibition of startle in healthy human volunteers depend on interstimulus interval. *Neuropsychopharmacology* 32: 1876–1887.
- Vollenweider FX, Leenders KL, Scharfetter C, et al. (1997) Positron emission tomography and fluorodeoxyglucose studies of metabolic hyperfrontality and psychopathology in the psilocybin model of psychosis. *Neuropsychopharmacology* 16: 357–372.
- Wickramasekera IE II and Szlyk JP (2003) Could empathy be a predictor of hypnotic ability? *Int J Clin Exp Hypn* 51: 390–399.
- Woody EZ and Barnier A (2008) Hypnosis scales for the twenty-first century: What do we need and how should we use them? In: Nash M and Barnier A (eds) *The Oxford Handbook of Hypnosis: Theory, Research, and Practice*. Oxford: Oxford University Press, pp. 255–281.
- Woody E and Szechtman H (2011) Using hypnosis to develop and test models of psychopathology. *J Mind-Body Regul* 1: 4–16.
- Woody E, Drugovic M and Oakman J (1997) A reexamination of the role of nonhypnotic suggestibility in hypnotic responding. *J Pers Soc Psychol* 72: 399–407.
- Yaden DB, Le Nguyen KD, Kern ML, et al. (2017) Of roots and fruits: A comparison of psychedelic and nonpsychedelic mystical experiences. *J Human Psychol* 57: 338–353.
- Zeev-Wolf M, Dor-Ziderman Y, Goldstein A, et al. (2017) Oscillatory brain mechanisms of the hypnotically-induced out-of-body experience. *Cortex* 96: 19–30.
- Zinberg NE (1986) Drug, set, and setting: The basis for controlled intoxicant use. *Am J Psychiatry* 143: 548–549.