

6. The unstable, sirupy halogeno compounds obtained from the second octaacetate have the regular structures of the α -halogeno-acetylturanoses (VII). Inasmuch as the properties of these substances are extremely different from those of the " α -chloroacetylfructose," it has been concluded that the latter substance cannot be the stereoisomer of the unstable β -chloroacetylfructose.

PRINCETON, NEW JERSEY

[CONTRIBUTION FROM THE EXPERIMENTAL RESEARCH LABORATORIES, BURROUGHS, WELLCOME AND COMPANY]

HYDROXY- AND DIHYDROXYPHENYLETHYLMETHYLAMINES AND THEIR ETHERS

BY JOHANNES S. BUCK


RECEIVED MARCH 28, 1932

PUBLISHED SEPTEMBER 5, 1932

In an earlier paper¹ the synthesis of Epinine, 3,4-dihydroxyphenylethylmethylamine, was described. This, with two other members of the same series, has been examined pharmacologically.² Owing to their pressor activity, it was decided to attempt the synthesis of the remaining members of the series and to carry out a pharmacological examination of these. The compounds recorded up to the present are the unsubstituted amine, the 4-hydroxyamine and the 3,4-dihydroxyamine. Those so far not described are the 2-hydroxy, 3-hydroxy, and the 2,3-, 2,5-, 2,4-, 3,5- and the 2,6-dihydroxyphenylethylmethylamines. Of these, all but the three latter have been synthesized and are described, together with their intermediates, in the present paper. The synthesis of the 2,4-dihydroxy compound broke down at the last stage, the demethylation proceeding abnormally. This reaction will form the subject of another communication. The syntheses of the remaining two members of the series, the 3,5- and 2,6-dihydroxyamines, have not been carried out, the first owing to lack of starting material, and the second owing to difficulties caused by the powerful steric hindrance of the 2,6-groups. It is hoped to complete the series in the future. The pharmacological action will be described in another place.

Experimental

The preparation of the hydroxyphenylethylmethylamines was carried out in all cases by a series of reactions similar to that used in the synthesis of epinine.¹



¹ Buck, *THIS JOURNAL*, **52**, 4119 (1930).

² Barger and Dale, *J. Physiol.*, **41**, 19 (1910).

The cinnamic acid in each case was made by the well-known Knoevenagel method, by condensation of the aldehyde with malonic acid in the presence of pyridine, piperidine being used as catalyst. On reduction with 4% sodium amalgam (excess = 4H being used), the corresponding phenylpropionic acid was produced, usually in very good yield. From the acid, the amide was made by heating in a stream of ammonia at 220–230° for two hours, the liquors, after removal of the first crop from benzene, being evaporated and the treatment repeated. Conversion to the phenylethylamine was carried out by a Hofmann reaction, the theoretical amount of sodium hypochlorite being used. Monomethylation of the phenylethylamine was done by the elegant Decker method,⁸ except that the reaction mixture was heated for eighteen hours at 37° in place of the higher temperatures used by Decker. The Schiff bases and the methyl iodide addition products were used directly, the first being thick sirups and the second being too unstable to manipulate. Subsequent O-demethylation was carried out in one of two ways, either by heating with hydriodic acid (52% colorless, sp. gr. 1.7) at the boiling point, removal of excess of acid and conversion to the chloride by means of silver chloride, or by demethylating the free amine by concentrated hydrochloric acid at 180°. In the latter case the amine

INTERMEDIATES, PREVIOUSLY PREPARED AND DESCRIBED

Compound	Yield, %	Lit. refs.	Compound	Yield, %	Lit. refs.
<i>o</i> -Methoxycinnamic acid ^{a,b}	95	4	2,3-Dimethoxyphenylpropionic acid	..	11
<i>o</i> -Methoxyphenylpropionic acid	87	5, 6	2,3-Dimethoxyphenylpropionamide ^{b,g}	84	11
<i>m</i> -Methoxybenzaldehyde ^{b,c,d}	83	8	2,4-Dimethoxybenzaldehyde ^d	..	12, 8
<i>m</i> -Methoxycinnamic acid	..	9	<i>Trans</i> -2,4-dimethoxycinnamic acid ^{a,b}	97	13
<i>m</i> -Methoxyphenylpropionic acid	..	10	2,4-Dimethoxyphenylpropionamide ^{b,e,f}	86	10
<i>m</i> -Methoxyphenylpropionamide ^{b,e,f}	86	10	2,4-Dimethoxyphenylpropionic acid	89	14
2,3-Dimethoxybenzaldehyde	..	8	2,5-Dimethoxybenzaldehyde	37	15
2,3-Dimethoxycinnamic acid	..	11	2,5-Dimethoxycinnamic acid ^{a,b}	95	15

⁸ Decker, *Ann.*, 395, 390 (1913).⁴ Perkin, *J. Chem. Soc.*, 31, 414 (1877).⁶ Perkin, *ibid.*, 39, 415 (1881).⁶ Tiemann and Ludwig, *Ber.*, 15, 2043 (1882).⁷ Mentioned but not described by Shoesmith and Connor, *J. Chem. Soc.*, 2230 (1927).⁸ Douetteau, *Bull. soc. chim.*, [iv] 9, 932 (1911).⁹ Chakravarti, Haworth and Perkin, *J. Chem. Soc.*, 2265 (1927).¹⁰ Helfer, *Helv. Chim. Acta*, 7, 945 (1924).¹¹ Haworth, *J. Chem. Soc.*, 2281 (1927).¹² Adams and Levine, *THIS JOURNAL*, 45, 2373 (1923).¹³ Perkin and Schiess, *J. Chem. Soc.*, 85, 159 (1904).¹⁴ Will, *Ber.*, 16, 2106 (1883).¹⁵ Kauffmann and Burr, *ibid.*, 40, 2352 (1907).

NEW INTERMEDIATES

Compound	Cryst. form	Sol. in	Yield, %	M. p., °C.	Analyses Found, %	
					C	H
<i>o</i> -Methoxyphenylpropionamide ^{7,8}	Prisms	Bz, EtOH, Et ₂ O	73	110	66.61	7.23
2,4-Dimethoxyphenylpropionamide ⁸	Prisms	Bz, EtOH, Et ₂ O	71	113	63.23	7.35
2,5-Dimethoxyphenylpropionamide ⁸	Needles	EtOH	85	111	63.24	6.99
2,5-Dimethoxyphenylpropionic acid ⁹	Poor prisms	EtOH	94	101	63.02	6.49

^a Malonic acid method of preparation. ^b Method differs from that in literature.

^c B. p. 122° (22 mm.). ^d Method of Ref. 8 adapted to present compound. ^e Ammonia method of preparation. ^f Purification difficult, b. p. 180–205° (1 mm.). ^g Sodium amalgam method of preparation.

hydrochloride was obtained directly. Repeated efforts failed to give the 2,4-dihydroxyamine, the reaction taking another course and giving products of a different type.

β -Phenylethylamines.—The amines are all colorless, somewhat viscous liquids, which rapidly absorb carbon dioxide from the air, forming solid carbonates. With the exception of the 2,3-compound, which is miscible with water, they are rather slightly soluble in water, giving strongly alkaline solutions. They are miscible with alcohol, benzene and ether. The odors are much fainter than would be expected. The mediocre yields in some cases are doubtless due to nuclear halogenation, a not infrequent accompaniment to the Hofmann reaction.

Amine	Odor	°C.	B. p., mm.	n_D^{25}	d_4^{25}	M_D	Formula	Analyses		Yield, %
								Calcd. C	Calcd. H	
2-Methoxy amine	Musty,	117	9	1.5391	1.0408	45.49 ^a	C ₉ H ₁₁ ON	71.48	8.66	65
3-Methoxy ^c amine	Musty, sweet	118	6	1.5370	1.0385	45.44 ^a	C ₉ H ₁₁ ON	71.48	8.66	33
2,3-Di-methoxy amine	Slight	138	8	1.5348	1.0839	52.01 ^b	C ₁₀ H ₁₃ O ₂ N	66.25	8.33	79
2,4-Di-methoxy amine	Faint	140	1	1.5402	1.0915	52.08 ^b	C ₁₀ H ₁₃ O ₂ N	66.25	8.33	31
2,5-Di-methoxy ester	Faint	150	8	1.5399	1.0889	52.18 ^b	C ₁₀ H ₁₃ O ₂ N	66.25	8.33	63

^a Calcd. (Brühl), 45.42. ^b Calcd. (Brühl), 51.73. ^c Helfer¹⁰ uses hypobromite.

β -Phenylethylmethylanines.—The amines were prepared by adding excess of 30% potassium hydroxide solution to the aqueous solution of the hydriodide. The resulting oil was extracted with ether, dried over solid potassium hydroxide, and distilled under reduced pressure. Small amounts of white crystalline material usually separate during the ether extraction. The amines form colorless, viscous liquids, slightly soluble in water, with the exception of the 2,3-compound, which is more soluble. The aqueous solutions are strongly alkaline. The amines do not form solid carbonates in the air.

Amine	Odor	B. p., °C.	Mm.	n_D^{25}	d_4^{25}	M_D	Formula	Analyses			
								Calcd. C	Calcd. H	Found C	Found H
2-Methoxy	Strong amine	115	9	1.5249	1.0043	50.38 ^a	$C_{10}H_{15}ON$	72.68	9.16	73.04	9.21
3-Methoxy	Strong amine	118	7	1.5225	0.9999	50.40 ^a	$C_{10}H_{15}ON$	72.68	9.16	72.83	8.99
2,3-Dimethoxy	Faint amine	136	6	1.5218	1.0471	56.82 ^b	$C_{11}H_{17}O_2N$	67.64	8.78	68.02	9.01
2,4-Dimethoxy	Faint, musty	136	4	1.5283	1.0534	57.07 ^b	$C_{11}H_{17}O_2N$	67.64	8.78	67.82	8.87
2,5-Dimethoxy	Pract. odorless	155	8	1.5278	1.0545	56.96 ^b	$C_{11}H_{17}O_2N$	67.64	8.78	68.01	8.57

^aCalcd. (Brühl), 50.23. ^bCalcd. (Brühl), 56.51.

β -Phenylethylmethylamines—Hydriodides.—With the exception of the 2,4-compound, which could not be crystallized, the hydriodides were recrystallized from alcohol-ether mixture. They are all white when pure, and are very soluble in water and in alcohol. The hydriodides show marked supercooling and are difficult to crystallize.

Hydriodide	Appearance	M. p., °C.	Formula	Analyses			
				Calcd. C	Calcd. H	Found C	Found H
2-Methoxy	Glittering rounded prisins	101	$C_{10}H_{15}ONI$	40.95	5.50	41.16	5.62
3-Methoxy	Glittering tiny rect. plates	108	$C_{10}H_{15}ONI$	40.95	5.50	41.27	5.26
2,3-Dimethoxy	Glittering prisms	90	$C_{11}H_{18}O_2NI$	40.86	5.62	41.01	5.60
2,4-Dimethoxy	Uncrystallizable sirup	...	...	...	...	...	...
2,5-Dimethoxy	Pearly plates	137	$C_{11}H_{18}O_2NI$	40.86	5.62	41.00	5.62

^aThe yields are given for unrecrystallized hydriodide. ^bWeighed as amine, 47% over both stages.

β -Phenylethylmethylamines—Hydrochlorides.—The hydrochlorides were prepared either by passing hydrogen chloride into an ether solution of the pure amine, or by the action of silver chloride on the hydriodide. They form bulky, pure white masses of micro crystals. Alcohol-ether mixture is the best solvent for recrystallization, as the hydrochlorides are all very soluble in cold water and in cold alcohol, but sparingly soluble in ether.

Hydrochloride	Appearance ^a	M. p., °C.	Formula	Analyses			
				Calcd. C	Calcd. H	Found C	Found H
2-Methoxy	Mesh of long thin plates	119	$C_{10}H_{15}ONCl$	59.52	8.00	59.31	8.05
3-Methoxy	Tiny rhombic plates	119	$C_{10}H_{15}ONCl$	59.52	8.00	59.48	8.08
2,3-Dimethoxy	Mesh of long thin plates	117	$C_{11}H_{18}O_2NCl$	56.99	7.83	57.11	7.80
2,4-Dimethoxy	Mesh of flat needle-prisms	145	$C_{11}H_{18}O_2NCl$	56.99	7.83	56.91	8.20
2,5-Dimethoxy	Mass of tiny needles	110	$C_{11}H_{18}O_2NCl$	56.99	7.83	56.80	7.60

^a Under microscope, from alcohol.

Hydroxyphenylethylmethylamines—Hydrochlorides.—The hydrochlorides all form white crystals, very soluble in water, alcohol, acetic acid and concd. hydrochloric acid. They are very difficult to obtain in a crystalline form, the 3-hydroxy compound being particularly so. When pure, however, they are quite stable. No bases could be isolated on account of their solubility and instability. The dihydroxy compounds are powerful reduc-

ing agents. Alcohol-ether or acetone-ether mixtures are the best solvents for crystallizing the salts, as they are sparingly soluble in ether.

No.	Hydrochloride	Appearance	FeCl ₃	Ammon. AgNO ₃	Excess NaOH
1	2-Hydroxy	Thin, pearly plates	Dull violet	No redn.	Colorless, stable
2	3-Hydroxy	Glittering plates	Pale violet	No redn.	Colorless, stable
3	2,3-Dihydroxy	Stars of prisms	Dull green	Reduced in cold	Rapidly blackens
4	2,5-Dihydroxy	Felted tiny plates	Transient pale green	Reduced in cold	Rapidly blackens

No.	Hydrochloride	M. p., °C.	Formula	Analyses		Found	
				C	H	C	H
1	2-Hydroxy	148	C ₉ H ₁₁ ONCl	57.63	7.53	57.98	7.49
2	3-Hydroxy	89	C ₉ H ₁₁ ONCl	57.63	7.53	57.43	7.60
3	2,3-Dihydroxy	149	C ₉ H ₁₁ O ₂ NCl	53.05	6.93	53.30	6.72
4	2,5-Dihydroxy	128	C ₉ H ₁₁ O ₂ NCl	53.05	6.93	53.23	7.01

The analyses given in this paper are all micro-analyses (Pregl). The author is indebted to Mr. W. S. Ide for these.

Summary

The paper describes the synthesis of four of the seven previously unknown hydroxy- and dihydroxyphenylethylmethylamines. The synthesis of one of the remaining three broke down at the last stage. It has not been possible to obtain starting material to attempt the syntheses of the other two.

TUCKAHOE, NEW YORK

[CONTRIBUTION FROM THE DEPARTMENT OF CHEMISTRY OF THE UNIVERSITY OF ILLINOIS]

THE HALOFORM REACTION. VI. ALPHA-HALOGEN DERIVATIVES OF HINDERED KETONES

BY C. HAROLD FISHER, HAROLD R. SNYDER AND REYNOLD C. FUSON

RECEIVED APRIL 15, 1932

PUBLISHED SEPTEMBER 5, 1932

The study of the halogenation phase of the haloform reaction has now been extended to types other than methyl ketones, and it has been found that the replacement of α -hydrogen by halogen is a fairly general reaction between hindered ketones and hypohalite solutions. This investigation has also included a study of certain unusual reactions which are apparently general for this type of compound. These reactions are (1) cleavage by alkali, (2) the dehalogenation by phenol and (3) dehalogenation by the Grignard reagent. These several phases of the work will be discussed under separate headings.

The Halogenation.—From the previous work¹ in this series it appears that the hindered methyl ketones are generally converted directly into the corresponding trihalomethyl compounds. In no case has the mono- or the di-halomethyl derivative been isolated as an intermediate.

¹ (a) Fuson and Walker, *THIS JOURNAL*, **52**, 3269 (1930); (b) Gray, Walker and Fuson, *ibid.*, **53**, 3494 (1931); (c) Fuson, Farlow and Stehman, *ibid.*, **53**, 4097 (1931); (d) Fuson, Lewis and Du Puis, *ibid.*, **54**, 1114 (1932).