

Molecular phylogeny of *Psilocybe cyanescens* complex in Europe, with reference to the position of the secotioid *Weraroa novae-zelandiae*

Jan Borovička · Machiel E. Noordeloos ·
Milan Gryndler · Miroslav Oborník

Received: 10 March 2010 / Revised: 10 May 2010 / Accepted: 18 May 2010 / Published online: 8 June 2010

Abstract A phylogenetic analysis with three molecular markers was undertaken to test the hypothesis that the complex of *Psilocybe cyanescens* in Europe consists of several, morphologically distinct species. The results support the existence of two molecularly well-supported morphological groups, that of *Psilocybe cyanescens* and *P. azurescens* on the one hand, and the complex of *P. serbica* on the other. However, in the last group, no sequence

variability within the three molecular markers from *P. serbica* and related taxa *P. bohémica*, *P. arcana*, and *P. moravica* was found. It was decided, therefore, to merge these taxa into *P. serbica*, and to distinguish them below species level. It was also demonstrated that the secotioid *Weraroa novae-zelandiae* belongs to the *P. cyanescens* species complex. Accordingly, it was transferred to *Psilocybe* as *P. weraroa*, nomen novum.

Electronic supplementary material The online version of this article (doi:10.1007/s11557-010-0684-3) contains supplementary material, which is available to authorized users.

J. Borovička
Institute of Geology, Academy of Sciences of the Czech Republic,
Rozvojová 269,
165 00 Prague 6, Czech Republic

J. Borovička
Nuclear Physics Institute, Academy of Sciences of the Czech Republic,
250 68 Řež,
Prague, Czech Republic

M. E. Noordeloos
Netherlands Centre for Biodiversity Naturalis (section NHN),
Leiden University,
P.O. Box 9514, 2300 RA Leiden, The Netherlands

M. Gryndler
Institute of Microbiology, Academy of Sciences of the Czech Republic,
Videňská 1083,
142 20 Prague 4, Czech Republic

M. Oborník (✉)
Biology Centre, Institute of Parasitology,
Academy of Sciences of the Czech Republic,
Branišovská 31,
37005 České Budějovice, Czech Republic
e-mail: obornik@paru.cas.cz

Keywords Strophariaceae · *Psilocybe serbica* · *Psilocybe cyanescens* · *Weraroa novae-zelandiae* · DNA markers · Phylogeny

Introduction

The discovery of the Mexican cult of “magic mushrooms” in the 1950s has started an intensive investigation of taxonomy, ecology and biochemistry of psychoactive fungi throughout the world (Heim and Wasson 1958). The major part of the known psychoactive species belongs to the genus *Psilocybe* (Basidiomycota, Agaricales, Strophariaceae) (for reviews, see Courtecuisse and Deveau 2004; Anderson et al. 2008; Guzmán 2009). *Psilocybe* species grow throughout most of the world and more than 200 taxa have been described to date (Guzmán 1983, 1995); more than 25 species are reported from Europe (Horak 2005; Noordeloos 1999; Knudsen and Vesterholt 2008).

Based on their ecology, the European psychoactive species, which are sometimes used as recreational drugs (Hillebrand et al. 2006), are divided into two main groups: (1) species growing in pastures and meadows, often on dung, and (2) species growing on plant debris in forests, parks and gardens. The latter is known as the *Psilocybe cyanescens* complex. During the 1980s, the European

Psilocybe species of the *P. cyanescens* complex were studied by the German mycologist, G.J. Krieglsteiner, who proposed a rather wide species concept of *Psilocybe cyanescens* Wakef., including as synonyms *P. serbica* M.M. Moser & E. Horak, *P. mairei* Singer and *P. bohémica* Šebek ex Šebek (Krieglsteiner 1984, 1986); this concept has been widely accepted in Europe (Ludwig 2001; Bon and Roux 2003; Horak 2005; Knudsen and Vesterholt 2008). Guzmán (1983, 1995), however, recognizes *P. serbica* and *P. cyanescens* as distinct species, but makes no reference to the works of Krieglsteiner.

However, as a result of a thorough morphological and ecological study of this species complex in the Czech Republic (Borovička and Hlaváček 2001a; b; Borovička 2003, 2005, 2006, 2008), seven taxa have been distinguished. These are divided over two stirpes (groups): (1) stirps *Cyanescens*, including *P. cyanescens* Wakef. (Dennis and Wakefield 1946) and *P. azurescens* Stamets and Gartz (1995), and (2) stirps *Serbica* including *P. serbica* Moser and Horak (1968), *P. bohémica* Šebek ex Šebek (1983), *P. arcana* Borovička and Hlaváček (2001a), *P. moravica* Borovička (2003), *P. moravica* var. *sternberkiana* Borovička (2006), and possibly *P. mairei* Singer (1973).

The aim of this study was to explore the relationships within the proposed *P. cyanescens* species complex by use of molecular markers and to compare that to the analysis of morphological, phenological and ecological data. Furthermore, the secotioid genus *Weraroa* was shown to be polyphyletic, as its species were shown to be distributed in different subclades of the Stropharioid clade (Bridge et al. 2008; Moncalvo et al. 2002). The type species, *Weraroa novae-zelandiae*, appeared in the clade “Psychodelia” close to *P. cyanescens*. For that reason, this species has been included in our study.

Materials and methods

Collections used

This study includes material of all taxa of the stirpes *Cyanescens* and *Serbica* except for *Psilocybe mairei* Singer. This last species was described from North Africa as *Hypholoma cyanescens* Maire (1928), but no recent collections were available. All of the collections used for DNA extraction fit well with the morphological and ecological concept by Borovička (2008) (Table 1). When possible, holotype collections were used for sequencing (*P. arcana*, *P. moravica*, *P. moravica* var. *sternberkiana*) or the collections were sampled in original localities (*P. bohémica*, locality of epitype; *P. arcana*, topotype). We were not successful in obtaining DNA data from the paratype of *P. serbica* (herb. E. Horak 63/2536), so we used a collection

donated from Croatia whose microcharacters (spore size, cheilocystidia shape and pleurocystidia occurrence) agree well to those observed in the original collection. Available sequences of related species were downloaded from GeneBank™ (Electronic supplementary material, ESM, Table S1). The secotioid fungus *Weraroa novae-zelandiae* was also included in this study because earlier reports show that this species might well be very closely related to members of the *P. cyanescens* complex (Bridge et al. 2008).

Molecular phylogeny

Nuclear DNA was extracted from a small piece of dried or frozen fungal biomass (a piece of basidiocarp) using the MoBio Power Soil DNA isolation kit (Mo-Bio Laboratories, Solana Beach, CA, USA) according to the manufacturer's instructions. We have used three molecular markers in particular: partial sequences of the nuclear LSU rRNA gene (Moncalvo et al. 2000, 2002; Maruyama et al. 2003, 2006; Nugent and Saville 2004; Garnica et al. 2007) and ITS1+5.8S+ITS2 region (Guzmán-Dávalos et al. 2003; Watling and Martin 2003; Nugent and Saville 2004; Matheny et al. 2006; Pornpakakul et al. 2009), which have previously been used to infer phylogenetic relationship of the genus *Psilocybe*; in addition to those, we have sequenced part of the elongation factor 1- α (EF1 α) which has not previously been used for phylogenetic studies in *Psilocybe*, but appeared useful in other groups of fungi (Kausserud and Schumacher 2001; Antonín et al. 2009).

Primer pair NL-1 and NL-4 was used to amplify the part of 28 S rDNA. To amplify the complete sequence of ITS1 +5.8S+ITS2 region and a part of 18S rDNA, the forward, fungal-specific ITS-1F primer and universal reverse primer ITS-4 or basidiomycete-specific reverse primer ITS-4B were used as recommended (White et al. 1990; Gardes and Bruns 1993). The partial elongation factor EF1 α sequence was amplified using the forward primer EF-595F and reverse primer EF-1160R. The EF1 α sequence included two partial and one complete exon and two introns (Kausserud and Schumacher 2001).

The reaction to amplify both rRNA genes was performed in 50 μ l volume with 1 μ l undiluted DNA preparation (approx. 5 ng μ l⁻¹) as a template, PPP Master Mix (*Taq*-Purple DNA polymerase PCR Master) supplying reaction with 2.5 U of *Taq* DNA polymerase, deoxyribonucleotides (200 μ M each) and 2.5 mM MgCl₂, and 200 nM of each primer. The reaction was, following the initial denaturation (5 min at 94°C), subjected to 34 cycles of 1 min at 94°C, 52°C for 1 min and 2 min at 72°C. To complete the amplicons, the reaction was further incubated for 10 min at 72°C. The elongation factor EF1 α was amplified in total volume of 50 μ l and amplification program consisted of pre-denaturation for 4 min at 95°C, which was followed by

Table 1 *Psilocybe* species under study

ID	Species	Collection	Origin	GeneBank™		
				ITS	LSU	EF1-a
Stirps <i>Cyanescens</i> (Borovička 2005)						
P 02	<i>Psilocybe cyanescens</i> Wakef.	PRM 902040	Belgium	GU565176	-	GU565161
P 21	<i>Psilocybe cyanescens</i> Wakef.	PRM 901481	Germany	GU565175	GU565167	GU565158
P 01*	<i>Psilocybe azurescens</i> Stamets & Gartz	PRM 901020	Oregon, USA	GU565173	GU565166	GU565159
Stirps <i>Serbica</i> (Borovička 2005)						
P 15	<i>Psilocybe serbica</i> M.M. Moser E. Horak	PRM 903176	Croatia	GU565177	GU565168	GU565165
P 14	<i>Psilocybe bohémica</i> Šebek ex Šebek	L 0342813	Czech Rep.	GU565178	GU565169	GU565162
P 05	<i>Psilocybe arcana</i> Borov. & Hlaváček	PRM 895093, holotype	Czech Rep.	GU565180	-	-
P 05*	<i>Psilocybe arcana</i> Borov. & Hlaváček	PRM 915262, topotype	Czech Rep.	GU565181	GU565172	GU565160
P 06	<i>Psilocybe moravica</i> Borov.	PRM 900455, holotype	Czech Rep.	GU565182	GU565171	GU565164
P 07	<i>P. moravica</i> var. <i>sternberkiana</i> Borov.	PRM 901650, holotype	Czech Rep.	GU565179	GU565170	GU565163

PRM Herbarium of the Mycological Department, National Museum, Prague; L Nationaal Herbarium Nederland, Leiden University branch

37 cycles containing temperature profile 94°C for 30 s, 55°C for 35 s and 72°C for 40 s. The program was terminated by final extension at 72°C for 10 min. Obtained amplicons were purified using the MoBio UltraClean PCR Clean-up Kit and both strands were directly sequenced with ABI Prism 3130XL sequence analyzer.

The sequences coding for rRNA genes were aligned to their homologues downloaded from GeneBank™ using Kalign program (Lassmann and Sonnhammer 2005) (available at <http://www.ebi.ac.uk/Tools/kalign/index.html>). Gaps and ambiguously aligned regions were excluded from further analysis. Nucleotide sequences coding for homologues of elongation factor EF1 α were downloaded from GeneBank™ and they were in the form of deduced amino acid sequences aligned to the genes from *Psilocybe* spp. using ClustalW algorithm. The alignment was retranslated back to nucleotides (BioEdit; see Hall 1999). Both introns were identified and their phylogenetic signal was explored separately from the mature protein coding sequence. Both nucleotide and aminoacid alignments were subsequently used to construct phylogenetic trees by maximum parsimony (MP) and maximum likelihood (ML) methods. Models for ML were selected using Modeltest program (GTR+ Γ +I) (Posada and Crandall 1998). All parameters such as gamma shape parameter and proportion of invariant sites were estimated from particular datasets. Robustness of constructed trees was tested by bootstrap analysis in 1,000 and 300 replicates for MP and ML, respectively.

To compare molecular analysis with that using phenotypic data, we have analyzed 34 morphological, ecological and phenological characters and computed a dendrogram based on them. We have used both approaches to test the validity of the proposed morphospecies with a phylogenetic analysis using molecular markers and to elucidate the

evolution of this group. The morphological matrix was constructed from 34 morphological, ecological and phenological data (ESM, Table S2) including, e.g., habitus, fruit-body size, shape of pileus, coloration of pileus when moist and in drying, occurrence of olive-green tinges, shape of stipe, character and occurrence of velar remnants, smell, size and shape of spores, shape and occurrence of pleuro- and cheilocystidia, distribution in Europe, ecology (preferred substrate, habitat), and fruiting maximum in Central Europe. The tree based on this matrix was computed using maximum parsimony and minimum evolution as implemented in PAUP*4b10 (Swofford 1998). Robustness of the tree was tested by bootstrap analysis from 1,000 replicates for both methods.

Results and discussion

A computational matrix reflecting phenotypic characters (morphology, ecological and phenological features, 34 countable characters) was established for *P. cyanescens*, *P. azurescens*, *P. serbica*, *P. arcana*, *P. bohémica*, *P. moravica*, and *P. moravica* var. *sternberkiana*, and was used to infer phylogeny of species of interest. Distant *P. cubensis* was used as outgroup (ESM, Table S2). Analysis of phenotypic characters gave a tree showing two distinct stirpes *Cyanescens* and *Serbica*. Within the stirps *Serbica*, *P. moravica* appeared as a sister to *P. moravica* var. *sternberkiana* with *P. bohémica* as the most basal species. The remaining species of this stirps, *P. serbica* and *P. arcana*, form a separated but related cluster (Fig. 1a).

Molecular phylogeny shows two distinct groups within the species of the *P. cyanescens* complex in Europe (Fig. 1b–e): group 1 encompassing *P. cyanescens* and *P.*




Fig. 1 Phylogenetic trees as inferred from various molecular and morphological data. **a** Minimum evolution tree based on morphological, phenological and ecological data (bootstraps minimum evolution/maximum parsimony; both 1,000 replicates). **b** Maximum likelihood tree (loglik = -2707.37145; gamma shape 0.015; proportion of invariant sites 0.000) based on partial LSU rDNA sequences. **c** Topology of species of interest as inferred by maximum parsimony method. **d**

Maximum likelihood tree (loglik = -1938.37061; gamma shape 0.496; proportion of invariant sites 0.386) as inferred from ITS1+5.8 S+ITS2 sequences. **e** Maximum likelihood tree (loglik = -1693.32043; gamma shape 1.026; proportion of invariant sites 0.509) as inferred from nucleotide sequences coding for elongation factor 1- α (EF1 α). Numbers above branches indicate ML/MP bootstraps (number of replicates 300/1,000)

azurescens, and group 2 encompassing *P. serbica*, *P. bohemica*, *P. arcana* and *P. moravica* (including its var. *sternberkiana*). The latter group is very well supported by bootstrap values. Furthermore, the taxa of the second group do not differ among each other in the sequence of all three DNA markers. We have also analyzed concatenated introns from EF1 α and have obtained the same topology as from the coding sequence (not shown). The absence of DNA polymorphism among the taxa of the *P. serbica* complex in

studied markers indicates that these taxa represent, in fact, a single species despite their phenotypic variability.

In general, molecular markers can successfully uncover the existence of cryptic species. Such species do not display morphological variation but can be well distinguished thanks to the polymorphism of DNA markers (Geiser et al. 1998; Roy et al. 1998; Cruse et al. 2002). In our case, however, the opposite situation seems to be at hand: a group of morphologically distinguishable fungal taxa that

display no sequence variation in three widely used DNA markers. In this particular case, the Morphological Species Concepts (MSC) and Ecological Species Concept (ESC) are in conflict with the Phylogenetic Species Concept (Giraud et al. 2008).

When only morphological and ecological characters are used for classification, *P. serbica* complex is very variable and this variability has led to description of several species within it in the past. However, when the DNA sequence characters are used, the previously described taxa within the stirps *Serbica* appear to be identical. This is surprising particularly in the case of ITS, which represents a highly polymorphic marker and consequently a powerful tool for taxonomic purposes at species level. It was suggested, based on analysis of 1,373 ITS2 sequences, that in 93% of cases where two taxa differed in any property of the ITS2 (sequential or secondary structural) they represented different species (Müller et al. 2007). Furthermore, this is contrasting with, for example, the results of Nugent and Saville (2004) and Maruyama et al. (2003) who successfully used partial sequences of LSU rRNA gene to infer phylogeny of closely related hallucinogenic species of the genus *Psilocybe*. Another example can be found in the *Armillaria* study of Antonín et al. (2009).

We therefore conclude that the *P. cyanescens* complex in Europe consists of three well-defined species, *P. cyanescens*, *P. azurescens* and *P. serbica*. The last species is, furthermore, very variable in macro- and microcharacters. The most distinct morphological entities, so far described at species level, namely *P. arcana*, *P. bohemica* and *P. moravica*, need to get a formal rank below species level. Consequently, we propose to treat *P. moravica* var. *sternberkiana* on the level of forma.

We have to note that our analysis led to the taxonomic suggestions also concerning other *Psilocybe* species. A secotioid fungus traditionally classified as *Weraroa novae-zelandiae* (G. Cunn.) Singer is obviously closely related to the investigated bluing species of the *Psilocybe cyanescens* complex (as already suggested by Bridge et al. 2008) and belongs to the genus *Psilocybe*. A similar phenomenon has also been observed in the case of other secotioid fungi, e.g., *Gyrophragmium dunalii* (Fr.) Zeller and *Endoptychum agaricoides* Czern., which are currently classified as *Agaricus aridicola* Geml, Geiser & Royse (Geml et al. 2004) and *Chlorophyllum agaricoides* (Czern.) Vellinga (Vellinga 2002), respectively.

Conclusion

Our analysis does not support the widely accepted view that there are only two wood-rotting bluing *Psilocybe* species

(*P. cyanescens* and *P. azurescens*) in Europe. Molecular analysis recognizes two sibling groups within the *P. cyanescens* complex and three distinct species: *P. cyanescens*, *P. azurescens* and *P. serbica*. The recently described morphospecies *P. bohemica*, *P. arcana*, and *P. moravica* apparently represent just morphological and ecological varieties of *P. serbica*, and should accordingly be treated below species level. The species of the two groups can be unambiguously recognized microscopically by occurrence and shape of pleurocystidia (Borovička 2008). *Psilocybe serbica* now appears as a highly variable species characterized, e.g., by various robustness, shape and coloration of fruit-bodies, and microscopically by a large range of spore size and varying abundance/shape of pleuro- and cheilocystidia. The phylogeny also shows that the secotioid *Weraroa novae-zelandiae* fits well in the agaricoid *P. cyanescens* group.

Taxonomic changes

Psilocybe serbica* var. *bohemica (Šebek ex Šebek) Borov., Oborník & Noordel., comb. & stat. nov.

Mycobank: MB 515725

Basionym: *Psilocybe bohemica* Šebek ex Šebek, Česká Mykol. 37(3): 177 (1983).

Psilocybe serbica* var. *arcana (Borov. & Hlaváček) Borov., Oborník & Noordel., comb. & stat. nov.

Mycobank: MB 515724

Basionym: *Psilocybe arcana* Borov. & Hlaváček, Mykol. Sborn. 78(1): 3 (2001a).

Psilocybe serbica* var. *moravica (Borov.) Borov., Oborník & Noordel., comb. & stat. nov.

Mycobank: MB 515726

Basionym: *Psilocybe moravica* Borov., Mykol. Sborn. 80(4): 127 (2003).

Psilocybe serbica* f. *sternberkiana (Borov.) Borov., Oborník & Noordel., comb. & stat. nov.

Mycobank: MB 518345

Basionym: *Psilocybe moravica* var. *sternberkiana* Borov., Czech Mycol. 58(1-2): 76 (2006).

Psilocybe weraroa Borov., Oborník & Noordel., nom. nov.

Mycobank: MB 515728

Replaced synonym: *Secotium novae-zelandiae* G. Cunn., Proc. Linn. Soc. N.S.W. 49(2): 107 (1924).

Etymology This species has been known as *Weraroa novae-zelandiae* (G. Cunn.) Singer; the epithet of the replacement name (nomen novum) refers to the former generic name. It was necessary to create the replacement name because the binomial *Psilocybe novae-zelandiae* was preoccupied by *P. novae-zelandiae* Guzmán & E. Horak, Sydowia 31(1-6): 51 (1979) [1978]; see also Redhead et al. (2007).

Acknowledgements We are very grateful to Michal Tomšovský for valuable advice and to our colleagues who kindly provided their collections for study: anonymous donors, Jürgen Hechler, Egon Horak, Zdenko Tkálčec, and Ruben Walley (†). The kind assistance of curators Lenka Edrová and Jan Holec (herbarium PRM; Mycological Department, National Museum, Prague) is also greatly appreciated. This research was supported by Institutional Research Plans (IRP) AV0Z30130516 (Institute of Geology, ASCR, Prague), AV0Z10480505 (Nuclear Physics Institute, ASCR, Řež near Prague), IRP AV0Z50200510 (Institute of Microbiology, ASCR, Prague), research project of the Institute of Parasitology ASCR (Z60220518) and Ministry of Education of the Czech Republic (6007665801). An indirect support was obtained from the project IAA600480801 (The Grant Agency of the Academy of Sciences of the Czech Republic).

References

- Anderson C, Kristinsson J, Gry J (2008) Occurrence and use of hallucinogenic mushrooms containing psilocybin alkaloids. *TemaNord* 2008:606, Nordic Council of Ministers, Copenhagen
- Antonín V, Tomšovský M, Sedláč P, Májek T, Jankovský L (2009) Morphological and molecular characterization of the *Armillaria cepistipes* – *A. gallica* complex in the Czech Republic and Slovakia. *Mycol Prog* 8:259–271
- Bon M, Roux P (2003) Clé analytique de la famille *Strophariaceae* Singer & A.H. Smith. *Doc Mycol* 129:3–54
- Borovička J (2003) The bluing *Psilocybe* species of the Czech Republic III. *Psilocybe moravica* sp. nova, the Moravian *Psilocybe*. *Mykol Sborn* 80:126–141
- Borovička J (2005) The bluing species of *Psilocybe* of the Czech Republic IV. The problem of *Psilocybe cyanescens* Wakef. *Mykol Sborn* 82:1–21
- Borovička J (2006) New variety of *Psilocybe moravica* and notes on *Psilocybe bohemia*. *Czech Mycol* 58:75–80
- Borovička J (2008) The wood-rotting bluing *Psilocybe* species in Central Europe – an identification key. *Czech Mycol* 60:173–192
- Borovička J, Hlaváček J (2001a) The bluing *Psilocybe* species of the Czech Republic I. *Psilocybe arcana* Borovička et Hlaváček, the mysterious *Psilocybe* [In Czech]. *Mykol Sborn* 78:2–7
- Borovička J, Hlaváček J (2001b) The bluing *Psilocybe* species of the Czech Republic II. *Psilocybe bohemia* Šebek, the Bohemian *Psilocybe* [In Czech]. *Mykol Sborn* 78:57–65
- Bridge PD, Spooner BM, Beever RE, Park DC (2008) Taxonomy of the fungus commonly known as *Stropharia aurantiacea*, with new combinations in *Leratiomyces*. *Mycotaxon* 103:109–121
- Courtecuisse R, Deveau M (2004) Mycology and toxicology of hallucinogenic mushrooms in Europe and the Americas [In French]. *Ann Toxicol Anal* 16:36–64
- Cruse M, Telerant R, Gallagher T, Taylor JW (2002) Cryptic species in *Stachybotrys chartarum*. *Mycologia* 94:814–822
- Dennis RWG, Wakefield EM (1946) New or interesting British fungi. *Trans Br Mycol Soc* 29:141–142
- Gardes M, Bruns TD (1993) ITS primers with enhanced specificity for basidiomycetes – application to the identification of mycorrhizae and rusts. *Mol Ecol* 2:113–118
- Garnica S, Weiss M, Walther G, Oberwinkler F (2007) Reconstructing the evolution of agarics from nuclear gene sequences and basidiospore ultrastructure. *Mycol Res* 111:1019–1029
- Geiser DM, Pitt IJ, Taylor JW (1998) Cryptic speciation and recombination in the aflatoxin-producing fungus *Aspergillus flavus*. *Proc Natl Acad Sci USA* 95:388–393
- Geml J, Geiser DM, Royse DJ (2004) Molecular evolution of *Agaricus* species based on ITS and LSU rDNA sequences. *Mycol Prog* 3:157–176
- Giraud T, Refrégier G, Le Gac M, de Vienne DM, Hood ME (2008) Speciation in fungi. *Fungal Genet Biol* 45:791–802
- Guzmán G (1983) The genus *Psilocybe*. *Nova Hedwigia, Beiheft* 74, Vaduz
- Guzmán G (1995) Supplement to the monograph of the genus *Psilocybe*. In: Petrini O, Horak E (eds) Taxonomic monographs of Agaricales. *Bibliotheca Mycol* 159:91–141
- Guzmán G (2009) The hallucinogenic mushrooms: diversity, traditions, use and abuse with special reference to the genus *Psilocybe*. In: Misra JK, Desmukh SK (eds) Fungi from difference environments. Science Publishers, Enfield (NH), pp 256–277
- Guzmán-Dávalos L, Mueller GM, Cifuentes J, Miller AN, Santerre A (2003) Traditional infrageneric classification of *Gymnopilus* is not supported by ribosomal DNA sequence data. *Mycologia* 95:1204–1214
- Hall TA (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp Ser* 41:95–98
- Heim R, Wasson RG (1958) Hallucinogenic mushrooms of Mexico [In French]. *Archives Du Muséum National d'Histoire Naturelle, 7e Série, Tome VI, Éditions du Muséum, Paris*
- Hillebrand J, Olszewski D, Sedefov R (2006) EMCDDA Thematic Papers, Hallucinogenic mushrooms: an emerging trend case study. European Monitoring Centre for Drugs and Drug Addiction, Lisbon
- Horak E (2005) European Agarics and Boleti [In German]. Elsevier, München
- Kausserud H, Schumacher T (2001) Outcrossing or inbreeding: DNA markers provide evidence for type of reproductive mode in *Phellinus nigrolimitatus* (Basidiomycota). *Mycol Res* 105:676–683
- Knudsen H, Vesterholt J (eds) (2008) *Funga Nordica: Agaricoid, boletoid and cyphelloid genera*. Copenhagen
- Kriegelsteiner GJ (1984) Studien zum *Psilocybe cyanescens*-Komplex in Europa. *Beitr Kenntn Pilze Mitteleur* 1:61–94
- Kriegelsteiner GJ (1986) Studien zum *Psilocybe cyanescens-callosa-semilanceata*-Komplex in Europa. *Beitr Kenntn Pilze Mitteleur* 2:57–72
- Lassmann T, Sonnhammer EL (2005) Kalign – an accurate and fast multiple sequence alignment algorithm. *BMC Bioinf* 6:298
- Ludwig E (2001) Pilzkompendium, Part 1, Descriptions [In German]. Eching
- Maire R (1928) Diagnoses de champignons inédits de l'Afrique du Nord. *Bull Soc Mycol Fr* 44:37–56
- Maruyama T, Yokoyama K, Makino Y, Goda Y (2003) Phylogenetic relationship of psychoactive fungi based on the rRNA gene for a large subunit and their identification using the TaqMan assay. *Chem Pharm Bull* 51:710–714
- Maruyama T, Kawahara N, Yokoyama K, Makino Y, Fukiharuru T, Goda Y (2006) Phylogenetic relationship of psychoactive fungi based on rRNA gene for a large subunit and their identification using the TaqMan assay (II). *Forensic Sci Int* 163:51–58
- Matheny PB, Curtis JM, Hofstetter V, Aime MC, Moncalvo JM, Ge ZW, Slot JC, Ammirati JF, Baroni TJ, Bougher NL, Hughes KW, Lodge DJ, Kerrigan RW, Seidl MT, Aanen DK, DeNitis M, Daniele GM, Desjardin DE, Kropp BR, Norvell LL, Parker A, Vellinga EC, Vilgalys R, Hibbett DS (2006) Major clades of Agaricales: a multilocus phylogenetic overview. *Mycologia* 98:982–995
- Moncalvo JM, Lutzoni FM, Rehner SA, Johnson J, Vilgalys R (2000) Phylogenetic relationships of agaric fungi based on nuclear large subunit ribosomal DNA sequences. *Syst Biol* 49:278–305

- Moncalvo JM, Vilgalys R, Redhead SA, Johnson JE, James TY, Aime MC, Hofstetter V, Verduin SJW, Larsson E, Baroni TJ, Thom RG, Jacobsson S, Clémence H, Miller OK Jr (2002) One hundred and seventeen clades of euagarics. *Mol Phylogenet Evol* 23:357–400
- Moser M, Horak E (1968) *Psilocybe serbica* spec. nov., eine neue Psilocybin und Psilocin Bildende Art aus Serbien. *Z Pilzk* 34:37–144
- Müller T, Philippi N, Dandekar T, Schultz J, Wolf M (2007) Distinguishing species. *RNA* 13:1469–1472
- Noordeloos ME (1999) Genus *Psilocybe*. In: Bas C, Kuyper TW, Noordeloos ME, Vellinga EC (eds) *Flora agaricina neerlandica*, vol IV. Brookfield, Rotterdam, pp 28–79
- Nugent KG, Saville BJ (2004) Forensic analysis of hallucinogenic fungi: a DNA-based approach. *Forensic Sci Int* 140:147–157
- Oborník M, Jirků M, Doležal D (2001) Phylogeny of mitosporic entomopathogenic fungi: Is the genus *Paecilomyces* polyphyletic? *Can J Microbiol* 47:813–819
- Pornpakakul S, Suwanchaoen S, Petsom A, Roengsumran S, Muangsins N, Chaichit N, Piapukiew J, Sihanonth P, Allen JW (2009) A new sesquiterpenoid metabolite from *Psilocybe samuiensis*. *J Asian Nat Prod Res* 11:12–17
- Posada D, Crandall KA (1998) Modeltest: testing the model of DNA substitution. *Bioinformatics* 14:817–818
- Redhead SA, Moncalvo JM, Vilgalys R, Matheny PB, Guzmán-Dávalos L, Guzmán G (2007) Proposal to conserve the name *Psilocybe* (Basidiomycota) with a conserved type. *Taxon* 56:255–257
- Roy BA, Vogler DR, Bruns TD, Szaro TM (1998) Cryptic species in the *Puccinia monoica* complex. *Mycologia* 90:846–853
- Šebek S (1983) The Bohemian *Psilocybe* – *Psilocybe bohemica* [In Czech]. *Czech Mycol* 37:177–181
- Singer R (1973) Diagnoses Fungorum Novorum Agaricalium III. *Beih Sydowia* 7:1–106
- Stamets P, Gartz J (1995) A new caeruleuscent *Psilocybe* from the Pacific Coast of Northwestern America. *Integration* 6:21–27
- Swofford DL (1998) *Phylogenetic Analysis Using Parsimony* (and other methods). Version 4b10. Sinauer, Sunderland, Mass
- Vellinga EC (2002) New combinations in *Chlorophyllum*. *Mycotaxon* 83:415–417
- Watling R, Martin MP (2003) A sequestrate *Psilocybe* from Scotland. *Bot J Scotl* 55:245–257
- White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ (eds) *PCR Protocols, a guide to methods and applications*. Academic, San Diego, CA, pp 315–322