

Studies on the role of dopamine in the degeneration of 5-HT nerve endings in the brain of Dark Agouti rats following 3,4-methylenedioxymethamphetamine (MDMA or 'ecstasy') administration

*¹M.I. Colado, ¹E. O'Shea, ¹R. Granados, ¹B. Esteban, ¹A.B. Martín & ²A.R. Green

¹Departamento de Farmacología, Facultad de Medicina, Universidad Complutense, Madrid 28040, Spain; ²School of Pharmacy and Pharmaceutical Sciences, De Montfort University, Leicester LE1 9RH, England, U.K.

1 We investigated whether dopamine plays a role in the neurodegeneration of 5-hydroxytryptamine (5-HT) nerve endings occurring in Dark Agouti rat brain after 3,4-methylenedioxymethamphetamine (MDMA or 'ecstasy') administration.

2 Haloperidol (2 mg kg⁻¹ i.p.) injected 5 min prior and 55 min post MDMA (15 mg kg⁻¹ i.p.) abolished the acute MDMA-induced hyperthermia and attenuated the neurotoxic loss of 5-HT 7 days later. When the rectal temperature of MDMA + haloperidol treated rats was kept elevated, this protective effect was marginal.

3 MDMA (15 mg kg⁻¹) increased the dopamine concentration in the dialysate from a striatal microdialysis probe by 800%. L-DOPA (25 mg kg⁻¹ i.p., plus benserazide, 6.25 mg kg⁻¹ i.p.) injected 2 h after MDMA (15 mg kg⁻¹) enhanced the increase in dopamine in the dialysate, but subsequent neurodegeneration was unaltered. L-DOPA (25 mg kg⁻¹) injected before a sub-toxic dose of MDMA (5 mg kg⁻¹) failed to induce neurodegeneration.

4 The MDMA-induced increase in free radical formation in the hippocampus (indicated by increased 2,3- and 2,5-dihydroxybenzoic acid in a microdialysis probe perfused with salicylic acid) was unaltered by L-DOPA.

5 The neuroprotective drug clomethiazole (50 mg kg⁻¹ i.p.) did not influence the MDMA-induced increase in extracellular dopamine.

6 These data suggest that previous observations on the protective effect of haloperidol and potentiating effect of L-DOPA on MDMA-induced neurodegeneration may have resulted from effects on MDMA-induced hyperthermia.

7 The increased extracellular dopamine concentration following MDMA may result from effects of MDMA on dopamine re-uptake, monoamine oxidase and 5-HT release rather than an 'amphetamine-like' action on dopamine release, thus explaining why the drug does not induce degeneration of dopamine nerve endings.

Keywords: 3,4-methylenedioxymethamphetamine; ecstasy; dopamine; free radicals; neurodegeneration; clomethiazole; 5-hydroxytryptamine; L-DOPA, haloperidol

Abbreviations: ANOVA, analysis of variance; CMZ, chlomethiazole; DA, dopamine; DHBA, dihydroxybenzoic acid; DOPAC, dihydroxyphenylacetic acid; Hal, haloperidol; 5-HIAA, 5-hydroxyindoleacetic acid; h.p.l.c., high performance liquid chromatography; 5-HT, 5-hydroxytryptamine; HVA, homovanillic acid; MAO, monoamine oxidase; MDMA, (±) 3,4-methylenedioxymethamphetamine; 6-OHDA, 6-hydroxydopamine; PCA, p-chloroamphetamine

Introduction

The commonly misused recreational drug 3,4-methylenedioxymethamphetamine (MDMA or 'ecstasy') has two well established phases of action in experimental animals. The acute phase consists of a rapid release of 5-hydroxytryptamine (5-HT) from neuronal stores, thereby producing many of the acute behavioural changes, including behavioural excitation and hyperthermia, that are seen in both experimental animals and humans (see reviews of White *et al.*, 1996; Green *et al.*, 1995; Steele *et al.*, 1994).

The second action of MDMA in the brain of experimental animals is to produce a long lasting neurotoxic loss of 5-HT nerve terminals in several areas of the brain, an effect demonstrated both histologically (O'Hearn *et al.*, 1988; Molliver *et al.*, 1990) and biochemically (Battaglia *et al.*,

1987; Sharkey *et al.*, 1991; Hewitt & Green, 1994). The mechanisms involved in producing this neurodegeneration are not fully understood at present, but recent data have given substantial evidence to support the contention that increased free radical production is involved (see Colado *et al.*, 1997b).

Studies on both the pharmacology of MDMA and putative neuroprotective agents have also indicated a role for dopamine in the neurodegenerative process. For example Stone *et al.*, (1988) proposed a role for dopamine based on their studies which showed that the long term depletion of cerebral 5-HT which follows administration of MDMA was attenuated by the dopamine synthesis inhibitor α -methyl p-tyrosine or the monoamine depletor reserpine. Schmidt *et al.* (1990b) reported that injection into the striatum of the selective dopamine neurotoxin 6-hydroxydopamine (6-OHDA) blocked the neurotoxic effects of MDMA on 5-HT neurones not only in the striatum but also other forebrain regions such as the

*Author for correspondence.

hippocampus and cortex. In addition Stone *et al.* (1988) reported that the dopamine uptake inhibitor GBR 12909 was neuroprotective. Finally, both Schmidt *et al.* (1990b) and Hewitt & Green (1994) have shown that acute administration of the dopamine antagonist haloperidol together with MDMA provides effective protection against neurotoxic damage. It is acknowledged that none of these studies demonstrate authoritatively the involvement of dopamine in the MDMA-induced neurodegeneration because none of the compounds used in these investigations (haloperidol, 6-OHDA, α -methyl p-tyrosine) selectively alter dopamine function. Nevertheless, when taken with other evidence (see below) that MDMA administration alters dopamine biochemistry in the brain, the interpretation that dopamine was involved in the mechanisms by which MDMA causes neurodegeneration of 5-HT nerve terminals appeared plausible.

Whilst MDMA has generally been reported to not produce neurotoxic degeneration of dopamine neurones (Stone *et al.*, 1986; Schmidt, 1987; Schmidt & Kehne, 1990; Colado *et al.*, 1997a, b), both *in vitro* and *in vivo* studies have found that MDMA enhances dopamine efflux from cerebral tissue. For example MDMA increases release of [3 H]-dopamine from striatal slices (Johnson *et al.*, 1986; Schmidt *et al.*, 1987) and the drug has also been reported to inhibit [3 H]-dopamine uptake into striatal synaptosomes (Steele *et al.*, 1987). Use of *in vivo* voltammetry (Yamamoto & Spanos, 1988) and *in vivo* microdialysis (Hiramatsu & Cho, 1990; Nash, 1990; Nash & Brodtkin, 1991) has demonstrated that MDMA increases dopamine release and these results are supported by the observation that MDMA produces an acute increase in striatal dopamine content and a decrease in the dihydroxyphenylacetic acid (DOPAC) content (Schmidt *et al.*, 1986; Johnson *et al.*, 1991; Colado & Green 1994).

These data led us to suggest that the reason that drugs which enhanced GABA function such as pentobarbitone (Colado & Green, 1994) and clomethiazole (Colado *et al.*, 1993) were neuroprotective was because their administration resulted in an inhibition of dopamine release (Colado & Green, 1994; Green *et al.*, 1995). Recently however it has been apparent that some re-evaluation is necessary of reports on the neuroprotective action of various compounds. This is because it is now clear that some drugs attenuate damage solely because they either induce hypothermia (Farfel & Seiden, 1995a, b; Malberg *et al.*, 1996) or merely prevent the MDMA-induced hyperthermia (Colado *et al.*, 1998b). In particular the report that α -methyl p-tyrosine is neuroprotective because of its hypothermic effect (Malberg *et al.*, 1996) and our recent observation that pentobarbitone is not protective when the animals are kept warm (Colado *et al.*, 1998a) has encouraged us both to re-examine our earlier data on haloperidol and also to try and clarify further the possible role of dopamine in the neurodegenerative changes which follow MDMA administration.

Methods

Animals, drugs and reagents

Adult male Dark Agouti rats (Interfauna, Barcelona, Spain) weighing 150–170 g were used. They were housed in groups of five, in conditions of constant temperature ($21^{\circ}\text{C} \pm 2^{\circ}\text{C}$) and a 12 h light/dark cycle (lights on: 07 h 00 min) and given free access to food and water.

The following drugs were used: ($\pm$) 3,4-methylenedioxymethamphetamine HCl (Ministry of Health, Spain), clo-

methiazole edisylate (Astra Arcus, Södertälje, Sweden), haloperidol (Sigma-Aldrich Química S.A., Spain), L-3,4-dihydroxyphenylalanine methyl ester (Methyl L-DOPA, Sigma-Aldrich Química S.A., Spain) and benserazide (Sigma-Aldrich Química S.A., Spain).

In all the experiments using L-DOPA, the dose of L-DOPA was preceded 1 min earlier by an injection of benserazide, a peripheral DOPA decarboxylase inhibitor, at a dose of 6.25 mg kg^{-1} i.p. in order to avoid the peripheral metabolism of L-DOPA.

All compounds except haloperidol were dissolved in 0.9% w/v NaCl (saline). Haloperidol was suspended in peanut oil. Control animals were injected with the appropriate vehicle. All drugs were injected i.p. in a volume of 1 ml kg^{-1} . Doses are always quoted in terms of the base.

Measurement of rectal temperature

Temperature was measured by use of a digital readout thermocouple (Type K thermometer, Portec, U.K.) with a resolution of $\pm 0.1^{\circ}\text{C}$ and accuracy of $\pm 0.2^{\circ}\text{C}$ attached to a CAC-005 Rodent Sensor which was inserted 2.5 cm into the rectum, the rat being lightly restrained by holding in the hand. A steady readout was obtained within 10 s of probe insertion.

Elevation of body temperature with a homeothermic blanket

In one experiment rats were injected with MDMA + haloperidol and had their rectal temperature kept elevated to near that seen in rats given MDMA alone. This was achieved by placing the rats in a cage with a Harvard Homeothermic Blanket (Model 50-7087) covering the base.

Measurement of monoamines and their metabolites in cerebral tissue

Rats were killed by cervical dislocation and decapitation, the brains rapidly removed and cortex, hippocampus and striatum dissected out on ice. Tissue was homogenized and 5-hydroxytryptamine (5-HT), 5-hydroxyindoleacetic acid (5-HIAA), dopamine, homovanillic acid (HVA) and 3,4-dihydroxyphenylacetic acid (DOPAC) measured by high performance liquid chromatography (h.p.l.c.). Briefly, the mobile phase consisted of: KH_2PO_4 0.05 M, octanesulphonic acid 0.16 mM, EDTA 0.1 mM and methanol (16%), and was adjusted to pH 3 with phosphoric acid, filtered and degassed. The flow rate was 1 ml min^{-1} and the working electrode potential was set at 0.85 V.

The h.p.l.c. system consisted of a pump (Waters 510) linked to an automatic sample injector (Loop 200 μl , Waters 712 WISP), a stainless steel reversed-phase column (Spherisorb ODS2, 5 μm , $150 \times 4.6 \text{ mm}$) with a precolumn and an amperometric detector (Waters M460). The current produced was monitored by using an integrator (Waters M745).

[3 H]-Paroxetine binding in tissue homogenates

[3 H]-Paroxetine binding was measured by the method described in detail by Hewitt & Green (1994). The animals were killed, the brain rapidly removed and dissected on ice within 2 min. Tissue from individual animals was homogenized in ice-cold Tris-HCl (50 mM; pH 7.4) containing (in mM): NaCl 120 and KCl 5, using an Ultra-Turrax. The homogenate was centrifuged at $30,000 \times g$ for 10 min at 4°C . The supernatant was discarded and the wash procedure

tälje, Sweden) was resuspended in a buffer at a concentration of 10 mg tissue ml⁻¹. In order to obtain an estimate of the maximal density of [³H]-paroxetine-labelled 5-HT uptake sites the assay solution (1 ml) contained saturating concentration of [³H]-paroxetine (1 nM) and 100 µl tissue preparation with the addition of 5-HT (100 µM) for determination of non specific binding. Incubation was for 30 min at room temperature. Assays were terminated by rapid filtration and counting of the radioactivity by scintillation spectrometry. Protein concentrations were measured by the method of Lowry *et al.* (1951).

Implantation of microdialysis probe in the striatum

Rats were anaesthetized with sodium pentobarbitone (Euta-Lender, 40 mg kg⁻¹ i.p.) and secured in a Kopf stereotaxic frame with the tooth bar at -3.3 mm below the interaural line. The dialysis probe (3.5 mm × 200 µm; Cuprophane) was implanted in the right striatum +7.9 mm from the interaural line, -2.5 mm lateral and -8 mm below the surface of the brain (König & Klippel, 1963). Probes were secured to the skull as described by Baldwin *et al.* (1994).

Measurement of dopamine and their metabolites in the striatal dialysate

Twenty-four hours after implantation, probes were perfused with artificial cerebrospinal fluid (in mM): KCl 2.5, NaCl 125, MgCl₂·6H₂O 1.18, CaCl₂·2H₂O 1.26, at a rate of 1 µl min⁻¹. Samples collected from the freely moving animals at 10 min intervals in tubes containing 5 µl of a solution composed of HClO₄ (0.01 M), cysteine (0.2%) and sodium metabisulphite (0.2%). The first 60 min sample was discarded and the next three 30 min baseline samples collected.

Dopamine, DOPAC and HVA were measured in the dialysate by h.p.l.c. and electrochemical detection. The mobile phase consisted of KH₂PO₄ (0.05 M), octanesulphonic acid (1 mM), EDTA (0.1 mM) and methanol (14%), and was adjusted to pH 3 with phosphoric acid, filtered and degassed. The flow rate was 1 ml min⁻¹. The h.p.l.c. system consisted of a pump (Waters 510) linked to manual sample injector (Loop 10 µl Rheodyne), a stainless steel reversed-phase column (Spherisorb ODS2, 5 µm, 150 × 4.6 mm) with a precolumn and a coulometric detector (Coulchem 5100A) with a 5011 analytical cell. The working electrode potential was set at 100 mV with 500 nA gain. The current produced was monitored by using a computer data handling system (AXXIOM 747).

The limit of the sensitivity of these assays was approximately 8 pg (dopamine), 30 pg (DOPAC) and 125 pg (HVA). The per cent recovery of a solution containing dopamine (1 pg µl⁻¹), DOPAC (100 pg µl⁻¹) and HVA (150 pg µl⁻¹) was 26.5 ± 5.5% (n=4) for dopamine, 25.8 ± 1.7% (n=4) for DOPAC and 28.0 ± 1.4% (n=4) for HVA. The percentage of recovery is maintained constant for the following ranges of concentrations in the external medium: dopamine 1–27 pg µl⁻¹, DOPAC 100–2700 pg µl⁻¹, and HVA 150–4050 pg µl⁻¹. These ranges were comparable to the amounts of dopamine and metabolites recovered from the brain *in vivo*.

Implantation of microdialysis probe in the hippocampus

Rats were anaesthetized with sodium pentobarbitone ('Euta-Lender', 40 mg kg⁻¹ i.p.) and secured in a Kopf stereotaxic frame with the tooth bar at -3.3 mm below

the interaural zero. The dialysis probe (3.5 mm × 200 µm; Cuprophane) was implanted in the right hippocampus +2.2 mm from the interaural line, -4.3 mm lateral and -8 mm below the surface of the brain (König & Klippel, 1963). Probes were secured to the skull as described by Baldwin *et al.* (1994).

Measurement of free radical formation in vivo using microdialysis

Free radical formation in the brain *in vivo* was determined by the method recently described in detail by Colado *et al.*




Figure 1 Effect of haloperidol (Hal) on MDMA-induced hyperthermia. (a) Rectal temperature of rats injected with saline or MDMA (15 mg kg⁻¹ i.p.) at time zero and rats injected with haloperidol (2 mg kg⁻¹ i.p.) at -5 min and +55 min and treated with MDMA at time zero. There was no difference in the basal temperature of the groups. MDMA produced a significant rise in body temperature ($F(1,10)=19.89$, $P<0.01$) compared with the saline injected group. Administration of haloperidol to MDMA-treated rats prevented the MDMA-induced hyperthermia ($F(1,11)=27.16$, $P<0.001$). (b) Effect of haloperidol on MDMA-induced hyperthermia when rats injected with haloperidol+MDMA were kept at high ambient temperature. Temperature of rats given haloperidol and saline is also shown. There was no difference in basal temperature of the groups. Using the homeothermic blanket, the rectal temperature of the haloperidol+MDMA-treated rats was significantly higher than that of the saline group ($F(1,10)=141.69$, $P<0.001$). All results shown as means ± s.e. mean of $n=4-6$ rats.

(1997b). The method relies on the fact that hydroxyl free radicals react with salicylic acid to generate 2,3- and 2,5-dihydroxybenzoic acids (2,3-DHBA and 2,5-DHBA) and this reaction is utilized by measuring the formation of 2,3- and 2,5-DHBA in the dialysate of a microdialysis probe implanted in the hippocampus (see above) and which is being perfused with salicylic acid (see Chieuh *et al.*, 1992; Giovanni *et al.*, 1995). Twenty-four hours after implantation, probes were perfused with artificial cerebrospinal fluid (in mM): KCl 2.5, NaCl 125, $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ 1.18, $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ 1.26, containing salicylic acid (0.5 mM) at a rate of $1 \mu\text{l min}^{-1}$ and samples collected from the freely moving animals at 30 min intervals. The first 60 min sample was

discarded and the next three 30 min baseline samples collected.

The two compounds 2,3-DHBA and 2,5-DHBA were measured by h.p.l.c. and electrochemical detection. The mobile phase consisted of KH_2PO_4 (0.025 M), acetonitrile (20%) and methanol (10%) and was adjusted to pH 3.7 with phosphoric acid, filtered and degassed. The flow rate was 1 ml min^{-1} .

The h.p.l.c. system consisted of a pump (Waters 510) linked to manual sample injector (Loop $20 \mu\text{l}$ Rheodyne), a stainless steel reversed-phase column ($250 \times 4.6 \text{ mm}$, $5 \mu\text{m}$ C8 Ultracarb, Phenomenex) with a precolumn ($30 \times 4.60 \text{ mm}$, $5 \mu\text{m}$ C8 Ultracarb, Phenomenex) and a coulometric detector (Coulo-




Figure 2 Indole concentrations in (a) hippocampus, (b) cortex and (c) striatum and (d) density of [³H]-paroxetine labelled 5-HT uptake sites in cortex 7 days following saline or haloperidol (Hal, 2 mg kg^{-1} i.p.) 5 min before and 55 min after MDMA (15 mg kg^{-1} i.p.). Results shown as means $\pm$ s.e. mean, $n = 4-7$. Different from saline-treated: * $P < 0.05$, ** $P < 0.01$. Different from MDMA-treated: $\Delta P < 0.05$, $\Delta\Delta P < 0.01$.

baseline samples

2,5-DHBA were
1 detection. The
5 M), acetonitrile
ed to pH 3.7 with
the flow rate was

Vaters 510) linked
odyne), a stainless
5 μ m C8 Ultra-
1.60 mm, 5 μ m C8
detector (Coulo-

chem 5100A) with a 5011 analytical cell. The working electrode potential was set at 400 mV with 1 μ A gain. The current produced was monitored by using a computer data handling system (AXXIOM 747).

Statistics

Statistical analyses of the microdialysis experiments were performed using the statistical computer package BMDP/386 Dynamic (BMDP Statistical Solutions, Cork, Eire). Data were analysed by analysis of variance (ANOVA) with repeated measures (program 2 V) or, where missing values occurred, an unbalanced repeated measure model (program 5 V) was used. Both used treatment as the between subjects factor and time as the repeated measure. ANOVA was performed on both pre-treatment and post-treatment data. Temperature data were also analysed by ANOVA with repeated measures. Data from

monoamine and [3 H]-paroxetine binding experiments were analysed by the Newman-Keuls test.

Results

The effect of haloperidol on rectal temperature of MDMA-treated rats kept at normal ambient temperature

Administration of MDMA (15 mg kg⁻¹ i.p.) produced a clear and sustained hyperthermia (Figure 1a). In contrast, when haloperidol (2 mg kg⁻¹ i.p.) was injected 5 min before and 55 min after the MDMA injection the hyperthermic response was totally abolished (Figure 1a). Administration of haloperidol (2 mg kg⁻¹ i.p.) alone had no effect on rectal temperature (Figure 1b).




Figure 3 Indole concentrations in (a) hippocampus, (b) cortex and (c) striatum and (d) density of 3 H-paroxetine labelled 5-HT uptake sites in cortex 7 days following saline or haloperidol (Hal, 2 mg kg⁻¹ i.p.) 5 min before and 55 min after saline or MDMA (15 mg kg⁻¹ i.p.). The MDMA + haloperidol group was kept at high ambient temperature during the 6 h after the first haloperidol injection. Results shown as means $\pm$ s.e.mean, $n = 4-6$. Different from saline-treated: * $P < 0.05$, ** $P < 0.01$. Different from MDMA-treated: $\Delta P < 0.05$.

e labelled 5-HT
1 after MDMA
Different from

The effect of haloperidol on the cerebral monoamine content and [³H]-paroxetine binding in MDMA-treated rats kept at normal ambient temperature

MDMA (15 mg kg⁻¹ i.p.) administration produced a substantial loss in 5-HT and 5-HIAA concentration in the three brain areas examined and also in [³H]-paroxetine binding values in the cortex (Figure 2) 7 days later. Haloperidol administration provided significant protection against this neurotoxic degeneration (Figure 2).

The effect of haloperidol on rectal temperature of MDMA-treated rats kept at high ambient temperature

The experiment above was repeated except that the rats which were injected with both MDMA and haloperidol were kept at high ambient temperature (see Methods) in order to prevent the decrease in rectal temperature that occurred when this group was kept at normal ambient temperature. This aim was achieved (Figure 1b).

The effect of haloperidol on the cerebral monoamine content and [³H]-paroxetine binding in MDMA-treated rats kept at high ambient temperature

Administration of MDMA (15 mg kg⁻¹ i.p.) again produced a substantial loss in 5-HT and 5-HIAA concentration and [³H]-paroxetine binding in the brain 7 days later (Figure 3). However little or no protection was seen in the group given haloperidol plus MDMA when the temperature of the rats had been kept elevated to match the temperature seen in the rats given only MDMA (Figure 3).

Effect of L-DOPA on rectal temperature of MDMA treated rats

Administration of L-DOPA (25 mg kg⁻¹ + benserazide 6.25 mg kg⁻¹) had no effect on rectal temperature (Figure 4). However, when this dose was administered 2 h after MDMA (15 mg kg⁻¹) it extended the duration of the major MDMA-induced hyperthermia (Figure 4). Higher doses of L-DOPA or administering this dose of the compound at the same time as MDMA resulted in death of the animals, apparently as the result of severe hyperthermia so such protocols could not be investigated further.

Effect of MDMA and L-DOPA on dopamine metabolism

Injection of MDMA (15 mg kg⁻¹) produced a rapid and substantial increase in extracellular dopamine in the striatum, as measured by microdialysis (Figure 5a). This peak was followed by a sustained decrease in both HVA and DOPAC, the decrease in the latter starting earlier than that of HVA (1 h after MDMA, the DOPAC and HVA levels compared to basal values were 23 ± 2% and 60 ± 3%, respectively, *P* < 0.01) (Figure 5b and c). Injection of L-DOPA (25 mg kg⁻¹ + benserazide 6.25 mg kg⁻¹) alone produced no change in the concentration of dopamine in the dialysate, but did produce a sustained and large increase in both HVA and DOPAC (Figure 5b and c). When L-DOPA was injected 2 h after MDMA it did produce a clear increase in dopamine, thereby enhancing the size of the total dopamine response (Figure 5a). A prolonged increase in both HVA and DOPAC was also seen, thereby reversing the MDMA-induced decrease in the concentration of these two metabolites (Figure 5b and c).




Figure 4 Rectal temperature of rats receiving L-DOPA (25 mg kg⁻¹ i.p. + benserazide 6.25 mg kg⁻¹ i.p.) 2 h after MDMA (15 mg kg⁻¹ i.p.) or saline. Also shown are the rectal temperature of animals injected with saline instead of L-DOPA and administered 2 h before saline or MDMA. There was no difference in the basal temperature of the groups. MDMA produced a significant rise in body temperature (*F*(1,9) = 129.36, *P* < 0.001) compared with the saline injected group. Administration of L-DOPA to MDMA-treated rats produced a prolongation of the hyperthermic response induced by MDMA (*F*(1,11) = 4.12, *P* < 0.05).

Effect of MDMA and L-DOPA on free radical formation in the brain

In confirmation of previous findings (see Discussion) MDMA injection increased the concentration of 2,3- and 2,5-DHBA in the dialysate from a microdialysis probe implanted in the hippocampus and perfused with salicylic acid (Figure 6). L-DOPA (25 mg kg⁻¹ i.p. + benserazide 6.25 mg kg⁻¹ i.p.) alone did not produced significant changes in the concentration of 2,3 DHBA or 2,5 DHBA (Figure 6). The increase in formation of 2,3- and 2,5-DHBA resulting from MDMA injection was unaltered by an injection of L-DOPA (25 mg kg⁻¹) 2 h later (Figure 6).

The effect of L-DOPA on MDMA-induced degeneration of 5-HT neurones


A single injection of MDMA (15 mg kg⁻¹) resulted in a major loss of 5-HT and 5-HIAA in the hippocampus, cortex and striatum 7 days later and a similar loss of [³H]-paroxetine binding in the cortex (Figure 7). The degenerative loss of 5-HT was similar in rats injected with MDMA and those given MDMA plus L-DOPA, while L-DOPA alone was without effect on the concentration of either the indole concentration or [³H]-paroxetine binding in the brain (Figure 7). MDMA injection either alone or with L-DOPA also had no effect on the concentration of dopamine or DOPAC in the striatum 7 days later (Table 1).

Effect of low dose MDMA and L-DOPA on rectal temperature, dopamine metabolism and subsequent neurotoxicity of cerebral 5-HT neurones

Rats were injected with L-DOPA (25 mg kg⁻¹ + benserazide 6.25 mg kg⁻¹) followed 30 min later with a sub-neurotoxic dose of MDMA (5 mg kg⁻¹) to examine whether pretreatment with the amino acid would result in MDMA now producing toxicity. Pretreatment with L-DOPA did not enhance the hyperthermic effect of MDMA (Figure 8). Although L-DOPA did not enhance the MDMA-induced increase in dopamine in the striatal dialysate, it induced marked increases in both HVA and DOPAC (Figure 5b and c).

Figure 5 (C) Dopamine (DOPAC) in the striatal dialysate of rats injected with L-DOPA (25 mg kg⁻¹ + benserazide 6.25 mg kg⁻¹) alone or 2 h after MDMA (15 mg kg⁻¹). The increase in HVA and DOPAC levels compared to basal values were 23 ± 2% and 60 ± 3%, respectively, *P* < 0.01) (Figure 5b and c). Injection of L-DOPA (25 mg kg⁻¹ + benserazide 6.25 mg kg⁻¹) alone produced no change in the concentration of dopamine in the dialysate, but did produce a sustained and large increase in both HVA and DOPAC (Figure 5b and c). When L-DOPA was injected 2 h after MDMA it did produce a clear increase in dopamine, thereby enhancing the size of the total dopamine response (Figure 5a). A prolonged increase in both HVA and DOPAC was also seen, thereby reversing the MDMA-induced decrease in the concentration of these two metabolites (Figure 5b and c).

○ Saline
□ L-DOPA
● MDMA
■ MDMA+L-DOPA



L-DOPA (25 mg kg⁻¹)
MDMA (15 mg kg⁻¹)
temperature of animals
administered 2 h before
the basal temperature
significant rise in body
temperature with the saline
MDMA-treated rats
response induced by

free radical

discussion) MDMA
and 2,5-DHBA in
implanted in the
acid (Figure 6). L-
mg kg⁻¹ i.p.) alone
the concentration of
crease in formation
MA injection was
mg kg⁻¹) 2 h later

neurodegeneration

resulted in a major
impus, cortex and
of [³H]-paroxetine
relative loss of 5-HT
and those given
alone was without
dole concentration
Figure 7). MDMA
had no effect on
in the striatum 7

MA on rectal
subsequent
s

kg⁻¹ + benserazide
a sub-neurotoxic
ether pretreatment
A now producing
not enhance the
Although L-DOPA
use in dopamine in
cases in both HVA




Figure 5 Changes in the levels of (a) dopamine (DA), (b) homovanillic acid (HVA) and (c) 3,4-dihydroxyphenylacetic acid (DOPAC) in the striatal dialysate of rats treated with L-DOPA 25 mg kg⁻¹ i.p. + benserazide 6.25 mg kg⁻¹ i.p. or saline 2 h after MDMA (15 mg kg⁻¹ i.p.). MDMA produced an increase in the content of DA ($F(1,9)=74.34$, $P<0.001$) and a decrease in the levels of HVA ($F(1,9)=81.07$, $P<0.001$) and DOPAC ($F(1,9)=52.61$, $P<0.001$). Administration of L-DOPA enhanced the effect of MDMA on extracellular dopamine ($F(1,10)=9.57$, $P<0.001$) and reversed the MDMA-induced decrease in HVA ($F(1,10)=66.72$, $P<0.001$) and DOPAC ($F(1,10)=43.10$, $P<0.001$). L-DOPA, when given alone, increased the extracellular levels of HVA ($F(1,10)=27.30$, $P<0.001$) and DOPAC ($F(1,10)=24.54$, $P<0.001$). Values are expressed as a percentage of the mean of three measurements before drug administration. Each value is the mean $\pm$ s.e. mean of 5–7 experiments. The basal concentrations in saline-

and DOPAC in place of the decrease in the concentration of these metabolites produced by MDMA alone (Figure 9). The low dose of MDMA did not result in neurodegeneration occurring 7 days later, nor did the pretreatment with L-DOPA result in this dose of MDMA producing neurodegeneration (Figure 10).

Effect of clomethiazole on the MDMA-induced change in dopamine metabolism in vivo

The increase in dopamine and decrease in HVA and DOPAC induced by MDMA (15 mg kg⁻¹) was totally unaltered by administration of clomethiazole (50 mg kg⁻¹) 5 min prior and 55 min post MDMA injection (Figure 11).




Figure 6 Changes in the levels of (a) 2,3 dihydroxybenzoic acid (DHBA) and (b) 2,5 DHBA in the hippocampal dialysate of rats treated with L-DOPA 25 mg kg⁻¹ i.p. + benserazide 6.25 mg kg⁻¹ i.p. or saline 2 h after MDMA (15 mg kg⁻¹ i.p.). MDMA increased the extracellular levels of 2,3 DHBA ($F(1,11)=10.34$, $P<0.001$) and 2,5 DHBA ($F(1,11)=4.66$, $P<0.05$). This effect was not modified by L-DOPA injection 2 h after MDMA. L-DOPA, when given to saline-treated rats did not significantly modify the extracellular levels of 2,3 DHBA or 2,5 DHBA. Values are expressed as a percentage of the mean of three measurements before drug administration. Each value is the mean $\pm$ s.e. mean of 6–10 experiments. The basal concentrations in saline-treated rats ($n=6$) were: 2,3 DHBA (10.1 ± 1.1 pg μ l⁻¹) and 2,5 DHBA (11.6 ± 1.22 pg μ l⁻¹).

treated rats were: DA (0.83 ± 15 pg μ l⁻¹), HVA (246 ± 16 pg μ l⁻¹) and DOPAC (252 ± 23 pg μ l⁻¹).




Figure 7 Indole concentrations in (a) hippocampus, (b) cortex and (c) striatum and (d) density of ³H-paroxetine labelled 5-HT uptake sites in cortex 7 days following saline or L-DOPA (25 mg kg⁻¹ i.p. + benserazide 6.25 mg kg⁻¹ i.p.) 2 h after saline or MDMA (15 mg kg⁻¹ i.p.). Results shown as means $\pm$ s.e. mean, $n = 5-8$. Different from saline-treated: * $P < 0.05$, ** $P < 0.01$.

Table 1 The concentration of dopamine and DOPAC in striatum 7 days after injection of MDMA (15 mg kg⁻¹ i.p.) followed 2 h later by L-DOPA (25 mg kg⁻¹ i.p.)

1st injection	2nd injection	n	Catechol concentration (ng g ⁻¹ tissue)	
			Dopamine	DOPAC
Saline	Saline	5	6110 $\pm$ 423	1588 $\pm$ 212
MDMA	Saline	5	6200 $\pm$ 204	1600 $\pm$ 78
MDMA	L-DOPA	8	6771 $\pm$ 310	1403 $\pm$ 137
Saline	L-DOPA	5	6495 $\pm$ 170	1662 $\pm$ 142

Results shown as means $\pm$ s.e. mean. No significant differences in tissue concentrations of the catechols were found between experimental groups.

Discussion

Both Schmidt *et al.* (1990b) and Hewitt & Green (1994) have reported that administration of haloperidol concurrently with MDMA attenuates the neurodegeneration of 5-HT terminals in the brain produced by this neurotoxic amphetamine and this observation was confirmed by the current study. Schmidt *et al.* (1990b) did not measure body temperature but Hewitt & Green (1994) observed that the combination of haloperidol and MDMA, whilst not producing hypothermia, did markedly attenuate the hyperthermia normally produced by the drug, an




Figure 8 Rectal temperature of rats receiving L-DOPA (25 mg kg⁻¹ i.p. + benserazide 6.25 mg kg⁻¹ i.p.) 30 min before MDMA (5 mg kg⁻¹ i.p.). Also shown are the rectal temperature of animals injected with saline instead of L-DOPA and administered saline or MDMA 30 min later. There was no difference in the basal temperature of the groups. MDMA produced a rise in body temperature ($F(1,10) = 7.15$, $P < 0.05$). Pretreatment with L-DOPA did not alter the hyperthermic response induced by MDMA.

observation also confirmed by the present study. We therefore repeated the experiment on the neuroprotective effect of

haloperidol
the rectal
to that s

10
8
6
4
2
DA (% of baseline)

2
1
HVA (% of baseline)

1
1
DOPAC (% of baseline)

Figure
homov:
(DOPA
(25 mg
MDM/
MDM/
the MI
DOPA
percent
admini:
experin

haloperidol but used a homeothermic blanket, thereby keeping the rectal temperature of the animals elevated to a similar level to that seen in the group given only MDMA (Figure 1). The




Figure 9 Changes in the levels of (a) dopamine (DA), (b) homovanillic acid (HVA) and (c) 3,4-dihydroxyphenylacetic acid (DOPAC) in the striatal dialysate of rats treated with L-DOPA (25 mg kg⁻¹ i.p. + benserazide 6.25 mg kg⁻¹ i.p.) 30 min before MDMA (5 mg kg⁻¹ i.p.). L-DOPA did not modify the effect of MDMA on extracellular dopamine ($F(1,8)=0.0908$, n.s.) but reversed the MDMA-induced decrease in HVA ($F(1,9)=47.82$, $P<0.001$) and DOPAC ($F(1,10)=20.55$, $P<0.001$). Values are expressed as a percentage of the mean of two measurements before drug administration. Each value is the mean $\pm$ s.e. mean of 4–6 experiments.

protective effect of haloperidol in this hyperthermic group was marginal. These data support the findings in our other recent investigations (Colado *et al.*, 1998a, b) which have indicated strongly that drugs can provide substantial neuroprotection not only by producing hypothermia as has previously been demonstrated by others (Farfel & Seiden 1995a, b; Malberg *et al.*, 1996; Colado *et al.*, 1998a) but also by merely preventing the MDMA-induced hyperthermia (Colado *et al.*, 1998b). Haloperidol therefore can be concluded to be protective because of its effect on temperature and not because of a selective neuroprotective mechanism.

This finding in turn raised the question as to the validity of earlier proposals that dopamine plays a key role in the expression of MDMA-induced neurotoxicity. Various observations have been used to support this tenet, including the fact that MDMA alters dopamine release (Johnson *et al.*, 1986; Schmidt *et al.*, 1987; Hiramatsu & Cho, 1990; Nash, 1990; Nash & Brodtkin, 1991) and the observation that L-DOPA administration potentiates MDMA-induced damage (Schmidt *et al.*, 1991). We therefore re-investigated both these claims and also investigated whether L-DOPA administration alters the increase in free radical formation which has been shown to follow MDMA injection (Colado *et al.*, 1997b; 1998c).

With regard to the alteration of dopamine metabolism which follows MDMA injection, both Johnson *et al.* (1986) and Schmidt *et al.* (1987) noted that MDMA released [³H]-dopamine from striatal slices and Steele *et al.* (1987) reported that the drug inhibited [³H]-dopamine uptake into striatal synaptosomes. Subsequent studies using either *in vivo* voltammetry (Yamamoto & Spanos, 1988) or microdialysis (Nash, 1990; Hiramatsu & Cho, 1990; Gough *et al.*, 1991) found that MDMA administration resulted in a rapid rise in extracellular dopamine and decrease in DOPAC and HVA. Both the size and the time course of these changes as reported by Nash (1990) and Gough *et al.* (1991) were replicated in the current study. The size of the dopamine increase and DOPAC decrease are considerably greater than those observed in postmortem tissue 4 h after a single injection of MDMA (Colado & Green, 1994).

The current investigation found that following MDMA the concentration of HVA in the dialysate was decreased in a similar fashion to DOPAC, but that the change in the extracellular level of HVA lagged behind the DOPAC change. The difference in the time courses of changes in the extracellular DOPAC and HVA levels following MDMA may reflect the fact that HVA is a secondary metabolite of dopamine, being predominantly derived from intraneuronally formed DOPAC but also from the released dopamine via 3-methoxytyramine (Westerink & Spaan, 1982). Therefore, the delayed decrease in HVA compared to DOPAC might be due to an initial rise in 3-methoxytyramine and its subsequent conversion to HVA by catechol O-methyl transferase. These data suggest that MDMA exerts two separate actions on dopaminergic activity *in vivo*, one causing an increase in dopamine release and another producing a decrease in the intraneuronal dopamine metabolism (reflected by the extracellular levels of DOPAC and HVA). These changes on dopamine metabolites are probably the results of MDMA blocking dopamine re-uptake (causing less dopamine to be metabolized to DOPAC) and also by inhibiting monoamine oxidase (MAO). Dopamine is preferentially metabolized by MAO-A in the rat striatum *in vivo* (Kato *et al.*, 1986) and MDMA acts as a competitive inhibitor of MAO-A activity (Leonardi & Azmitia, 1994). An additional explanation of the decrease in DOPAC and HVA would be a reduction in dopamine synthesis following MDMA. However this hypoth-

pelled 5-HT
or saline or
<0.01.

OPA (25 mg kg⁻¹
before MDMA
temperature of animals
inistered saline or
ce in the basal
a rise in body
nt with L-DOPA
y MDMA.

ly. We therefore
ective effect of




Figure 10 Indole concentrations in (a) hippocampus, (b) cortex and (c) striatum and (d) density of ³H paroxetine labelled 5-HT uptake sites in cortex 7 days following saline or L-DOPA (25 mg kg⁻¹ i.p. + benserazide 6.25 mg kg⁻¹ i.p.) 30 min before MDMA (5 mg kg⁻¹ i.p.). Results shown as means $\pm$ s.e. mean, $n = 5-8$. Different from saline-treated: *** $P < 0.01$.

esis is unlikely since MDMA does not alter tyrosine hydroxylase activity in the striatum (Stone *et al.*, 1986).

The data raise a further point; that is, the nature of the mechanism of MDMA-induced dopamine release. It has often been assumed that the dopamine release is by a carrier-mediated exchange mechanism as occurs with other amphetamine-like drugs (Raiteri *et al.*, 1979; Fischer & Cho, 1979; also see McMillen, 1983). However it is worth noting that several investigators have provided convincing evidence that 5-HT₂ agonists enhance MDMA-induced dopamine release while 5-HT₂ antagonists inhibit release (Nash, 1990; Schmidt *et al.*, 1990a, 1994; Yamamoto *et al.*, 1995; Gudelsky *et al.*, 1994). MDMA-induced 5-HT release therefore appears to play a major role in the release of dopamine. When one adds the known effects of MDMA on dopamine uptake inhibition (Steele *et al.*, 1987) and MAO inhibition (Leonardi & Azmitia, 1994) then it becomes plausible to suggest that carrier-mediated exchange need not be a major determinant in the increase of extracellular dopamine which follows MDMA administration, a suggestion supported by recent *in vitro* studies (Crespi *et al.*, 1997). Further support for this proposal

comes from our study on the effect of clomethiazole. This drug, which enhances GABA_A receptor activity (see Green 1998), inhibited the increase in extracellular dopamine which follows administration of methamphetamine, a known carrier-mediated mechanism (McMillen, 1983). Therefore, the failure of this drug to alter the MDMA-induced increase in extracellular dopamine may reflect a difference in the release mechanism between these two amphetamine derivatives. The implications of this to the selective neurotoxic action of MDMA are discussed later.

Interestingly the size of the increase in extracellular dopamine in the striatum following MDMA seen *in vivo* was similar in magnitude to that seen after a high dose of methamphetamine (Baldwin *et al.*, 1993) which raises questions as to why the locomotor and behavioural activity seen after MDMA administration appears to be 5-HT rather than dopamine receptor mediated, having a different profile than that seen after amphetamine (Callaway *et al.*, 1990; Callaway & Geyer, 1992). It could be that the serotonin-mediated behaviours predominate over any dopamine-mediated effects. While MDMA releases dopamine in both the accumbens and nucleus accumbens, it is possible that the release in the accumbens is more significant.

Figure 10
metabolite
treated v
before a:
did not
extracellu
of three
means $\pm$ s

Callawa
this pape
following
However
MDMA
neuronal
Ca²⁺ de
dopamine
allowing
including
1989; Col
When
that extrac
the MDM
that HV/
markedly
reflect incr
clear *in vi*
possibly al
city was ob
of the repr
neurotoxic
colleagues
impossible
trial. This
train of rat
neurotoxicit
of MDMA
required wit
Murray *et al*
dose of L-D
was detected
rats given a s
DOPA, as w
In order
between our
examined w/
medical form
substrate cor
viously be




Figure 11 Changes in the levels of dopamine (DA) and the metabolites (shown in the insert) in the striatal dialysate of rats treated with clomethiazole (CMZ, 50 mg kg⁻¹ i.p.) or saline 5 min before and 55 min after MDMA (15 mg kg⁻¹ i.p.). Clomethiazole did not modify the effect of MDMA on DA, DOPAC and HVA extracellular levels. Values are expressed as a percentage of the mean of three measurements before drug administration. Each value is the mean $\pm$ s.e. mean of 6–7 experiments.

Callaway & Geyer, 1992) and striatal regions (Nash, 1990; this paper) the accumbens is the locus for locomotor activity following amphetamine administration (Kelly *et al.*, 1975). However, recent data suggest that the predominant action of MDMA in this region may be that of inhibiting dopamine neuronal firing (Gifford *et al.*, 1996). Since the drug can induce Ca²⁺ dependent dopamine release (Crespi *et al.*, 1997) dopamine mediated function would be suppressed, thereby allowing the expression of 5-HT mediated behaviours including the serotonin syndrome (Spanos & Yamamoto, 1989; Colado *et al.*, 1993; Green *et al.*, 1995).

When L-DOPA was given 2 h after MDMA we observed that extracellular dopamine was increased above that seen in the MDMA alone treated group for approximately 3 h and that HVA and DOPAC concentrations also increased markedly over a similar period. These changes presumably reflect increased dopamine synthesis and release. Despite this clear *in vivo* indication of increased dopamine release and possibly also function, no enhancement of MDMA neurotoxicity was observed 7 days later. These data are in contradiction of the report of Schmidt *et al.* (1991) that MDMA-induced neurotoxicity is enhanced by L-DOPA. While Schmidt and colleagues used higher doses of L-DOPA these proved impossible to administer in the current study as they proved fatal. This is probably because of the use of the Dark Agouti strain of rat in our study since we have previously found that neurotoxicity occurs in these animals after much lower doses of MDMA and p-chloroamphetamine (PCA) than are required with other strains (see Colado *et al.*, 1995; 1997b; Murray *et al.*, 1996). Nevertheless using the highest practical dose of L-DOPA, no potentiation of MDMA-induced toxicity was detected. We were also unable to induce neurotoxicity in rats given a sub-toxic dose of MDMA by also administering L-DOPA, as was also reported by Schmidt *et al.* (1991).

In order to investigate further this apparent discrepancy between our data and that of Schmidt *et al.* (1991) we also examined whether L-DOPA altered MDMA-induced free radical formation. The MDMA-induced increase in the dialysate concentration of 2,3- and 2,5-DHBA that has previously been reported to occur after MDMA administra-

tion (Colado *et al.*, 1997b; 1998c) was again seen in the current study. While the concentration of 2,5-DHBA appeared to be increased by L-DOPA, this change was not statistically significant. In any event, as Halliwell *et al.* (1991) have pointed out, a 2,3-DHBA increase is the only reliable indicator of increased free radical formation because of enzyme-induced conversion of salicylate to 2,5-DHBA. Nevertheless we found no evidence that the MDMA-induced increase in either 2,3- or 2,5-DHBA was even modestly enhanced by L-DOPA administration. Thus one plausible explanation for the results obtained by Schmidt *et al.* (1991), namely that tissue damage resulting from free radicals produced by MDMA metabolism (see Colado *et al.*, 1997b) would be further potentiated by administration of L-DOPA, a compound that through its metabolism to dopamine would also increase free radical production through auto-oxidation (see Chiueh *et al.*, 1992; 1993), could not be supported.

What does remain a possible explanation for the discrepant findings is the degree of hyperthermia produced by addition of L-DOPA to MDMA-treated rats. As we and others have shown, sustained hyperthermia markedly enhances the neurotoxic damage produced by MDMA (Colado *et al.*, 1998b; Broening *et al.*, 1995) presumably because hyperthermia is conducive to enhancing free radical production (Kil *et al.*, 1996). If L-DOPA produced a sustained hyperthermia in the strain of rats used by Schmidt *et al.* (1990) rather than the more transient effect we observed, then neurotoxic damage would probably be increased. However, such a mechanism does not indicate that dopamine has a core role in MDMA-induced neurodegeneration as has been proposed by others (Stone *et al.*, 1988; Schmidt *et al.*, 1991). In this regard it is also worth pointing out that in our current study the increase in extracellular dopamine was almost the same whether a neurotoxic dose of MDMA of 15 mg kg⁻¹ or a non-toxic dose of 5 mg kg⁻¹ was given.

Finally we examined the effect of injecting clomethiazole on MDMA-induced changes in dopamine metabolism. Previously in an *in vivo* study we found that clomethiazole inhibited both the increase in dopamine and reduction in HVA and DOPAC which followed an injection of methamphetamine (Baldwin *et al.*, 1993). In contrast, clomethiazole appeared to have no effect on MDMA-induced changes in dopamine metabolism. This finding is perhaps surprising in the light of a report suggesting that MDMA-induced dopamine release is regulated by the activity of a GABAergic input into the substantia nigra (Yamamoto *et al.*, 1995). Clomethiazole is one of the few compounds to have been shown unequivocally to protect against MDMA-induced damage by a mechanism which does not involve a reduction in body temperature (Colado *et al.*, 1998b). It is also a compound that has been shown to have a powerful neuroprotective action in a wide variety of animal models of acute ischaemic stroke (Green, 1998). These data suggest that dopamine is not involved in the neuroprotective effect of clomethiazole against MDMA-induced neurotoxicity.

Finally it is worth pointing out that a major conundrum is raised by the current study. In rats, but not mice (see Laverty & Logan 1990), MDMA produces a neurodegenerative loss of 5-HT nerve terminals, whilst sparing dopamine terminals (Schmidt & Kehne, 1990; Colado *et al.*, 1997a, b), a fact again supported by the findings in this current study (see Table 1). The question arises as to the mechanisms involved in this selectivity. It is noteworthy that methamphetamine which also increases extracellular 5-HT and dopamine (see for example Baldwin *et al.*, 1993) and also increases free radical formation (Giovanni *et al.*, 1995) produces neurotoxic degeneration of both 5-HT and dopamine neurones (Gibb *et al.*, 1990; Green *et*

al., 1992; Baldwin *et al.*, 1993). If we are correct and the extracellular dopamine increase does not reflect carrier-mediated exchange then relatively little MDMA may be entering the dopamine nerve ending and being metabolised to the neurotoxic metabolites which are then auto-oxidised thereby producing free radicals (see Colado *et al.*, 1997b). This may, in part, explain its lack of neurotoxic action on dopamine nerve endings since it would be reasonable to assume that MDMA would be metabolized in the same way in dopamine and 5-HT neurones. Against this hypothesis however there is good evidence that dopamine uptake inhibitors markedly inhibit MDMA-induced dopamine release (Nash & Brodtkin, 1991). This observation would strongly support the idea that MDMA acts to release dopamine by a facilitated exchange mechanism which utilises the dopamine uptake carrier mechanism (see Raiteri *et al.*, 1979; Fischer & Cho, 1979; McMillen, 1983).

Another possible explanation for the selective damage which follows MDMA is that the neurotoxic metabolites of

methamphetamine and MDMA differ in their pattern of damage. A major postulated neurotoxic metabolite of MDMA, namely dihydroxymethylamphetamine (Hiramatsu *et al.*, 1990; Colado *et al.*, 1995, 1997b), has been reported not to be toxic to dopamine neurones (Johnson *et al.*, 1992).

In conclusion, the current results demonstrate that MDMA has a major acute effect on dopamine metabolism in the brain but we suggest that there is little evidence to indicate that the changes reported are crucially involved in the long term neurotoxic degeneration of 5-HT neurones produced by this commonly used recreational drug.

M.I. Colado thanks CICYT (SAF 98-0074) and Astra Arcus for financial support. The authors are grateful to Servicio de Restriccion de Estupefacientes, Ministry of Health, Spain for supplying MDMA.

References

- BALDWIN, H.A., COLADO, M.I., MURRAY, T.K., DE SOUZA, R.J. & GREEN, A.R. (1993). Striatal dopamine release *in vivo* following neurotoxic doses of methamphetamine and effect of the neuroprotective drugs, chlormethiazole and dizocilpine. *Br. J. Pharmacol.*, **108**, 590–596.
- BALDWIN, H.A., WILLIAMS, J.L., SNARES, M., FERREIRA, T., CROSS, A.J. & GREEN, A.R. (1994). Attenuation by chlormethiazole administration of the rise in extracellular amino acids following focal ischaemia in the cerebral cortex of the rat. *Br. J. Pharmacol.*, **112**, 188–194.
- BATTAGLIA, G., YEH, S.Y., O'HEARN, E., MOLLIVER, M.E., KUJAR, K.J. & DE SOUZA, E.B. (1987). 3,4-methylenedioxymphetamine destroys serotonin terminals in rat brain: quantification of neurodegeneration by measurement of [³H]-paroxetine-labelled serotonin uptake sites. *J. Pharmacol. Exp. Ther.*, **242**, 911–916.
- BROENING, H.W., BOWYER, J.F. & SLIKKER JR. W. (1995). Age dependent sensitivity of rats to the long term effects of the serotonergic neurotoxicant (±)-3,4-methylenedioxymphetamine (MDMA) correlates with the magnitude of the MDMA-induced thermal response. *J. Pharmacol. Exp. Ther.*, **275**, 325–333.
- CALLAWAY, C.W. & GEYER, M.A. (1992). Stimulant effects of 3,4-methylenedioxymphetamine in the nucleus accumbens of rats. *Eur. J. Pharmacol.*, **214**, 45–51.
- CALLAWAY, C.W., WING, L.L. & GEYER, M.A. (1990). Serotonin release contributes to the locomotor stimulant effects of 3,4-methylenedioxymphetamine in rats. *J. Pharmacol. Exp. Ther.*, **254**, 456–464.
- CHIUH, C.C., KRISHNA, G., TULSI, P., OBATA, T., LANG, K., HUANG, S.J. & MURPHY, D.L. (1992). Intracranial microdialysis of salicylic acid to detect hydroxyl radical generation through dopamine autooxidation in the caudate nucleus: effect of MPP⁺. *Free Rad. Biol. Med.*, **13**, 581–583.
- CHIUH, C.C., MIYAKE, H. & PENG, M.-T. (1993). Role of dopamine autooxidation, hydroxyl radical generation and calcium overload in underlying mechanisms involved in MPTP-induced Parkinsonism. *Adv. Neurol.*, **60**, 251–258.
- COLADO, M.I., ESTEBAN, B., O'SHEA, E., GRANADOS, R. & GREEN, A.R. (1998a). Studies on the protective effect of pentobarbitone on MDMA-induced neurodegeneration. *Psychopharmacology*. (In press).
- COLADO, M.I., GRANADOS, R., O'SHEA, E., ESTEBAN, B. & GREEN, A.R. (1998b). Role of hyperthermia in the protective action of clomethiazole against MDMA ('ecstasy')-induced degeneration, comparison with the novel NMDA channel blocker AR-R15896. *Br. J. Pharmacol.*, **124**, 479–484.
- COLADO, M.I., O'SHEA, E., ESTEBAN, B., GRANADOS, R. & GREEN, A.R. (1998c). *In vivo* evidence against clomethiazole being neuroprotective against MDMA ('ecstasy')-induced degeneration of rat brain 5-HT nerve terminals by a free radical scavenging mechanism. *Neuropharmacology*. (In press).
- COLADO, M.I. & GREEN, A.R. (1994). A study of the mechanism of MDMA ('ecstasy')-induced neurotoxicity of 5-HT neurones using chlormethiazole, dizocilpine and other protective compounds. *Br. J. Pharmacol.*, **111**, 131–136.
- COLADO, M.I., MURRAY, T.K. & GREEN, A.R. (1993). 5-HT loss in rat brain following 3,4-methylenedioxymphetamine (MDMA), p-chloroamphetamine and fenfluramine administration and effects of chlormethiazole and dizocilpine. *Br. J. Pharmacol.*, **108**, 583–589.
- COLADO, M.I., O'SHEA, E., GRANADOS, R., MISRA, A., MURRAY, T.K. & GREEN, A.R. (1997a). A study of the neurotoxic effect of MDMA ('ecstasy') on 5-HT neurones in the brains of mothers and neonates following administration of the drug during pregnancy. *Br. J. Pharmacol.*, **121**, 827–833.
- COLADO, M.I., O'SHEA, E., GRANADOS, R., MURRAY, T.K., WILLIAMS, J.L. & GREEN, A.R. (1997b). *In vivo* evidence for free radical involvement in the degeneration of rat brain 5-HT following administration of MDMA ('Ecstasy') and p-chloroamphetamine but not the degeneration following fenfluramine. *Br. J. Pharmacol.*, **121**, 889–900.
- COLADO, M.I., WILLIAMS, J.L. & GREEN, A.R. (1995). The hyperthermic and neurotoxic effects of 'Ecstasy' (MDMA) and 3,4-methylenedioxymphetamine (MDA) in the Dark Agouti (DA) rat, a model of the CYP2D6 poor metaboliser phenotype. *Br. J. Pharmacol.*, **115**, 1281–1289.
- CRESPI, D., MENNINI, T. & GOBBI, M. (1997). Carrier dependent and Ca²⁺ dependent 5-HT and dopamine release induced by (+)-amphetamine, 3,4-methylenedioxymphetamine, p-chloroamphetamine and fenfluramine. *Br. J. Pharmacol.*, **121**, 1735–1743.
- FARFEL, C.M. & SEIDEN, L.S. (1995a). Role of hypothermia in the mechanism of protection against serotonergic toxicity. I. Experiments with 3,4-methylenedioxymphetamine, dizocilpine, CGS 19755 and NBQX. *J. Pharmacol. Exp. Ther.*, **272**, 860–867.
- FARFEL, C.M. & SEIDEN, L.S. (1995b). Role of hypothermia in the mechanism of protection against serotonergic toxicity. II. Experiments with methamphetamine, p-chloroamphetamine, fenfluramine, dizocilpine and dextromethorphan. *J. Pharmacol. Exp. Ther.*, **272**, 868–875.
- FISCHER, J.F. & CHO, A.K. (1979). Chemical release of dopamine from striatal homogenates: evidence for an exchange diffusion model. *J. Pharmacol. Exp. Ther.*, **208**, 203–209.
- GIBB, J.W., JOHNSON, M., STONE, D. & HANSON, G.R. (1990). MDMA: historical perspectives. *Ann. N.Y. Acad. Sci.*, **600**, 601–611.
- GIFFORD, A.N., MINABE, Y., TOOR, A., WANG, R.Y. & ASHBY JR. C.R. (1996). Examination of the action of 3,4-methylenedioxymphetamine on rat A10 dopamine neurons. *Synapse*, **23**, 52–57.

GIOVANNI (1995) system tamine
GOUGH. effects mono-
619-6
GREEN. stroke efficac
GREEN. / pharm ymeth ogy, 1
GREEN. CROSS on 5-HT protec
315-
GUDELSON o releas
Eur. J
HALLIWILL Hydro cautic
HEWITT. and h termi
Neuro
HIRAMATA effect monc
rats: 269-
HIRAMATA Meta dihyd
gluta
JOHNSON & LIN and 2 meth
dopa
JOHNSON the e [³H]-
brain
JOHNSON neuro nergi
ymet
Beha
KATO. dialy mon
rats.
KELLY. and of th
94, 5
KIL, H.Y alter repe
KÖNIG. atlas
York
LAVERI drug
city
LEONA inhib
fenfl
10, 2
LOWRY (195
Biol

- er in their pattern of neurotoxic metabolite of methamphetamine (Hiramatsu 7b), has been reported in monkeys (Johnson et al., 1995).
- nonstrate that MDMA metabolism in the brain needs to indicate that the metabolites produced by this
- 74) and Astra Arcus for grateful to Servicio de Salud, Spain for
- study of the mechanism of toxicity of 5-HT neurones and other protective com-
- 6.
- 1993). 5-HT loss in methylenedioxymethamphetamine (MDMA) administration and dizocilpine. *Br. J. Pharmacol.*, **119**, 247-260.
- MISRA, A., MURRAY, T.K. & GREEN, A.R. (1993). The neurotoxic effect of MDMA in the brains of mothers of the drug during pregnancy. *Br. J. Pharmacol.*, **119**, 247-260.
- S. R., MURRAY, T.K. (1993). In vivo evidence for free release of rat brain 5-HT ('Ecstasy') and p-chloroamphetamine following fenfluramine.
- EN, A.R. (1995). The 'Ecstasy' (MDMA) and its metabolites in the Dark Agouti metaboliser phenotype.
- Carrier dependent and release induced by (+)-amphetamine, p-chloroamphetamine, and p-chloroamphetamine. *J. Pharmacol. Exp. Ther.*, **272**, 1735-1743.
- of hypothermia in the rat following administration of methamphetamine, dizocilpine, and p-chloroamphetamine. *J. Pharmacol. Exp. Ther.*, **272**, 1735-1743.
- of hypothermia in the rat following administration of methamphetamine, dizocilpine, and p-chloroamphetamine. *J. Pharmacol. Exp. Ther.*, **272**, 1735-1743.
- al release of dopamine from the striatum: an exchange diffusion study. *J. Pharmacol. Exp. Ther.*, **272**, 1735-1743.
- HANSON, G.R. (1990). The neurotoxic effect of MDMA in the brains of mothers of the drug during pregnancy. *Br. J. Pharmacol.*, **119**, 247-260.
- NG, R.Y. & ASHBY JR, J.R. (1993). In vivo evidence for free release of rat brain 5-HT ('Ecstasy') and p-chloroamphetamine following fenfluramine.
- ANNI, A., LIANG, L.P., HASTINGS, T.G. & ZIGMOND, M.J. (1995). Estimating hydroxyl radical content in rat brain using chemiluminescence and intraventricular salicylate: impact of methamphetamine. *J. Neurochem.*, **64**, 1819-1825.
- GH, B., ALI, S.F., SLIKKER JR, W. & HOLSON, R.R. (1991). Acute effects of 3,4-methylenedioxymethamphetamine (MDMA) on monoamines in rat caudate. *Pharmacol. Biochem. Behav.*, **39**, 619-623.
- GREEN, A.R. (1998). Clomethiazole (Zendra) in acute ischemic stroke: basic pharmacology and biochemistry and clinical efficacy. *Pharmacol. Ther.*, **80**, 123-147.
- GREEN, A.R., CROSS, A.J. & GOODWIN, G.M. (1995). Review of the pharmacology and clinical pharmacology of 3,4-methylenedioxymethamphetamine (MDMA or 'Ecstasy'). *Psychopharmacology*, **119**, 247-260.
- GREEN, A.R., DE SOUZA, R.J., WILLIAMS, J.L., MURRAY, T.K. & CROSS, A.J. (1992). The neurotoxic effects of methamphetamine on 5-hydroxytryptamine and dopamine in brain: evidence for the protective effect of chlormethiazole. *Neuropharmacology*, **31**, 315-321.
- ODELSKY, G.A., YAMAMOTO, B.K. & NASH, J.F. (1994). Potentiation of 3,4-methylenedioxymethamphetamine-induced dopamine release and serotonin neurotoxicity by 5-HT₂ receptor agonists. *Eur. J. Pharmacol.*, **264**, 325-330.
- ALLIWEILL, B., KAUR, H. & INGELMAN-SUNDBERG, M. (1991). Hydroxylation of salicylate as an assay for hydroxyl radicals: a cautionary note. *Free Rad. Biol. Med.*, **10**, 439-441.
- WITT, K.E. & GREEN, A.R. (1994). Chlormethiazole, dizocilpine and haloperidol prevent the degeneration of serotonergic nerve terminals induced by administration of MDMA ('Ecstasy'). *Neuropharmacology*, **33**, 1589-1595.
- YAMAMOTO, M. & CHO, A.K. (1990). Enantiomeric differences in the effects of 3,4-methylenedioxymethamphetamine on extracellular monoamines and metabolites in the striatum of freely-moving rats: an in vivo microdialysis study. *Neuropharmacology*, **29**, 269-275.
- YAMAMOTO, M., KUMAGAI, Y., UNGER, S.E. & CHO, A.K. (1990). Metabolism of methylenedioxymethamphetamine: formation of dihydroxymethamphetamine and a quinone identified as its glutathione adduct. *J. Pharmacol. Exp. Ther.*, **254**, 521-527.
- HANSON, M., ELAYAN, L., HANSON, G.R., FOLTZ, R.L., GIBB, J.W. & LIM, H.K. (1992). Effects of 3,4-dihydroxymethamphetamine and 2,4,5-trihydroxymethamphetamine, two metabolites of 3,4-methylenedioxymethamphetamine, on central serotonergic and dopaminergic systems. *J. Pharmacol. Exp. Ther.*, **261**, 447-453.
- HANSON, M.P., HOFFMAN, A.J. & NICHOLS, D.E. (1986). Effects of the enantiomers of MDA, MDMA and related analogues on [³H]-serotonin and [³H]-dopamine release from superfused rat brain slices. *Eur. J. Pharmacol.*, **132**, 269-276.
- HANSON, M.P., HUANG, X. & NICHOLS, D.E. (1991). Serotonin neurotoxicity in rats after combined treatment with a dopaminergic agent followed by a non-neurotoxic 3,4-methylenedioxymethamphetamine (MDMA) analogue. *Pharmacol. Biochem. Behav.*, **40**, 915-922.
- ATO, T., DONG, B., ISHII, K. & KINEMUCHI, H. (1986). Brain dialysis: In vivo metabolism of dopamine and serotonin by monoamine oxidase A but not B in the striatum of unrestrained rats. *J. Neurochem.*, **46**, 1277-1282.
- ELLY, P.H., SEVIOUR, P.H. & IVERSEN, S.D. (1975). Amphetamine and apomorphine responses in the rat following 6-OHDA lesions of the nucleus accumbens septi and corpus striatum. *Brain Res.*, **94**, 507-522.
- YANG, H.Y., ZHANG, J. & PIANTADOSI, L.A. (1996). Brain temperature alters hydroxyl radical production during cerebral ischemia/reperfusion in rats. *J. Cereb. Blood Flow Metab.*, **16**, 100-106.
- NG, J.F.R. & KLIPPEL, R.A. (1963). The rat brain. A Stereotaxic atlas of the forebrain and lower parts of the brain stem. New York, Robert E Krieger Publishing Co. Inc.
- VERTY, R. & LOGAN, B.J. (1990). Protection by MK801 and other drugs of methylenedioxymethamphetamine (MDMA) neurotoxicity in rats and mice. *Eur. J. Pharmacol.*, **183**, 451-452.
- ONARDI, E.T.K. & AZMITIA, E.C. (1994). MDMA (Ecstasy) inhibition of MAO type A and type B: Comparisons with fenfluramine and fluoxetine (Prozac). *Neuropsychopharmacology*, **19**, 231-238.
- MURRAY, T.K., ROSEBROUGH, N.J., FARR, A.L. & RANDALL, R.J. (1991). Protein measurement with the Folin phenol reagent. *J. Biol. Chem.*, **193**, 265-275.
- MALBERG, J.E., SABOL, K.E. & SEIDEN, L.S. (1996). Co-administration of MDMA with drugs that protect against MDMA neurotoxicity produces different effects on body temperature in the rat. *J. Pharmacol. Exp. Ther.*, **278**, 258-267.
- McMILLEN, B.A. (1983). CNS stimulants: two distinct mechanisms of action for amphetamine-like drugs. *Trends Pharmacol. Sci.*, **4**, 429-432.
- MOLLIVER, M.E., BERGER, U.V., MANOUNAS, L.A., MOLLIVER, D.L., O'HEARN, E. & WILSON, M.A. (1990). Neurotoxicity of MDMA and related compounds: anatomic studies. *Ann. N.Y. Acad. Sci.*, **600**, 640-661.
- MURRAY, T.K., WILLIAMS, J.L., MISRA, A., COLADO, M.I. & GREEN, A.R. (1996). The spin trap reagent PBN attenuates degeneration of 5-HT neurons in rat brain induced by p-chloroamphetamine but not fenfluramine. *Neuropharmacology*, **35**, 1615-1620.
- NASH, J.F. (1990). Ketanserin pretreatment attenuates MDMA-induced dopamine release in the striatum as measured by in vivo microdialysis. *Life Sci.*, **47**, 2401-2408.
- NASH, J.F. & BRODKIN, J. (1991). Microdialysis studies on 3,4-methylenedioxymethamphetamine-induced dopamine release: effect of dopamine uptake inhibitors. *J. Pharmacol. Exp. Ther.*, **259**, 820-825.
- O'HEARN, E., BATTAGLIA, G., DE SOUZA, E.B., KUJAR, M.J. & MOLLIVER, M.E. (1988). Methylenedioxymethamphetamine (MDA) and methylenedioxymethamphetamine (MDMA) cause ablation of serotonergic nerve terminals in forebrain: immunocytochemical evidence. *J. Neurosci.*, **8**, 2788-2803.
- RAITERI, M., CERRITO, F., CERVONI, A.M. & LEVI, G. (1979). Dopamine can be released by two mechanisms differentially affected by the dopamine transport inhibitor nomifensine. *J. Pharmacol. Exp. Ther.*, **208**, 195-202.
- SCHMIDT, C.J. (1987). Neurotoxicity of the psychedelic amphetamine, methylenedioxymethamphetamine. *J. Pharmacol. Exp. Ther.*, **240**, 1-7.
- SCHMIDT, C.J., ABBATE, G.M., BLACK, C.K. & TAYLOR, V.L. (1990a). Selective 5-hydroxytryptamine₂ receptor antagonists protect against the neurotoxicity of methylenedioxymethamphetamine in rats. *J. Pharmacol. Exp. Ther.*, **255**, 478-483.
- SCHMIDT, C.J., BLACK, C.K. & TAYLOR, V.L. (1990b). Antagonism of the neurotoxicity due to a single administration of methylenedioxymethamphetamine. *Eur. J. Pharmacol.*, **181**, 59-70.
- SCHMIDT, C.J., BLACK, C.K. & TAYLOR, V.L. (1991). L-DOPA potentiation of the serotonergic deficits due to a single administration of 3,4-methylenedioxymethamphetamine to rats. *Eur. J. Pharmacol.*, **203**, 41-49.
- SCHMIDT, C.J. & KEHNE, J.H. (1990). Neurotoxicity of MDMA: neurochemical effects. *Ann. N.Y. Acad. Sci.*, **600**, 665-681.
- SCHMIDT, C.J., RITTER, J.K., SONSALLA, P.K., HANSON, G.R. & GIBB, J.W. (1987). Role of dopamine in the neurotoxic effects of methamphetamine. *J. Pharmacol. Exp. Ther.*, **233**, 539-544.
- SCHMIDT, C.J., SULLIVAN, C.K. & FADAYEL, G.N. (1994). Blockade of striatal 5-hydroxytryptamine₂ receptors reduced the increase in extracellular concentrations of dopamine produced by the amphetamine analogue 3,4-methylenedioxymethamphetamine. *J. Neurochem.*, **62**, 1382-1389.
- SCHMIDT, C.J., WU, L. & LOVENBERG, W. (1986). Methylenedioxymethamphetamine: a potentially neurotoxic amphetamine analogue. *Eur. J. Pharmacol.*, **124**, 175-178.
- SHARKEY, J., McBEAN, D.E. & KELLY, P.A.T. (1991). Alterations in hippocampal function following repeated exposure to the amphetamine derivative methylenedioxymethamphetamine ('Ecstasy'). *Psychopharmacology*, **105**, 113-118.
- SPANOS, L.J. & YAMAMOTO, B.K. (1989). Acute and subchronic effects of methylenedioxymethamphetamine [(±)MDMA] on locomotion and serotonin syndrome behavior in the rat. *Pharmacol. Biochem. Behav.*, **32**, 835-840.
- STEELE, T.D., MCCANN, U.D. & RICAURTE, G.A. (1994). 3,4-methylenedioxymethamphetamine (MDMA, 'Ecstasy'); pharmacology and toxicology in animals and humans. *Addiction*, **89**, 539-551.
- STEELE, T.D., NICHOLAS, D.E. & YIM, G.K.W. (1987). Stereochemical effects of 3,4-methylenedioxymethamphetamine (MDMA) and related amphetamine derivatives on inhibition of uptake of [³H]-monoamines into synaptosomes from different regions of rat brain. *Biochem. Pharmacol.*, **36**, 2297-2303.

① No strong evidence for central DA role in mechanism of neurotox

uptake inhibitors block tox

SMT2 plays a role

analog toxicity correlated w/ DA release

② Neurotoxic metabolite $\leftarrow$ (should form from $\pm$ isomer)

- form in neuron? $\leftarrow$ may not be detectable

- enzymatically? $\leftarrow$ should be inhibitable

- non-enzymatically?

- form outside neuron? $\leftarrow$ should be detectable

- ~~the~~ maybe cause oxidative damage throughout?